dilatation. The lungs showed severe pulmonary oedema and there were serous transudates in both pleural cavities and in the peritoneal cavity. The right renal pedicle was free from disease, and the right kidney (84 g.), which was somewhat larger than usual, was embedded in very fibrous perirenal fat containing an apparently encapsulated tumour (diameter 3.5 cm.) showing foci of haemorrhage and necrosis. The lower pole of the kidney was distorted by this mass. At the upper pole the capsule stripped easily, leaving a smooth speckled surface, but over the lower pole adjacent to the tumour it was densely adherent. The cortex was a little broader than normal and showed accentuation of its markings and prominence of the glomeruli. The medulla was congested and the intrarenal vessels were not unduly prominent. The pelvis was normal, but the upper part of the right ureter, although embedded in the sclerotic perirenal tissue, was patent. There was an irregular haemorrhagic speckling of the lower part of the right lobe of the liver overlying the right kidney. No other significant abnormality was seen.

Histology.—The renal capsule was somewhat thickened and there was a mild diffuse interstitial fibrosis, although the renal architecture showed no gross distortion. Glomeruli were present in the usual numbers, but most showed fibrinoid and haemorrhagic necrosis of isolated tufts (Fig. 2). The tubules showed minimal dilatation and were lined with a flattened epithelium which showed foci containing hyaline droplets. Occasional tubules contained hyaline casts. Many of the arterioles and a few of the interlobular arteries showed severe fibrinoid necrosis; the arcuate arteries were normal. These appearances are characteristic of acute radiation nephritis as described by Russell (1953). In the liver the foci of dilated blood-filled sinuses were thought to be the late effects of hepatic irradiation.

Discussion

The first course of radiotherapy was directed to the left renal area. However, the metastasis was in the area of the right kidney. This required to be included in the area of treatment of the second course, and this course would probably be responsible for the radiation nephritis. The latent period before the onset of symptoms of nephritis was 128 days, although in retrospect it appears likely that the gain in weight in January, 1958, was due to fluid retention. This latent period is appreciably shorter than that encountered in the cases recorded in young adults (Luxton, 1953), and in this respect the present case is similar to the three cases described in young children by Zuelzer *et al.* (1950) except for the extremely rapid onset of uraemia and the short duration of the terminal illness.

The present case showed a severe acute haemolytic anaemia with well-marked anisocytosis, polychromasia,

poikilocytosis, and abundant "burr" cells in the peripheral blood. This blood picture is similar to that described in children by Gasser *et al.* (1955) and Macfarlane (1958) in association with the azotaemia of acute renal disease. The association of this blood picture with acute radiation nephritis has not been described previously, but in this case it is considered to be merely another manifestation of the fulminating uraemia.

My thanks are due to Mr. W. M. Dennison for permission to publish the clinical findings in this case; to Dr. T. M. Young for information on the radiotherapy administered; to Professor D. F. Cappel and Dr. A. M. MacDonald for advice and criticism; to Mr. W. Mason for the photographs; and to Miss M. Malcolm for technical assistance.

REFERENCES

- Gasser, C., Gautier, E., Steck, A., Siebenmann, R. E., and Oechslin, R. (1955). *Schweiz. med. Wschr.*, 85, 905.
Luxton, R. W. (1953). *Quart. J. Med.*, 22, 215.
Macfarlane, P. S. (1958). *Arch. Dis. Childh.*, 33, 181.
Russell, H. (1953). *Edinb. med. J.*, 60, 474.
Zuelzer, W. W., Palmer, H. D., and Newton, W. A., jun. (1950). *Amer. J. Path.*, 26, 1019.

GENERALIZED VACCINIA IN INFANCY

A REPORT OF FOUR CASES

BY

W. H. GALLOWAY, M.B., F.R.C.P.Ed., D.C.H.

Consultant Paediatrician

AND

L. M. MacBEAN, M.B., Ch.B., D.C.H.

Senior House Officer, Aberdeen Special Hospitals Group
and the Department of Child Health, University of
Aberdeen

Cutaneous complications of smallpox vaccination include generalized vaccinia, eczema vaccinatum, accidental vaccination, progressive vaccinia, and the erythema multiforme syndrome (Burgoon and Burgoon, 1956).

The term "generalized vaccinia" should be restricted to cases in which the primary site heals normally but where vaccinal lesions erupt on the body between the sixth and fourteenth days after vaccination. These develop in the same way as the primary, but because of the development of antibodies they progress at an accelerated rate and heal at the same time as the primary, without scarring (Barbero *et al.*, 1955).

Case Reports

Case 1.—A healthy male infant was vaccinated at 5 months of age and admitted to hospital 16 days later with generalized vaccinia. Death ensued a few hours later. A report of the case has already been published (Galloway and MacBean, 1956). No electrophoretic or antibody studies were carried out.

Case 2.—A healthy male infant was vaccinated at 3 months of age and admitted to hospital seven days later with generalized vaccinia. His mother had not been vaccinated; his father had. There were no other children. On examination he was pale, alert, irritable, and ill-looking. His temperature was 101° F. (38.3° C.). Over the left deltoid region was a large black scab about 1 in. (2.5 cm.) in diameter surrounded by a wide area of inflammation and induration. There were a few minute vesicles round the edge of the scab and a small secondary one ½ in. (1.3 cm.) from the primary site. Scattered vesicles were found on the left side of trunk, right axilla, both forearms, palms of both hands, sole of the left foot, and the left temporal region. Some were umbilicated. On admission intramuscular penicillin was given, also 4 ml. (500 mg.) of ordinary

FIG. 2.—Kidney showing interstitial fibrosis, fibrinoid change in glomeruli and arterioles, with minimal tubular dilatation. (Mallory. $\times 80$.)

gamma-globulin. Next morning fresh vesicles appeared over his face, scalp, napkin area, and feet. The following day 6 ml. (500 mg.) of hyperimmune vaccinia gamma-globulin was given, after which no further vesicles erupted. Penicillin was stopped and one of the tetracyclines started. The lesions went through the usual stages from vesicles to crusts, and healed at the same time as the primary, which was about the size of a normal "take." Electrophoresis of serum on admission showed no gamma-globulin fraction. One week after the administration of both ordinary and hyperimmune gamma-globulin electrophoretic study revealed a normal amount of gamma-globulin. Vaccinia virus was isolated from vesicular fluid.

Case 3.—A healthy male infant was vaccinated at 2 months of age and admitted to hospital nine days later because of vesicles on the face, scalp, arms, and legs. Both parents had been vaccinated in infancy. His brother had been vaccinated successfully. On examination the infant did not look particularly ill. The primary site was 1 in. (2.5 cm.) in diameter, and was moist and pustular. The lesions on the face and scalp were pustular; those on legs and hands were vesicular. On admission 6 ml. (500 mg.) of hyperimmune vaccinia gamma-globulin was given intramuscularly, and tetracycline therapy was begun. No fresh vesicles appeared, and he made an uneventful recovery. Electrophoresis of serum on admission showed gamma-globulin in concentration only slightly less than normal. Vaccinia virus was isolated from skin lesions.

Case 4.—A healthy male infant was 7 weeks old when vaccinated. He was admitted to hospital nine days later with generalized vaccinia. Both parents had been successfully vaccinated in infancy. There were no other children. On examination he was irritable but not particularly ill. The primary site was a crusted lesion 1 in. (2.5 cm.) in diameter, with surrounding vesicles and a wide area of inflammation. He had scattered rashes on the legs, feet, trunk, and back of the head, some of which were vesicular and some pustular. On admission 4 ml. (333 mg.) of hyperimmune vaccinia gamma-globulin was given intramuscularly, and tetracycline therapy was started. No fresh vesicles appeared, and the infant made an uncomplicated recovery. Electrophoresis of serum on admission showed gamma-globulin in concentration slightly less than normal. Vaccinia virus was isolated from skin lesions.

Serum Neutralizing Antibodies (Professor A. MacDonald).—Case 2 developed no neutralizing antibodies, and the only time that antibodies were detected was after the infant had had hyperimmune vaccinia gamma-globulin. Cases 3 and 4 developed vaccinia antibodies apparently in a normal manner.

Treatment

If the essential defect is in antibody formation it would seem rational to treat such patients as soon as possible with any substance rich in antibody content, so that passive immunity might be conferred. Keidan *et al.* (1953) treated his patient with blood and serum from a recently vaccinated donor, and followed this up with gamma-globulin. This patient had been vaccinated about one month previously, so that a considerable time had elapsed between onset of symptoms and treatment. In the cases here reported treatment was given within 48 hours of admission, and this may have had an important bearing on the successful outcome of the illness.

Kempe *et al.* (1956) reported on the use of hyperimmune vaccinia gamma-globulin in serious skin complications due to haematogenous spread following vaccination. This globulin was produced from blood obtained from donors who had been successfully vaccinated four to eight weeks previously. The antibody content was thought to be threefold that of gamma-globulin, and 5 ml. of the substance represented the antibodies of 1 pint (570 ml.) of donated blood. He suggested a dosage of 0.6 ml. per kg. body weight. Our cases had a slightly higher dose than this. Burgoon and Burgoon (1956) thought that the hyperimmune

vaccinia gamma-globulin might need to be repeated, and recommended a broad-spectrum antibiotic.

After admission two of our patients developed fresh lesions despite one of them having had 5 ml. of gamma-globulin. No fresh lesions developed in any patient after administration of hyperimmune vaccinia gamma-globulin. The temperature settled quickly, and the vesicles went through the usual stages, finally healing at about the same time as the primary. Most of the lesions healed without scarring except for a few which had been secondarily infected. The primary vaccination in all three healed completely, leaving a scar very little bigger than a normal "take."

In the fatal case the primary vaccination showed little sign of healing, involved the whole thickness of skin, and was approximately 5 cm. in diameter. This last case might indicate a lack of cutaneous immunity similar to the case of generalized vaccinia reported by Evans (1937). It is interesting to note that Evans suggested that such cases should be treated with serum from a recently vaccinated person.

Several of the cases reported in the literature had a long illness with a fatal outcome. None of our three patients who recovered was very ill, and it is felt that the prompt administration of hyperimmune vaccinia gamma-globulin prevented a serious illness from developing.

Discussion

The incidence of generalized vaccinia has been variously reported as 1 in 100,000 (Jubb, 1943) to 1 in 25,000 for infants vaccinated in the first year of life (Laurance *et al.*, 1952). The latter figure would seem to include all the various cutaneous complications of smallpox vaccination grouped under the heading of generalized vaccinia. Orlandini *et al.* (1955) have shown that the gamma-globulin level drops steeply from birth to the age of 1 month, when it is at its lowest level. It remains at this low level till the age of 3 months, after which it gradually rises to reach the adult level at the age of 2 years. Serum electrophoresis was not done in Case 1, but in Case 2 gamma-globulin was absent, and in Cases 3 and 4 it was present in slightly less than normal concentration.

Hayles *et al.* (1954) reported a successful uncomplicated vaccination in a patient with agammaglobulinaemia, and Apt (1953) knew of 24 patients with agammaglobulinaemia who were vaccinated without complication. The case reported by Keidan *et al.* (1953) had no neutralizing antibodies and lack gamma-globulin, while that of Barbero *et al.* (1955) had normal amounts of gamma-globulin although no neutralizing antibodies. Janeway and Gitlin (1957) state that it seems probable that the vulnerability of the young infant with a very low gamma-globulin level may be not so much the result of the low globulin level as that the low level reflects a lag in development of the capacity to form antibody gamma-globulins in response to infection. In this respect one patient (Case 2) was readmitted at the age of 8 months with meningococcal meningitis when electrophoresis showed a reasonable level of gamma-globulin. He made a successful recovery without the administration of gamma-globulin. He was seen at the age of 15 months, when his serum showed gamma-globulin in normal concentration.

The age of vaccination may be significant. Three of the patients here reported were vaccinated at an age when their gamma-globulin level would be at its lowest, as they were aged 3 months or under. In the one fatality (Case 1) the patient was 5 months old, when theoretically his level should have been higher than the younger infants. No electrophoretic studies were carried out on his serum. A section of spleen in this case showed absence of germinal follicles, as has been found in cases of agammaglobulinaemia (Somers, 1957). Recently a younger brother of this patient has been admitted to the Royal Hospital for Sick Children with widespread pneumonia. In view of the family history

electrophoresis of serum in the younger brother was carried out, and the result showed him to be lacking completely in gamma-globulin. It would seem reasonable, therefore, to presume that Case 1 likewise lacked gamma-globulin, and was probably an example of congenital agammaglobulinaemia. The other cases here reported were probably examples of "physiological transient hypogammaglobulinaemia of infancy" (Janeway and Gitlin, 1957). These workers quote Orlandini's figures for normal gamma-globulin levels in the first two years of life and suggest that the initial fall and subsequent rise to normal represents a physiological state rather than a disease.

Janeway and Gitlin also comment on the possible relationship between agammaglobulinaemia and progressive vaccinia. They think it possible that there are gradations of agammaglobulinaemia and that infants with complete agammaglobulinaemia, or even transient hypogammaglobulinaemia, who are vaccinated in early infancy may be overwhelmed because other mechanisms of resistance are less well developed. In their paper they quote L. Pillemer as stating that normal levels of serum complement and properdin are found in congenital agammaglobulinaemia.

The mothers of three of the infants (including the one who died) had been vaccinated in infancy or early childhood, and so presumably had very little antibody to pass on to their infants. Janeway and Gitlin (1957) stated that it is not so much the quantity of gamma-globulin that the infant receives from its mother as the quality—that is, its content of specific antibodies.

All our patients were male, and it is interesting to note that all cases of agammaglobulinaemia have been male except for Keidan's case.

The method of vaccination in three of the infants was by a $\frac{1}{4}$ -in. (6-mm.) linear scratch, and the other by two $\frac{1}{4}$ -in. (6-mm.) scratches at right angles to each other. Two had been vaccinated at a clinic and two by their family doctor. The lymph for three was from the same supplier but of different batch numbers, while the fourth was done by lymph from a different supplier. Several other infants had been vaccinated by these methods, the same batches of lymph being used without ill effects.

Summary and Conclusions

Four cases of generalized vaccinia are reported, with one death. The role of agammaglobulinaemia and antibody formation is discussed. The treatment with hyperimmune vaccinia gamma-globulin is described.

It is felt that generalized vaccinia occurs in some infants because of a failure of their resistance mechanism. Whether or not the failure to form gamma-globulin and/or antibodies is the essential defect, it would appear necessary that infants developing any of the serious cutaneous complications of smallpox vaccination should be treated as soon as possible with hyperimmune vaccinia gamma-globulin and a wide-spectrum antibiotic.

We thank Professor John Craig for his advice in compiling this paper, Dr. T. E. Anderson and Dr. Alan Lyell for permission to publish Cases 3 and 4, and Dr. H. Thom for carrying out the electrophoretic studies.

REFERENCES

- Apt, L. (1953). *Yearbook of Paediatrics*, p. 67. Year Book Publ., Chicago.
 Barbero, G. J., Gray, A., Scott, T. F. McN., and Kempe, C. H. (1955). *Pediatrics*, 16, 609.
 Burgoon, C. F., and Burgoon, J. S. (1956). *Ped. Clin. N. Amer.*, p. 521.
 Evans, P. R. (1937). *Arch. Dis. Childh.*, 12, 333.
 Galloway, W. H., and MacBean, L. M. (1956). *Brit. med. J.*, 2, 1102.
 Hayles, A. B., Stickler, G. B., and McKenzie, B. F. (1954). *Pediatrics*, 14, 449.
 Janeway, C. A., and Gitlin, D. (1957). *Advanc. Pediat.*, 9, 98, 109, 112.
 Jubb, A. A. (1943). *Brit. med. J.*, 1, 91.
 Keidan, S. E., McCarthy, K., and Haworth, J. C. (1953). *Arch. Dis. Childh.*, 28, 110.
 Kempe, C. H., Berge, T. O., and England, B. (1956). *Pediatrics*, 18, 177.
 Laurance, B., Cunliffe, A. C., and Dudgeon, J. A. (1952). *Arch. Dis. Childh.*, 27, 482.
 Orlandini, T. O., Sass-Kortsak, A., and Ebbs, J. H. (1955). *Pediatrics*, 16, 575.
 Somers, K. (1957). *Arch. Dis. Childh.*, 32, 220.

Medical Memoranda

Case of Progressive Vaccinia

A case of progressive vaccinia which responded well to hyperimmune vaccinia gamma-globulin is described below.

CASE REPORT

A West African infant aged 10 months was admitted to the South Western Hospital on March 25 with a generalized vaccinia eruption. He was first seen by his doctor on the previous day, when the lesions were reported to be pustular. The infant had been in hospital on two occasions with dietetic troubles, and on the second occasion was treated for ichthyosis. Primary vaccination was performed at a clinic on March 15. His mother stated that the rash appeared all over the body on March 22 and she noticed a swelling "under the left arm."

On admission his temperature was 102.8° F. (39.3° C.), pulse 150, and respirations 50. Apart from the skin, physical examination revealed nothing abnormal. The lungs were clear.

There was a raw granulating area 1 by $\frac{1}{4}$ in. (2.5 by 1.25 cm.) over the left deltoid, presumably the site of vaccination. Extending both upwards and downwards from this point on the outer surface was a profuse crop of umbilicated vesicles, circular in shape and quite superficial. Small crops of lesions were scattered over the rest of the body, including the face and scalp, in situations which the child could reach, and very possibly due to auto-inoculation. There were only two or three vesicles on the back. The colour of the lesions was difficult to gauge owing to the deep pigmentation. The axillary glands were enlarged, especially on the left side. Fresh lesions appeared daily up to March 28, especially on the face; the conjunctivae and oropharynx remained clear; and the granulating area round the site of vaccination extended to the shoulder and elbow on the external surface. The temperature remained raised, varying between 101 and 103° F. (38.3 and 39.4° C.), the appetite was poor and the infant obviously toxic.

On March 29 blood was withdrawn for antibody estimation and 1 g. of hyperimmune vaccinia gamma-globulin was given intramuscularly. This dose was repeated 24 hours later. The effect was dramatic: no fresh lesions appeared; scabs formed and separated quickly, leaving shallow foveate scars. 27 hours after the first injection the temperature fell to 98.8° F. (37.1° C.), and thereafter fluctuated between 98 and 99° F. (36.7 and 37.2° C.) until April 1, when it remained normal until discharge. The child's general condition improved rapidly and the appetite returned. By April 10 all scabs were separated and the granulating area was soundly healed. There was still some irritation, and gloves were necessary.

From admission until March 20 he was given 250,000 units of penicillin, six-hourly, without effect. This was supplemented on March 27 with sulphonamides with similar lack of response.

Dr. F. O. MacCallum, of the Virus Reference Laboratory, reported that the serum contained 640 mg. of gamma-globulin per 100 ml., which he stated was a normal level. A heavy growth of vaccinia virus was obtained from the scabs. The child was discharged on April 10.

COMMENT

The normal level of gamma-globulin, in spite of the fact that lesions were continuing to appear and the general condition of the child to deteriorate, followed by rapid response to hyperimmune gamma-globulin, suggests that this is an instance of dysgammaglobulinaemia as described