of working. But it is felt that the experience of other practitioners may be similar.

To guard against the intrusion of the human factor the doctor must first of all know himself and know his weaknesses. Secondly, he must enjoy his work, and have adequate spare time and leisure. Thirdly, he must not have a larger list than he is capable of dealing with. Fourthly, he must organize the practice for his own and his patients' benefit, even if this means making rules which may seem petty and tiresome to the patients themselves. In short, he must have an efficient mental discipline; and herein the fundamental difficulty of general practice lies. If his mental discipline is too rigid he is bound to lose some personal contact with his patients; if his personal contact is too strong, on the other hand, he cannot exert efficient mental discipline. A book could be written on the subject.

Summary

A record has been made of delays of diagnosis, following stated criteria, over a period of two years. Observations are made on these cases. In 5 out of 10 cases a human factor was thought to have played a major part in the delay. This factor is defined and discussed.

I wish to thank my partner, Dr. J. F. Mitchell, and Dr. T. A. Richards for their help and encouragement, and all those consultants of the Northampton General Hospital to whom most of the cases mentioned finally came for treatment and advice.

INFECTED CORK CLOSURES AND THE APPARENT SURVIVAL OF ORGANISMS IN ANTISEPTIC SOLUTIONS

BY

KEVIN ANDERSON, M.D.

AND

RUTH KEYNES, B.Sc.

From the Division of Bacteriology, Institute of Medical and Veterinary Science, Adelaide, South Australia

The dangers of infected cork closures have been repeatedly emphasized since the original description by Lowbury (1951). Antiseptic solutions are thought to be contaminated by contact with the cork during shaking and by being poured through the infected neck of the bottle. Since the organisms are frequently killed by the antiseptic when exposed *in vitro*, it has always seemed that such contamination must needs be intermittent, and that when the container is left undisturbed the contents should be sterile. A specimen of cetrimide sent to these laboratories for examination showed such persistent and repeated contamination, even when left undisturbed, that it gave support to the original comment that "organisms were growing in the solution." It was therefore decided to investigate the mechanisms of contamination which might lead to the apparent survival of organisms in an otherwise lethal fluid.

Examination of Infected Cetrimide Solution

The specimen consisted of half a gallon (2.2 litres) of 1% aqueous cetrimide contained in a glass flagon and sealed with a bark cork $1\frac{1}{2}$ in. (4 cm.) in diameter. On arrival in the laboratory the fluid level was below the bottle neck, which was dry. The cork was also dry, fissured but other-

wise intact. 0.1 ml. of the solution was placed in 100 ml. of nutrient broth and incubated for 18 hours. A segment of the cork, excised with a sterile scalpel, was cultured in a similar fashion. Both specimens yielded a heavy growth of a motile Gram-negative rod with the biochemical reactions of a paracolon bacillus. A pour-plate made at this time showed approximately 760 organisms per ml. of cetrimide solution. The container was left undisturbed at room temperature for seven days and the plate count was repeated. Samples were taken by removing the cork and passing a sterile pipette into the solution. Care was taken not to touch the neck of the bottle or the cork. The count on this occasion was 57×10^3 viable organisms per ml.

Experiment 1. Sensitivity of Infecting Organism to Cetrimide *in Vitro*.—Preliminary tests, using the tube titration method and pour-plates incorporating varying concentrations of fresh cetrimide, showed that the organism was consistently inhibited by a dilution of 1:1,000 cetrimide. The bacterioid concentration was determined by adding 1-ml. quantities of an overnight broth culture to 9 ml. of varying concentrations of aqueous cetrimide. After five minutes' exposure the inoculum was centrifuged down, washed twice with saline to remove residual cetrimide, and either placed in 100 ml. of nutrient broth or diluted for viable counting. With this technique, 1:1,000 cetrimide appeared bactericidal, in all but one of the experiments, for inocula of the order of $11,000 \times 10^6$ viable organisms per ml. The method used was more exacting than the random loop sample employed by Calman and Murray (1956), and this accounts for the greater concentration necessary to achieve a bactericidal effect in aqueous solution. The contaminating organism was none the less rapidly killed when exposed *in vitro* to one-tenth of the original concentration of cetrimide.

Experiment 2. To Exclude Growth Factors in Cetrimide Derived from Cork.—In view of the experimental findings of Nelson (1942), the cetrimide solution was examined for the presence of growth factors which might allow either survival or multiplication of contaminants. A portion of the solution was Seitz-filtered to remove organisms and examined by two-dimensional chromatography, using as solvent 80% alcohol and phenol saturated with water. The sample, which failed to reduce Benedict's solution, showed no hexoses, pentoses, or amino-acids. A cork extract, prepared by autoclaving 5 g. of cork in 100 ml. of distilled water, was similarly negative.

Experiment 3. To Exclude Physical Protection of Organisms by Cork Extracts.—10 ml. of the cork extract prepared in Experiment 2 was incorporated in 100 ml. of 1% aqueous cetrimide. This solution was used to repeat the sensitivity tests performed in Experiment 1. No protection of the inoculum could be demonstrated under these conditions, and the bactericidal figure remained unchanged. These findings agree with the experiments of Lowbury (1951), who concluded that his infected solutions contained insufficient solutes to protect added organisms.

Experiment 4. Self-sterilizing Properties of Contaminated Cetrimide Solutions.—Since the paracolon bacillus was rapidly killed by cetrimide *in vitro*, it was decided to investigate the behaviour of the contaminated solution after removal from its original container. 10 ml. of the contaminated cetrimide was carefully removed and transferred to a sterile screw-capped bottle. A time-clock was started and an immediate viable count performed (20 seconds elapsed between removal of the cork and the second tenfold dilution). The count showed 950×10^3 viable organisms per ml. Exactly five minutes after the transfer, volumes of 0.1 and 0.5 of the cetrimide were incorporated in pour-plates, and further samples were diluted for viable counting. No organisms were detected in these volumes of solution. This experiment was repeated on a number of occasions with identical results. It was therefore concluded that contaminated solutions of cetrimide are self-sterilizing within a short time of removal from their original container. Since, in this experiment, the solution had not come into contact with either the cork or

the bottle neck, it was thought that the physical act of removing the cork must in some way contaminate the solution at the moment of sampling.

Experiment 5. To Demonstrate the Mechanism of Contamination on Removing the Cork.—The neck of the flask was cut off with an emery wheel, leaving the handle intact (see Fig.). The infected cork was removed and the glass

sterilized by steaming. The apparatus was then held over an exposed agar plate, and the contaminated cork inserted and removed once. Inspection of the surface of the agar with a plate microscope showed countless particles of cork dust adhering to the surface. Subsequent incubation revealed numerous colonies of the infecting organism. The shape and size

of these colonies suggested that they had arisen from particles infected with numerous organisms rather than from single cells. The experiment was therefore repeated, holding the apparatus over 1 ml. of nutrient broth in a sterile container. This was shaken to disperse the organisms and a viable count performed. The contaminated broth contained 50×10^2 viable organisms per ml.

Experiment 6. Contamination of Other Antiseptic Solutions with Infected Cork.—The cork from the original flask was inserted, in turn, into flasks containing 10% aqueous dettol (new formula), 1:1,000 aqueous benzalkonium chloride ("zephiran"), and 1:2,000 aqueous chlorhexidine ("hibitane"). Paracolon bacilli could be readily recovered from all the solutions after one or two insertions of the infected cork. All solutions showed the same self-sterilizing properties as the original cetrimide when removed to another container.

Experiment 7. Histology of Infected Cork Closures.—A segment of the cork was excised and embedded in paraffin. Sections approximately 6μ in thickness were cut and stained by Gram's method. The main cellular structure of the cork was seen to be free of organisms. In certain parts of the section, however, the normal pattern was replaced by necrotic areas containing dark amorphous debris. In these fields large masses of Gram-negative bacilli could be seen, many lying free at the edge of the section. These areas seemed to represent the blackened fissures seen in many cork closures.

Discussion

The mechanism of contamination outlined above offers an explanation for the otherwise puzzling phenomenon of organisms recoverable from a bactericidal solution at any given moment of sampling. Although each manipulation of the infected cork releases a shower of infected cork dust on to the surface of the liquid, the actual numbers of organisms recovered will be inversely proportional to the time elapsing between removal of the cork and transfer of the sample to media which dilutes it beyond its bactericidal concentration. It is interesting to note that the viable count performed in Experiment 4, which was done in the shortest possible time after sampling, yielded the highest number of organisms recorded throughout this series of experiments.

In the routine culturing of such a contaminated solution, the sampling time is remarkably short and was found not to exceed 15 seconds in a number of experiments, even when manipulating a graduated pipette. It is therefore scarcely

surprising that positive cultures can be obtained repeatedly from solutions that have been contaminated in this way. Such results may give the impression that organisms are surviving in the antiseptic, and wide variations in the count due to different sampling times may even suggest that multiplication is taking place. Dust from infected corks can contaminate a variety of antiseptic solutions, and the phenomenon does not appear to be confined to cetrimide. Histology of the cork suggests that organisms colonize the fissures in large numbers, and that dry, friable closures are particularly likely to produce infection in this manner. Although the experiments of Nelson (1942) demonstrate that cork extracts can allow the multiplication of coliform organisms in distilled water, the nutrients concerned do not appear in solutions with cork closures in sufficient concentration to be detected by chromatography. Contaminating organisms, however, can clearly multiply in the interstices of corks, since it is scarcely credible that the large numbers seen in sections could have arisen from direct implantation only.

The self-sterilizing effect demonstrated in these experiments cannot excuse the continued widespread use of cork closures in bottles of all types containing solutions which may come into contact with an open wound. In addition to the mechanism described, contamination of a solution also occurs by direct contact with the cork or bottle neck, and only a few seconds may elapse between pouring the solution into a receiver and transferring it to the patient's skin.

Summary

A series of experiments which were carried out to investigate the apparent survival of a contaminating organism in an antiseptic solution are summarized.

A method of contamination is described which accounts for positive cultures obtained at any given moment of sampling. The apparent persistence of contaminants in a lethal solution is thereby explained.

The experimental contamination of other antiseptic solutions is demonstrated, and the dangers of cork closures are once again emphasized.

We are indebted to Miss Margaret Coles for chromatographic analysis of the solutions, to Mr. Martin Hansen for the histological sections, and to Mr. W. Nolan for the photograph. Our thanks are also due to Mr. Darwin, of I.C.I., for generous supplies of cetrimide and "hibitane" during this investigation.

REFERENCES

- Calman, R. M., and Murray, J. (1956). *Brit. med. J.*, 2, 200.
Lowbury, E. J. L. (1951). *Brit. J. Industr. Med.*, 8, 22.
Nelson, J. H. (1942). *J. Path. Bact.*, 54, 449.

Young Children in Hospital, by James Robertson (Tavistock Publications, Ltd., 4s. 6d. net), is a book reviewing the problem of the young child in hospital separated from his mother. It is based on a memorandum submitted on behalf of the Tavistock Clinic and the Tavistock Institute of Human Relations to the Committee on the Welfare of Children in Hospital appointed by the Central Health Services Council. It embodies experience the author obtained in the joint Child Development Research Unit of the Tavistock Clinic and Institute. Mr. Robertson has also made a documentary film, *Going to Hospital with Mother*, complementing the book. It describes how a mother of a young child can help when her child is admitted to hospital for an operation. The film is 16 mm. and runs for 45 minutes. It is available (hiring fee 25s.) from the Tavistock Institute of Human Relations, 2, Beaumont Street, London, W.1.

Billy Goes to Hospital is a "comic" intended for children about to enter hospital and is issued by the Department of Child Health, Guy's Hospital. Also from Guy's is a leaflet for the parents of such children, of which the cost of printing has been met by the hospital's Guild of Ex-patients and Friends. Its purpose is to help parents prepare children for this experience and so prevent consequent emotional upset.