Finally, comparison by sequential analysis showed that symptomatic remission and improvement in mucosal appearances were significantly more often induced by potent than by inert treatment.

My thanks are due to Glaxo Laboratories, who provided supplies of both potent and inert hydrocortisone for the trial; to Dr. P. Armitage, who designed the sequential analysis chart; and to Professor J. C. Goligher, who performed sigmoidoscopic examinations. I am grateful also to Professor J. C. Goligher, Mr. A. V. Pollock, Mr. G. Smiddy, Mr. E. T. McCartney, Dr. W. S. Suffern, Dr. M. Atkinson, and Mr. D. B. Feather, who referred patients for treatment.

REFERENCES

- Armitage, P. (1954). *Quart. J. Med.*, 23, 255.
 — (1957). *Biometrika*, 44, 9.
 Bross, I. (1952). *Biometrics*, 8, 188.
 Newton, D. R. L., and Tanner, J. M. (1956). *Brit. med. J.*, 2, 1096.
 Snell, E. S., and Armitage, P. (1957). *Lancet*, 1, 860.
 Truelove, S. C. (1956). *Brit. med. J.*, 2, 1267.
 — (1957). *Ibid.*, 1, 1437.

ANKYLOSING SPONDYLITIS

BY

F. DUDLEY HART, M.D., F.R.C.P.

Westminster Hospital, London

In the spring of 1958 a questionnaire was sent to 343 patients with ankylosing spondylitis who had been seen, and the majority treated, at Westminster Hospital since the war. Answers were received from 209; most of the rest were returned by the Post Office, the present address of those patients being unknown. Many of the patients had been seen and treated while in the armed Forces, and the difficulty in tracing many lay in this fact. In the vast majority of cases, however, where the questionnaire was sent to the correct address an answer was received within three weeks, demonstrating once again the high degree of co-operation found in our previous study with this group of patients (Hart, Robinson, Allchin, and MacLagan, 1949). Questions were related particularly to the effects of treatment as assessed by the patient himself and the ability to do his work and earn his own living.

Duration of Disease.—Of the 209 who returned their questionnaires, 23 had suffered from ankylosing spondylitis since before the onset of the second world war, in 58 the onset lay during 1940–4, in 75 during 1945–9, in 44 during 1950–4; in only 9 had it developed since that date.

Results

Work Capacity.—200 question sheets were returned fully completed. Of these, 21 were from women, 179 from men. This 8.5:1 ratio is an artificial one owing to the high number of Service men seen; it therefore does not reflect the true sex ratio in this disease. Of this number, 160 were working full time, and, if housewives, were doing their full duties in the home. Many stated that in addition to their ordinary duties they worked voluntarily in the evenings, and others did gardening and played games. 40 stated that they were incapable of full duties, but only 9 were incapable of any duties whatsoever. 19 were capable of modified duties and 12 of light or occasional duties. In addition the questionnaire had been returned by relatives of four patients, all male, who had died: one of amyloidosis, one of coronary thrombosis, one of carcinoma of the pancreas, and one of a neoplasm of uncertain nature. These figures—160 at full work, 19 at modified work, and only 21 capable of little or no duties—are typical of the group as a whole, and similar figures have been reported by Blumberg and Ragan (1956). Although working full time, many commented that the duties

in their particular post were not physically hard and some that the post had been selected because of their disability. In four instances a regular paid post had been given up for full-time independent work where self-imposed duties could be varied to avoid the stiffness and discomfort attendant on too little mobility at work. Once again the well-known fact became evident that a stiff spine alone does not cripple, though stiff hips and peripheral joint involvement may.

General Health.—Of 191 patients who answered questions relating to their general health, 36 stated it was excellent, 74 good, 71 "fairly good," and 10 poor. Asked specifically about indigestion, 11 complained of severe recurrent or chronic dyspepsia, 38 of more moderate but annoying symptoms. Radiological investigation in 37 cases had revealed peptic ulcers in 10. Of these 191 cases, therefore, 49, or roughly 25%, have dyspepsia to the point of being a definite and occasionally a major complaint. Although some symptoms were obviously related to or aggravated by therapy, this was clearly not so in all cases. This group of spondylitic dyspeptic patients remains a major therapeutic hazard, for almost all current forms of therapy may aggravate gastro-duodenal symptoms or cause complications. Of other disabilities, the next most common was iritis. Although no specific question was asked and although the eye was not mentioned on the questionnaire, 35 patients mentioned iridocyclitis when asked for general comments about their health. One was registered as a blind person from this disability. Three patients were under treatment for pulmonary tuberculosis; one had had a thoracoplasty. Vertigo was mentioned by 7, headaches by 4, anaemia by 4, and recurrent bronchitis by only 4, although 8 made major complaints regarding stiffness of the chest wall and inability to breathe deeply.

Analgesics.—To the question, "What medicines have you found to be most helpful for your spondylitis?" 43 answered none, 49 aspirin in some form, 54 compound codeine tablets, 56 phenylbutazone, and 9 steroid therapy. This last is a small group, as only a small minority had been offered steroid therapy. No other drug was mentioned more than once.

Radiotherapy

One of the main reasons for the follow-up was to learn what the patients thought of the long-term results of deep x-ray therapy and to find out what blood disorders had occurred, if any. The large majority of these patients were treated at Westminster Hospital by Dr. F. M. Allchin, Mr. Tom Prosser, and Dr. Kenneth Newton in the conservative dosage reported by us previously (Hart *et al.*, 1949); only a few treated elsewhere received a total dosage above 1,500 r to any part of the spine, and the majority received under 1,200 r. Of 143 patients who had received therapy to the spine, 114 (80%) stated they had received some benefit, 29 (20%) that they had not. Of those who stated that they had received benefit, 41 (28.7% of all patients treated with deep x rays) stated that some benefit still persisted, after follow-up periods varying from 2 months to 17 years since therapy was given; 27 of these 41 patients had received therapy more than four years previously. 30 patients had had symptomatic relief for six months or less, 18 for seven months to five years, 3 for periods over five years, the effect then completely wearing off. In 22 cases details of duration of relief were not adequately given.

While some of this improvement may be psychogenic and reflect the high optimism of the patients as a group, the stories given fall into fairly regular patterns, and suggest more than "therapeutic persuasion." In some cases the question of introduction of a natural remission has to be considered where for many years after therapy the condition appears to have remained largely or completely quiescent. On these figures it seems that 20% of irradiated spondylitics will derive no benefit whatsoever, 20% a temporary symptomatic improvement for a period of up to six months, the remaining 60% some degree of relief for periods varying from seven months to many years.

Discussion

With so many missing returns no conclusions could be drawn as regards mortality and morbidity, though the main reason for failure to return the questionnaire was the migratory nature of these patients and frequent change of addresses. Nevertheless, of the returns received 80% of the patients were at full work and only 5% capable of no work whatsoever of any sort, the remaining 15% being capable of modified, light, or occasional duties. These figures are the most encouraging we have in the rheumatism unit—far better than those for rheumatoid arthritis. 110 of 191 patients considered their general health good or excellent and only 10 poor.

As regards the most popular of the drugs used it is of interest that aspirin in some form, compound codeine tablets, and phenylbutazone were quoted in almost identical numbers, the last-named coming top in this poll by the narrow margin of two votes. It has been our impression in the unit for some time that, if tolerated, this drug is the single most useful one for spondylitics at all stages of their painful disease, and Bauer and Ropes (1956) found it more useful in this disease than in rheumatoid arthritis.

The effect of any drug, however, depends on its regular daily administration; its beneficial effects rapidly disappear on stopping it. In contrast, deep x rays appear, on this survey, to give some degree of lasting relief in many cases, for 32 of 114 patients (28%) obtained some amelioration of symptoms for over four years. Caution is necessary, however, in interpreting these figures, for a course of deep x-ray therapy, like a surgical operation, sticks in the memory as a landmark in the disease course, and subsequent improvement may falsely be attributed to it when, in fact, natural remission has occurred. Improvement, also, is as compared by the patient with pre-treatment symptoms which were severe enough to bring the patient to hospital. One may therefore be comparing symptoms of exacerbation with symptoms of remission, and the therapy given between the two may have been unrelated. A second point to emphasize is that it is impossible on the returns to tell whether symptomatic improvement was great or small; all that can be said is that some measure of symptomatic improvement occurred after therapy.

Wilkinson and Bywaters (1958) found that 43% of 200 irradiated patients required no regular analgesics within a year of completion of therapy, this figure dropping to 22% after a ten-year follow-up. These figures, obtained by clinicians from assessment of analgesic consumption, do not differ very greatly from those obtained by us from an assessment of our patients' own statements.

While I have hitherto been doubtful about the induction of natural remission in this disorder by this form of therapy, there is some indication on these figures that it may occur. The hazards of irradiation are well known, and have latterly received much attention; it is as well to record the reverse side of the coin.

Summary and Conclusions

Question sheets were sent to 343 patients with ankylosing spondylitis, and 209 answers were received. Of these returns, four were from relatives of deceased patients, and five were incomplete. Of the 200 fully completed returns, 179 were from men, 21 from women.

160 (80%) stated that they were working full time; 19 were capable of modified duties, 12 of light or occasional duties; and only 9 were permanently incapable of any work whatsoever.

Of 191 patients who answered questions relating to their general health, 36 stated that it was excellent, 74 good, 71 fairly good, and 10 poor. 11 complained of severe, 38 of moderate but annoying dyspepsia. X-ray examination in 37 cases revealed peptic ulcers in 10.

Of 143 patients given deep x-ray therapy to the spine, 114 (80%) derived some benefit; 41 (28.7%) claimed some measure of lasting relief, which still continues 2 months to 17 years after treatment. Of these 41 patients 27 were irradiated over four years previously. 30 claimed transient improvement lasting six months or less, 18 improvement for periods of seven months to five years, 3 for more than five years. 32 of 114 (28%) of these patients derived some measure of symptomatic relief for periods exceeding four years.

These results once again emphasize the relatively good prognosis as regards work capabilities in this group of patients and a reasonably satisfactory response to existing forms of therapy.

My thanks are due to Professor J. H. Kellgren and Dr. J. Sharp, of the Rheumatism Research Centre, Manchester University, for their helpful and constructive comments.

REFERENCES

- Bauer, W., and Ropes, M. W. (1956). *Practitioner*, 177, 431.
Blumberg, B., and Ragan, C. (1956). *Medicine (Baltimore)*, 35, 1.
Hart, F. D., Robinson, K. C., Allchin, F. M., and MacLagan, N. F. (1949). *Quart. J. Med.*, 18, 217.
Wilkinson, M., and Bywaters, E. G. L. (1958). *Ann. rheum. Dis.*, 17, 209.

VALUE OF AZACYCLONOL ON CHRONIC SCHIZOPHRENIC PATIENTS PRELIMINARY OBSERVATIONS

BY

SYLVIA REID, B.A., M.B., D.P.M.

Consultant Psychiatrist, Carlton Hayes Hospital,
Narborough, near Leicester

Azacyclonol hydrochloride ("frenquel") is the gamma isomer of pipradrol hydrochloride ("meratran"), the latter drug being used in this country as an antidepressant agent. Chemically, azacyclonol is α -(4-piperidyl) benzhydrol hydrochloride. Despite its structural resemblance to pipradrol, azacyclonol has been reported to possess different clinical properties inasmuch as it exerts a tranquillizing effect.

Both mescaline and L.S.D.25 have been reported to produce states resembling schizophrenia. Fabing (1955a, 1955b, 1955c), working in the United States, has reported a series of controlled experiments on "normal" volunteers who were each given L.S.D.25. This dose produced a characteristic psychosis with symptoms resembling schizophrenia. The experiment was then repeated, but some of the volunteers were given a course of azacyclonol prior to the administration of the L.S.D.25 and the other volunteers were given control tablets. It was observed that, although the effects of L.S.D.25 began as usual, the more acute symptoms were inhibited in the patients who had been pretreated with azacyclonol. The vivid hallucinatory experiences were absent and some of the group even succeeded in carrying out their normal duties. It would appear from this work that azacyclonol has some blocking effect in experimental psychosis or clinical conditions of disorientation.

The nature of this neurochemical effect is not yet fully understood.

Fabing and Hawkins (1955) have reported on clinical trials with azacyclonol in a series of acute and chronic schizophrenic patients. It appears that azacyclonol affords schizophrenic patients some amelioration of