BRITISH MEDICAL JOURNAL

LONDON SATURDAY FEBRUARY 15 1958

NEUROLOGY: PAST, PRESENT, AND FUTURE*

BY

Sir RUSSELL BRAIN, Bt., D.M., F.R.C.P., F.R.C.P.Ed.

Physician, London Hospital and Maida Vale Hospital

What is neurology? The subtitle of the journal *Brain* is "A Journal of Neurology," and the scope of that journal may therefore be taken as a useful index of the scope of neurology. Any volume of *Brain* contains articles on the physiology, anatomy, and biochemistry of the nervous system, on neuropathology, on the psychological and psychiatric aspects of nervous disorders, and on clinical neurology, including neurosurgery. In the most comprehensive sense, then, neurology comprises all aspects of the normal and abnormal functioning of the nervous system. In one important respect neurology differs from all other branches of physiology and of medicine: it is concerned with that part of the human organism which is most distinctively human, and it is therefore unique among all branches of medicine in the extent of its contributions to scientific knowledge beyond the scope of animal physiology, through its study of the neural basis of all the psychological and physiological functions of man.

The Century of Nosography

Before the nineteenth century, though many isolated neurological observations had been made from the earliest times, neurology as a science can hardly be said to have existed. The nineteenth century was pre-eminently a century of nosography—of the recognition of syndromes and diseases. It would take far too long to enumerate all those which the nineteenth-century clinicians and pathologists delineated in neurology; instead I will trace the development of thought about two which still present us with unsolved problems.

Disseminated Sclerosis

Disseminated sclerosis was first recognized as a pathological entity by Cruveilhier (1835) and by Carswell (1838). Cruveilhier noted also the association of the pathological lesions with paraplegia in one case. Limited clinical and pathological studies were made by German authors, but it was Charcot and his school who first presented a coherent clinical picture. Indeed, to read Charcot's (1877) lectures on disseminated sclerosis is to realize how comparatively little has been added to our clinical knowledge of the disease since his day. Though Charcot maintained that "the multiplication of nuclei and the concomitant hyperplasia of the reticulated fibres of the neuroglia constitute the initial, fundamental fact, and necessary antecedent," and that "the degenerative atrophy of the nerve elements is consecutive and secondary," the breakdown of the myelin had not escaped his notice, for he drew attention to bodies containing fatty droplets—"granular bodies" exactly

resembling "those which result from the disintegration of the medullary matter (myeline) in cases of nerve-section."

And then a century was to pass before our knowledge of the disease made a new and at present very tentative advance. Post-vaccinal encephalomyelitis was certainly not a new disease when an outbreak in the early 1920s attracted attention (Bastiaanse, 1925), and pathological studies of this and allied disorders following measles, and other exanthems, revealed acute inflammatory lesions of the nervous system both resembling and differing from disseminated sclerosis (Turnbull and McIntosh, 1926). This led experimental pathologists to study the production of acute demyelinating lesions in animals by the injection of brain tissue, a condition already recognized in man as a result of antirabic inoculation. Meanwhile other observations had suggested that some forms of demyelination, or failure of myelination, might be related to defects in the oligodendroglia, which stimulated research into the role of the glia in relation to the production of myelin. The present position of this research is well summarized in Lumsden's (1957) two chapters in "Modern Trends in Neurology" (second series) entitled "Some Aspects of the Chemistry of the Myelin and Sheath Cell Complex" and "Cell Structure and Cell Physiology in Relation to Myelin." Here we learn how biochemistry, tissue culture, the electron microscope, and electrophysiology are contributing to the fundamental knowledge of the nerve cell which we must possess if we are to understand the aetiology of disseminated sclerosis.

Parkinsonism

Now turn to another common nervous disease which exhibits a growth of knowledge parallel in time with that of disseminated sclerosis, but in some interesting respects different. In 1817 James Parkinson published his classical *Essay on the Shaking Palsy*. For a long time, however, this disease was confused with disseminated sclerosis, the differentiation being made in the mid-century by Charcot, amongst others, in the lectures from which I have already quoted. It was not until this century, however, that any light was thrown upon its pathology, chiefly owing to the discovery by Kinnier Wilson (1912) of the association between degeneration of the corpus striatum and symptoms resembling Parkinsonism. Lewy (1914) drew attention to the presence of cell degeneration in the putamen and globus pallidus and, later, in the substantia nigra. Almost exactly a century after Parkinson's publication epidemic encephalitis produced a new variety of Parkinsonism, and the relationship between the syndrome and lesions of the substantia nigra was confirmed. By this time it was

*The Frederick Price Lecture delivered at the Royal College of Physicians of Edinburgh on November 13, 1957.

recognized that Parkinsonism was merely a syndrome, resulting from a number of causes—cerebral atheroma, neurosyphilis, manganese and carbon monoxide poisoning amongst others—in addition to those already mentioned.

The association between the liver disease and the lenticular degeneration in Wilson's disease remained a puzzle until, quite recently, it was discovered that the fundamental disorder was an abnormality of copper metabolism, which is now being extensively studied. But why do the cells of the substantia nigra undergo degeneration in the spontaneous variety called paralysis agitans by Marshall Hall? In paralysis agitans we recognize one of the "presenile degenerations," a group which includes some of the most insidious and relentless nervous disorders, the presenile dementias, the progressive cerebellar degenerations, and motor neurone disease. Unless a lucky chance discloses unsuspected metabolic disturbances in patients suffering from these diseases, it is likely that we shall have to wait for more detailed knowledge of the metabolism of the nerve-cell before we can understand them.

While during the nineteenth century clinicians and pathologists, joined later by the bacteriologists, were establishing clinical syndromes and correlating them with their pathology and aetiology, during the second half of the century there was an increasing analysis of symptoms themselves, and the development of broad concepts of the functions of the nervous system. Examples of this are the discovery of the excitability of the brain by Fritsch and Hitzig (1870) and by Ferrier (1873), Jackson's observations on epilepsy and his ideas of the organization of function in the nervous system, Gowers's acute clinical observations, the large literature on aphasia, the concepts of localization and representation in the nervous system which emerged from all this work; and, just at the very end of the century, Babinski's (1896) account of his observations on the plantar reflex.

Contribution of the Twentieth Century

The contribution of the twentieth century to neurology has been so rich and varied that I can depict only the broad outlines. Fundamental to all subsequent work was the publication by Sherrington of *The Integrative Action of the Nervous System* in 1906. Sherrington's study of the reflex arc, the interaction between reflexes, inhibition and excitation, the synapse, a term which he had invented some years before, proprioceptors, decerebrate rigidity, and the physiological position and dominance of the brain opened for neurology a new world, the exploration of which, as we shall see, is still far from exhausted.

Between the beginning of the century and the first world war Head and Sir Gordon Holmes were advancing our knowledge of the neurology of sensation. Head's work on the peripheral nerves (Rivers and Head, 1908), though much of it has not stood the test of time, stimulated subsequent investigators, and I am sure he would have felt much interest in the recent ideas of the relation between quality of sensation and pattern of impulses in the peripheral nerves. We owe to this period also the work of Head and Thompson (1906) on sensation in the spinal cord, and Head and Holmes (1911) on the thalamic syndrome and sensory disturbances at the cortical level, and Sir Francis Walshe's (1914) application of Sherrington's ideas to the interpretation of the reflexes in paralysis of the lower limbs in man. The spinal injuries of the first world war stimulated still further the study of the reactions of the damaged spinal cord and bore fruit in the work of Head and Riddoch (1917), Riddoch (1917), and Walshe (1919). Gunshot wounds

of the brain gave Holmes, who had already contributed to our knowledge of the cerebellar degenerations, the opportunity for his masterly studies of the disordered physiology of the cerebellum in man (Holmes, 1917, 1922) and the cortical representation of vision, and the higher visual functions (Holmes, 1918a, 1918b; Holmes and Lister, 1916).

Neurosurgery

Meanwhile neurosurgery was growing up. In its modern form it begins with the pioneer work of Horsley, who laid the foundations of neurosurgery not only as a technique but also as a sphere of applied physiology, in which he was followed by Cushing and Foerster, amongst many others. Neurosurgery brought in its train advances in neuroradiology—especially ventriculography and pneumo-encephalography, which we owe to Dandy, and cerebral angiography, the invention of Moniz—which may be said to have introduced a new dimension into neurology. In the treatment of the gliomas neurosurgery has proved disappointing, and we must look to a greater understanding of both tumour formation and the biochemistry of neuroglia for an effective answer to this distressing disease.

The contribution of neurosurgery to neurological knowledge arises chiefly out of the increased accuracy of location of cerebral lesions which it, with its ancillary investigations, has made possible, and hence the differentiation of disorders previously merged in a general picture of increased intracranial pressure. Though Cushing and Bailey (1928) and others had drawn attention to the vascular malformations of the brain before angiography was introduced, it is only since then that we have been able to estimate the frequency of these lesions and to recognize their varied symptomatology. Our knowledge of intracranial aneurysms has followed a somewhat similar course. It is surprising that what was presumably a common lesion went so long unrecognized. But it was not until the work of Fearnside (1916) and Turnbull (1918) on their pathological aspects and, above all, Symonds's (1924-5) clinical and pathological contribution that we began to recognize subarachnoid haemorrhage. Even so, it was not until angiography became available that diagnosis could be in many cases accurate enough to raise the question of surgical treatment, the indications for which are still a matter of discussion. And it is largely to angiography that we owe the recognition of the symptomatology of atheroma of the internal carotid artery.

Electroencephalography

Electroencephalography was developing *pari passu* with these radiological techniques. Beginning with the discovery of Berger (1929) and Adrian and Matthews's (1934) pioneer applications of the method to neurophysiology, it was soon taken up in the clinical sphere by Gibbs, Gibbs, and Lennox (1937), Grey Walter (1938), and others. A quarter of a century's experience shows that in many patients suspected of an organic cerebral lesion or dysrhythmic disturbance it will add little to what can be learned by careful history-taking and clinical examination. Its chief diagnostic value lies in: (1) locating a lesion not otherwise localizable, or the confirmation of a localization otherwise in doubt; examples of this are the recognition and localization of a metastatic abscess or neoplasm in the brain, the presence of which might otherwise be unsuspected; (2) the demonstration of a dysrhythmia which confirms the suspicion of epilepsy made on clinical grounds; (3) the locating of an epileptogenic focus; and (4) the demonstration of diffuse degenerative or inflammatory processes, and even in some cases the detection of a type of dysrhythmia characteristic of the disease. It follows that, like all diagnostic aids, electroencephalography is of value only when the clinician knows what question to put to the machine, and how to interpret the answer it gives.

The uses of electroencephalography which I have enumerated have contributed greatly to the clarification of our understanding of epilepsy during the last twenty years. The idea that epilepsy should be regarded primarily as a symptom

is a very old one, expressed by Hughlings Jackson and stressed by Kinnier Wilson in his emphasis on "the epilepsies." For a long time, however, it seemed to be in conflict with the concept of idiopathic epilepsy. The effect of electroencephalography aided by neuroradiology has been greatly to extend our recognition of symptomatic or focal epilepsy by demonstrating foci of dysrhythmia or angiomas or areas of cerebral atrophy which elude clinical investigation. So it has proved possible more often to correlate the clinical features of an epileptic attack with an anatomical lesion; and here neurosurgery has contributed by making the cerebral cortex of the conscious epileptic patient available for electrical stimulation. Much, however, remains to be learned about the spread of focal epilepsy, the origin of generalized major epilepsy and petit mal, and the underlying basis of hereditary predisposition. The tendency to neural discharge must possess a biochemical basis, as the value of anticonvulsant drugs shows. The abnormal biochemical state may often be the result of a local structural lesion, but is it always so? The answer to this question may depend upon the neurophysiologists, who are now able to stimulate localized areas deep in the brain and study the resulting disturbances of cortical rhythm, which is but one application of electroencephalography to neurophysiology, about which I have more to say later.

Recent Developments in Neurophysiology

This brings me naturally to recent developments in neurophysiology. Encephalitis lethargica, epidemic in the years following 1916, showed a predilection for the basal structures of the cerebral hemispheres, the corpus striatum, the hypothalamus, and the upper part of the brain stem. Some patients, particularly children and adolescents, exhibited striking physiological and psychological aberrations which attracted much attention, including obesity and genital atrophy, polydipsia and polyuria, and impulsive, aggressive, and irresponsible behaviour. During the following decade physiologists were producing similar abnormalities in animals by means of lesions at the base of the brain.

Diabetes insipidus, obesity, and gonadal atrophy were all attributed to localized lesions of the hypothalamus by Bailey and Bremer (1921), Camus and Roussy (1922a, 1922b), Smith (1927), and others, and this initiated a vast amount of work devoted to the study of the mutual relationships of the hypothalamus and the pituitary. Bard (1929) produced what he called "sham rage" in the cat by a lesion between the frontal lobes and the anterior part of the hypothalamus, and showed that this behaviour could be abolished by a section posterior to the hypothalamus. At about the same time the production of sleep-like states in animals by lesions deep in the central part of the brain stem provided an interpretation of the sleep disturbances of encephalitis lethargica. This work, thirty years ago, marks the beginning of three important developments in neurology: the correlation between behaviour disturbances and focal lesions of the brain; the study of the influence of the nervous system upon the endocrine system and so on the biochemistry of behaviour; and, lastly, the investigation of the central nervous regulation of the state of consciousness.

Now let us jump thirty years and see where these studies stand to-day. There have been numerous investigations of behaviour disturbances in animals produced in various ways, either by the ablation of comparatively large areas, such as the temporal lobe of Klüver and Bucy (1939), or by the destruction of more localized areas of cortex. By means of deeply implanted electrodes it has been possible to study the effects of stimulating localized deep structures. The application of electroencephalography to physiology has introduced a functional method of tracing the connexions of areas of the nervous system by making it possible to detect the remote responses to a stimulus. This is one of the methods which have been used to elucidate the relationship of the central reticular formation of the brain stem to consciousness, the effect of the activity of that region

being correlated with the electrical state of the cerebral cortex and the behaviour of the animal.

The relationship between the hypothalamus and the endocrine system has been illuminated by such work as Verney's (1957) on the hypothalamic regulation of the renal excretion of water, and that of Harris and others, who have shown that individual nuclei of the diencephalon may act as chemoreceptors to particular hormones. Adrenaline, for example, acts upon the central reticular formation; and Harris, Michael, and Scott (1957) have shown that the implantation of very small amounts of stilboestrol into the hypothalamus of ovariectomized cats, but not elsewhere, results in full sexual behaviour in the absence of or with minimal development of the vaginal epithelium and uterus. Thus the central site of action of ovarian hormones in producing active sexual behaviour would seem to be in some part of the posterior hypothalamus or upper mid-brain.

Fifty years ago Sherrington was writing about excitation and inhibition, and the synapse. To-day the newest neurophysiological techniques are concentrated upon precisely these topics. Eccles (1957) has shown that it is possible to penetrate into the interior of the single nerve cell and analyse the transactions which are taking place across its membrane. Thus he has been able to differentiate the functions of the dendrites and the axon hillock and to show that discharge depends on depolarization and inhibition on hyperpolarization, stimulation and inhibition summing in their effect on the cell, as Sherrington foresaw. Hodgkin, Huxley, and Keynes have also studied the physico-chemical behaviour of the neurone. Thus we are beginning to learn about the role of chemical transmitters at the synapse in the central nervous system, and the chemical basis of excitation and inhibition. Similar studies are showing what goes on at the myoneural junction.

Clinical Neurology To-day

Where does clinical neurology stand to-day? People sometimes speak as though nosography in neurology was almost exhausted, but this is quite untrue. The discovery of new diseases or syndromes, and equally the elaboration of the clinical pictures of existing ones, are partly a matter of seeing differences in what has hitherto been thought to be homogeneous. I have already shown how new diagnostic methods have contributed to this process in the case of epilepsy, cerebral angioma, intracranial aneurysm, and internal carotid atheroma. Radiography also played an important part in the recognition of the neurological manifestations of cervical spondylosis. Compression of the median nerve in the carpal tunnel is an example of a syndrome which seems to have become much commoner since its recognition shortly after the war. New diseases and the increased incidence of old diseases, together with changes in the age distribution of the population, have also altered the clinical landscape of neurology, producing, for example, the carcinomatous neuromyopathies and new infections such as acute inclusion encephalitis and benign myalgic encephalomyelitis, the "Royal Free disease." The neurological aspects of hypertension and atheroma are of growing importance. Involvement of the nervous system and muscles in collagen diseases is a new field now being explored, and, among the metabolic disorders, porphyria. Electromyography is throwing new light upon disorders of the lower motor neurone, the myoneural junction, and the muscles.

It might have been thought that after a century of study the main disorders of cerebral function would have been elucidated. Yet since the beginning of the war considerable additions have been made to our knowledge of the functions of the frontal lobe, the symptomatology of lesions of the temporal lobe and of the minor hemisphere, the agnosias and disorders of the body image, and the central regulation of consciousness. But the greater our knowledge of the effects of lesions of different parts of the

brain, the more apparent becomes our need for a clarification of our thought on the meaning of localization and representation in the nervous system. Such considerations are not to be regarded as merely the preoccupation of a few neurological schoolmen: they are fundamental to our understanding of how the nervous system works, and hence to the interpretation not only of symptoms but also of the observations of the neurophysiologist.

The Contribution of Neurology to Psychiatry

Hughlings Jackson (1932), writing of insanity, said in 1881: "We require a rational generalization so wide as to show on the physical side relations of diseases of the mind, which are for physicians nothing but diseases of the highest centres, to all other diseases of the nervous system. We have to find some fundamental principle under which things so superficially different as the diseases empirically named hemiplegia, aphasia, acute mania, chorea, melancholia, permanent dementia, coma, etc., can be methodically classified." If nearly eighty years later we ask how far Hughlings Jackson's requirement has been or is likely to be fulfilled, we should discover what is the essential neurological contribution to psychiatry, and what are its limitations.

Both neurologists and psychiatrists recognize in practice the validity of four kinds of aetiological factor in their work. (1) There are the causes which produce disorders of function by acting physically upon the nervous system. This wide field ranges from pathophysiology, which requires normal physiology for its interpretation, through neurochemistry to the visible changes studied by the neuropathologist. It is essentially the study of the nervous system as a physical object, behaving in health and disease according to the laws of the physics and chemistry of its constituent parts. (2) There is the study of the effects of the bodily changes of the kind which I have just described upon the mind. (3) We may approach the mind-brain relationship from the other side and observe the effects of mental changes upon the body, where we enter the field of what is commonly called psychosomatic medicine. (4) Finally, we may limit our study of causation to the psychological sphere, by interpreting psychological disorders in psychological terms—that is, asking how the patient's present mental state has been caused by his previous mental states. This aspect of psychiatry comprises psychopathology and psychotherapy in its various forms.

Although I began with physical and ended with psychological aetiology, this does not mean that I regard any one form as more fundamental or valuable than the rest. I might equally well have arranged them in the reverse order, with the same qualification. Nor does the fact that we must recognize these four varieties of aetiology in "neuropsychiatry" absolve us from the duty of applying the most rigid criteria of validity to the arguments which may be used at any aetiological level. Thus, the recognition that symptoms may be explained as psychogenic does not imply the acceptance of one particular theory of psychopathology or technique of psychotherapy. And the practitioner who gives due weight to the factors in all these aetiological fields is unlikely to accept too readily a psychological explanation for severe mental illness, merely because at the moment we are unable to explain it on physical grounds. And, since this classification is pragmatic, one form of causation does not exclude another. It is in the sphere of the first two categories of causation, which deal with neuropathology and with the influence of physical changes in the nervous system upon the mind, that we may expect to find the main contribution of neurology to psychiatry. I need not discuss long-familiar disorders like the dementias and mental defect, nor the psychological importance of the ever-increasing knowledge of the aphasia, agnosias, and disorders of the body image.

The actual or potential contributions of neurology to psychiatry could be illustrated from examples of the progress of neurology, some of which I have already given—the numerous symptoms of disturbances of the temporal lobe

which may lead to disorders of awareness of the self and the external world, and may produce distortions of memory or states of fear, sorrow, disgust, loneliness, or depression (Williams, 1956). Whitty and Lewin's (1957) observations on cingulotomy throw an interesting light on the sense of reality. Sleep, dreams, illusions, and hallucinations are all subjects as appropriately studied by the neurologist as by the psychiatrist. Electric convulsion therapy and leucotomy act neurologically.

Thus the simple concept that the mind is dependent upon the brain has during recent years been increasingly refined and defined till now we know much, though still far too little, about the part played by the different areas of the brain in the regulation of consciousness, memory, emotional mood, and the integration of the personality.

If we compare the psychological disorders which can be explained in terms of psychological causation with those which we have reason to think are the result of physical changes in the nervous system, we note certain points of difference between them. The first relates to their neural basis. We know extremely little about the organization of memory in the brain, but it is surely clear that the nervous organization underlying one memory can differ from that underlying another only in respect of subtle differences in the arrangement of the finest nervous impulses, or perhaps synaptic resistances, over wide areas of the brain. It must follow, therefore, that no physical method which can be applied to the brain, whether surgical, electrical, or pharmacological, is capable of dealing with an individual past experience as such.

Interpretation of Mental Disorder

To put the matter in another way, when the mind is disordered as a result of disorder of the nervous system the neurological disorder is describable, actually or at any rate potentially, in physical terms, that is to say, in terms of the biochemistry of nerve cells or the gross structural changes familiar to the neuropathologist. When, however, the mind is disordered as a result of the reactions of the individual to his previous experiences, no such explanation in terms of biochemistry or morbid anatomy is possible. No doubt the neurological explanation is that certain nervous impulses are taking one pathway rather than another, *but the pathways themselves can be distinguished as healthy or unhealthy only in terms of the psychological states with which they are associated.* Here we reach the limitation of Hughlings Jackson's hopes for the neurological interpretation of mental disorder. Hemiplegia, delirium, and coma are states of body or mind which can be attributed to physical disorders of the nervous system, and it may well be that mania and melancholia, as Jackson suggested, and possibly ultimately schizophrenia, will be similarly explained. If so, the causes of these disorders could be described in purely physical terms and without reference to their symptoms. Psychogenic disorders belong to a different category of causation, which cannot be adequately described without using psychological terms or bringing in their symptoms.

This, of course, does not exclude the possibility that more than one type of causation may be operative in the same patient at the same time. For example, an individual's abnormal psychological reactions to past experiences may be explicable in psychological terms, but they may also be the result of an inherited predisposition depending upon the structure and functions of his nervous system. Conversely, a person with an inherited predisposition to recurrent attacks of depression, which presumably depends upon some abnormality in the behaviour of his nervous system, may have an attack of depression precipitated by some psychological stress. And it is still very difficult in the present state of our knowledge to evaluate the relative importance of physical and psychological factors in the causation of some psychological disorders, of which the obsessional states are a good example.

Hence a patient suffering from a psychological disorder needs to be considered from both neurological and psycho-

logical standpoints. In practice, in spite of the need for this dual approach and the frequent interaction of the two modes of causation, in many patients the physical or the psychological element will clearly predominate, and this will determine the method of treatment.

It seems to me to follow, from what I have said, that even if by neuropsychiatry is meant only the field common to both medical neurology and psychiatry, it is impossible for a single practitioner to be expert in all parts of it. But, even so, it does not follow that the present division between the two subjects is made in the right place. This is the outcome partly of historical and partly of social considerations, the need for the segregation of some of the mentally ill having in the past isolated in mental hospitals patients suffering from certain neurological disorders and from mental illnesses the causation of which was, and is still, unknown. Neurology grew up in equal isolation from the mental hospital, and the study of the psychological effects of cerebral lesions, however interesting to neurologists and psychologists, seemed to have little to contribute to the understanding of the major psychiatric problems of schizophrenia, cyclothymia, and the depressive states. The theory and practice of analytical psychopathology and psychotherapy were developed largely in isolation from both neurology and mental hospital psychiatry. Fortunately these barriers are in many places being steadily broken down.

Neurology and General Medicine

The Committee on Neurology of the Royal College of Physicians of London made the following statement: "Neurology remains a part, though an important part, of general medicine and cannot without danger to its own vitality allow itself to become separated from the larger whole. Much of the most valuable neurological material is to be found in the general medical and surgical wards and out-patient departments. Many neurological disorders are the direct result of disease of the cardiovascular system. Others are as directly related to general or local infective processes or to disorders of nutrition or metabolism." The Committee drew from this the conclusion that "to be able to play his part fully in the recognition and management of such cases, it is essential that the neurologist should base his specialized knowledge on a foundation of sound training and experience in general medicine and on a general familiarity with allied specialties such as is obtainable in a large general hospital.

I think that the facts to which the Committee drew attention imply not merely that the neurologist should "base his specialized knowledge on a foundation of sound training and experience in general medicine," but also that there should be a close continuing relationship between neurology and general medicine in the work of our hospitals. No branch of medicine can live on its past achievements, and, although the sphere of applied anatomy and physiology which has yielded such a rich harvest to neurology in the past is by no means exhausted, it may well prove to be subject to a law of diminishing returns, while fresh fields await cultivation. Nothing can take the place of the special neurological hospital. It has an essential part to play in the training of neurologists and provides a unique forum for the interchange of ideas and a continuing stimulus to the thought of the neurologist already in practice. Moreover, it is also the most suitable place for the expansion of neurology into some of the new fields which recent scientific developments, especially in biochemistry and electronics, are opening.

In the special hospital, however, there is a relative lack of certain types of clinical material, and it is inevitable that there should be some neurological disorders which are to be found chiefly, if not exclusively, in relation to general medical diseases and not therefore at a special hospital, while much of the neurology of childhood is to be found in children's hospitals. An increasing amount of research work is being done in these fields, and has already demonstrated its value both to neurology and to medicine. This can be developed as it should only if the work of the neurologist in

the general hospital is given the fullest possible scope. It would be a mistake to suppose that his place there can be supplied by the general physician. The scope of neurology as outlined in this lecture is surely a sufficient general refutation of that. The recognition of new syndromes requires expert knowledge to differentiate them from other clinically similar conditions or out of what has hitherto been regarded as a homogeneous group of disorders. Similarly the assessment of the indications for therapeutic procedures must in difficult cases depend upon a knowledge of the multiplicity of factors involved which only the expert possesses.

The contribution of the specialist to diagnosis is equally important. We are all of us apt to regard the field of work of others as possibly less complex than our own, and there exist physicians who, though well enough aware of the complex effects of disturbances of suprarenal function, regard the diagnosis of paraplegia as something which presents no great difficulty. The neurologist who is an active participant in the work of his general hospital will do far more than merely help his colleagues with the diagnosis of cases which seem to them difficult; he will educate them in the difficulty of cases which seem to them easy. He also has a valuable contribution to make to medical teaching, since neurology is a subject which lends itself especially to the education of the student in the importance of accurate history-taking, careful clinical examination, and the logical processes by which a diagnosis is reached.

Conclusions

What I have said well illustrates, if indeed it needed to be illustrated, the rich contribution which specialization has made to medicine. To-day, far from knowing more and more about less and less, the specialist, like every one of his colleagues in medicine, needs to know more and more about more and more. Hughlings Jackson said in 1887: "As scientific medical research goes on, there is greater specialization of investigation, just as, in the development of society, there is that continually increasing specialization, called division of labour. This being so, all the more need is there that there should be greater integration, just as along with division of labour there is need for co-operation of labourers." Failing the appearance of a race of medical supermen who can take all medical knowledge for their province, this integration can be accomplished only by the exchange of information between workers in different fields, and this is most easily achieved by enabling them to meet and work together under the same roof.

It is true that the rapid expansion of scientific knowledge in medicine calls for a review of our methods of organization, and possibly, even, for new tools of thought, but, judging by the results so far achieved, there is nothing to suggest that the human mind is anywhere near the limits of its ability to deal with the challenge created by its own productivity. If we ask what part neurology is to play in this process the answer begins to emerge from our survey of the scope of the subject. Neurology plays a part in physiology, psychology, psychiatry, pathology, bacteriology, medicine, surgery, and no doubt other disciplines, and all of these in their turn have a contribution to make to neurology. What, then, is the role of the neurological physician? The fact that he alone is commonly called a neurologist surely reflects his key position in relation to neurology.

The process of integration, to which I have just alluded, is more likely to be successful if there is an integrator; and the neurologist, though he cannot possibly be an expert in all these fields, is the person most fitted by training, experience, and opportunity to be a nodal point in the relations of neurology. Even so, as the College report recognized, within the work of the

neurologist there is room for some differentiation. "In the future, as has always been the case in the past, some neurologists will be particularly interested in those aspects of neurology which are relatively remote from psychiatry. Others will find their chief interest in those neurological disorders in which the psychiatric aspect is prominent. These latter should be in a position to pursue their interests unfettered, for they have a most valuable contribution to make to the study of neurology as a whole, and, especially at a time when psychiatry is turning more and more to physical methods of treatment, they have similarly a valuable contribution to make to the study of psychiatry." The Committee also saw the need for the general physician, who combines an interest and training in neurology with his general medicine. The growing relationship between neurology and general medicine might well now be more widely recognized in the training of the neurologist, as it has been in the arrangement recently entered into between the Maida Vale Hospital and the London Hospital for two alternating senior registrarships tenable at both.

I have been speaking of definitions: let me end with a parable. The Pigeon in *Alice in Wonderland* insisted that Alice must be a serpent because she ate eggs. "Little girls," said Alice, 'eat eggs quite as much as serpents do, you know.' 'I don't believe it,' said the Pigeon, 'but if they do, they're a kind of serpent, that's all I can say.'"

The logic of the Pigeon sometimes finds expression to-day. Many kinds of doctors rightly see patients with nervous disorders, but the future of neurology in this country still depends upon its ability to attract, train, and provide satisfying work for neurologists of equal calibre to those to whom we owe the achievements of this branch of medicine and science in the past.

REFERENCES

- Adrian, E. D., and Matthews, B. H. C. (1934). *Brain*, 57, 355.
 Babinski, J. (1896). *C.R. Soc. Biol. (Paris)*, 3, 207.
 Bailey, P., and Bremer, F. (1921). *Arch. Intern. Med.*, 28, 773.
 Bard, P. (1929). *Arch. Neurol. Psychiat. (Chicago)*, 22, 230.
 Bastiaanse, F. S. van B. (1925). *Bull. Acad. Méd.*, 94, 815.
 Berger, H. (1929). *Arch. Psychiat. Nervenkr.*, 87, 527.
 Camus, J., and Roussy, G. (1922a). *J. méd. franç.*, 11, 332.
 — (1922b). *Rev. neurol.*, 38, 622.
 Carswell, R. (1838). *Pathological Anatomy*. London.
 Charcot, J. M. (1877). *Lectures on the Diseases of the Nervous System*. New Sydenham Society, London.
 Cruveilhier, J. (1835). *Anatomie Pathologique du Corps Humain*. Paris.
 Cushing, H., and Bailey, P. (1928). *Tumors Arising from the Blood-vessels of the Brain*. Springfield, Ill.
 Eccles, J. C. (1957). *The Physiology of Nerve Cells*. Johns Hopkins Press, Baltimore.
 Fearnside, E. G. (1916). *Brain*, 39, 224.
 Ferrier, D. (1873). West Riding Lunatic Asylum Medical Report, 3, 30.
 Fritsch, G., and Hitzig, E. (1870). *Arch. Anat. Physiol. (Lpz.)*, 37, 300.
 Gibbs, F. A., Gibbs, E. L., and Lennox, W. G. (1937). *Brain*, 60, 377.
 Harris, G. W., Michael, R. P., and Scott, P. Ciba Foundation Symposium on Neurological Basis of Behaviour, 1957 (in the press).
 Head, H., and Holmes, G. (1911). *Brain*, 34, 102.
 — and Riddoch, G. (1917). *Ibid.*, 40, 188.
 — and Thompson, T. (1906). *Ibid.*, 29, 537.
 Holmes, G. (1917). *Ibid.*, 40, 461.
 — (1918a). *Brit. J. Ophthalm.*, 2, 353.
 — (1918b). *Ibid.*, 2, 449, 506.
 — (1922). *Lancet*, 1, 1177, 1231; 2, 59, 111.
 — and Lister, W. T. (1916). *Brain*, 39, 34.
 Jackson, J. Hughlings (1932). *Selected Writings*. Hodder and Stoughton, London.
 Klüver, H., and Bucy, P. C. (1939). *Arch. Neurol. Psychiat. (Chicago)*, 42, 979.
 Lewy, F. H. (1914). *Dtsch. Z. Nervenheilk.*, 50, 50.
 Lumsden, C. E. (1957). In *Modern Trends in Neurology* (second series), edited by Denis Williams. Butterworth, London.
 Riddoch, G. (1917). *Brain*, 40, 264.
 Rivers, W. H. R., and Head, H. (1908). *Ibid.*, 31, 323.
 Sherrington, C. S. (1906). *The Integrative Action of the Nervous System*. New York.
 Smith, P. E. (1927). *J. Amer. med. Ass.*, 88, 158.
 Symonds, C. P. (1924-5). *Quart. J. Med.*, 18, 93.
 Turnbull, H. M. (1918). *Brain*, 41, 50.
 — and McIntosh, J. (1926). *Brit. J. exp. Path.*, 7, 181.
 Verney, E. B. (1957). *Lancet*, 2, 1237, 1295.
 Walshe, F. M. R. (1914). *Brain*, 37, 269.
 — (1919). *Ibid.*, 42, 1.
 Walter, W. G. (1938). *J. Neurol. Psychiat.*, 1, 359.
 Whitty, C. W. M., and Lewin, W. (1957). *Brain*, 80, 72.
 Williams, D. (1956). *Ibid.*, 79, 29.
 Wilson, S. A. K. (1912). *Ibid.*, 34, 295.

THE CHOLECYSTOGRAM AND THE CLINICIAN*

BY

J. DUDFIELD ROSE, F.R.C.S.

Senior Surgeon, Newcastle General Hospital; Associate Surgeon, Royal Victoria Infirmary, Newcastle upon Tyne

The first step to any form of treatment, whether medical or surgical, is to make an accurate diagnosis. The selection of patients for operation is made on clinical grounds and on the results of appropriate technical investigations. The usual investigation to clarify a diagnosis of disorder of the biliary tract is cholecystography. Great reliance has been placed on this type of investigation despite the fact that it gives a positive diagnosis in less than half the cases of suspected disorder of the biliary tract submitted to it. In an analysis of four separate series of 100 consecutive cholecystographic examinations from three hospitals and my own personal series, the numbers of cases in which no abnormality was reported were 56, 55, 66, and 66. This seemed to be a high incidence, particularly as all the patients had definite symptoms of a disorder of the biliary tract which in many cases was severe. The problem that arises is, What is the diagnosis of these patients, and how may the diagnosis be made?

The "classical cholecystogram" will demonstrate the presence or absence of a gall-bladder shadow, it may demonstrate the presence of gall-stones, it may demonstrate morphological abnormalities of the gall-bladder, it may show a subnormal power of concentration, it will demonstrate whether the gall-bladder contracts or not after a fatty meal. It will not give exact information on the evacuation of the gall-bladder, nor will it in any way give information concerning the various motor dysfunctions of the biliary tract. There are also morphological abnormalities which are not demonstrated by the classical method.

Serial Cholecystography

Experience with the technique of per-operative manometry and radiology (Mallet-Guy and Rose, 1956) over the past three years has demonstrated quite decisively the incidence and importance of disturbances of the motor function of the biliary tract—biliary dyskinesia. A dyskinesia was demonstrated in no fewer than 126 cases out of the first 200 consecutive cases investigated and treated during that time. In 38 of these 126 cases there was a dyskinesia in the absence of any morphological abnormality. This experience indicated the necessity for a more accurate pre-operative method of diagnosing dyskinesia. It was thought that cholecystography should be able to provide the answer to this problem. It was also obvious that cholecystography in its usual form did not supply the answer.

Boyden (1926), in his original work, studied the motor function of the gall-bladder by means of a series of cholecystographic pictures, from which he calculated the volume of the gall-bladder and drew a graph of its evacuation. Clinical radiologists abbreviated this method, and the present technique of cholecystography is to take control pictures before the ingestion of an opaque medium, 14 hours after it, and further radiographs at a variable time after a fatty meal. In some departments this refinement is omitted. This abbreviation destroyed in great part the value of

*Read to the Sections of Radiology and Surgery at the Annual Meeting of the British Medical Association, Newcastle upon Tyne, 1957.