Drug Treatment of Disease

DRUG TREATMENT OF NEURITIS

BY

J. W. ALDREN TURNER, D.M., F.R.C.P.*Neurologist, St. Bartholomew's Hospital*

Neuritis is a much misused term. Many of the conditions to which it is sometimes applied are in fact of mechanical origin. For instance, brachial neuritis may be used as a diagnostic label for pain in the arm accompanied by paraesthesiae and neurological signs when in fact the basic cause is compression or irritation of posterior roots by a protruding intervertebral disk or cervical spondylosis, pressure on the brachial plexus by a cervical rib or by drooping of the shoulder girdle in the axillary-inlet syndrome, or injury of a peripheral nerve, as by compression of the median nerve in the carpal tunnel or friction to the ulnar nerve at the elbow in association with cubitus valgus or osteoarthritis of the elbow. Similarly, in the lower limb, sciatic neuritis and femoral neuritis probably do not exist, the majority of cases of sciatica and anterior crural pain being due to mechanical causes, such as protrusion of intervertebral disk or other lesions compressing the nerves or posterior roots. In such cases drug treatment can only be of very limited value and consists of suitable analgesics. The essential treatment for the condition—immobilization, physiotherapy of various types, manipulation, or surgical treatment—varies with the aetiology. The administration of vitamin-B preparations for such conditions has no rational place in treatment.

Multiple peripheral neuritis or polyneuritis is to some extent also a misnomer, but in many of such cases there are changes in the parenchyma of the peripheral nerves. There are many possible causes of polyneuritis, and in some of them drug treatment is of value.

Deficiency of Vitamin B₁

Deficiency of aneurine (vitamin B₁) in relation to the carbohydrate intake may cause peripheral neuritis as a result of disturbance of carbohydrate metabolism in the nervous system at the pyruvate stage. In such cases the amount of pyruvate in the blood is increased, and this can, if necessary, be shown by a pyruvate tolerance curve. Actual deficiency of aneurine in the diet is rare in the United Kingdom, but defective absorption may occur in cases of steatorrhoea, long-continued diarrhoea, extensive resection of the small intestine, and long-continued fever; and a deficiency peripheral neuritis may occur in these conditions. The peripheral neuritis of alcoholism is to a large extent due to aneurine deficiency in relation to the carbohydrate intake.

In such cases treatment should be started at the earliest opportunity before degenerative changes have occurred in the peripheral nerves and when the lesion is still a biochemical one. The first essential is large doses of aneurine by injection, and, as in many cases there is also a deficiency of other elements of the vitamin-B complex, and of vitamin C, these should be given in addition, together with a normal diet. A suitable preparation for starting treatment is "parentrovite" high potency intramuscular injection, each injection of 7 ml. containing aneurine hydrochloride 250 mg., nicotinamide 160 mg., riboflavin 4 mg., pyridoxine 50 mg., and ascorbic acid 500 mg. After daily injections have been given for a week oral treatment should be continued with tab. aneurin. co. fort. (B.N.F.), which contains aneurine hydrochloride 5 mg., riboflavin 2 mg., nicotinamide 20 mg., and pyridoxine 2 mg., a tablet being taken four times a day.

Acute Infectious, Toxic, or Febrile Polyneuritis (Guillain-Barré Syndrome)

Although the aetiology of this type of polyneuritis is uncertain, the evidence suggests that it may be allergic in nature. For this reason it has been treated with cortisone or prednisone. The natural history of the disease is for rapid deterioration to take place over the course of a few days up to two to three weeks; if the patient survives the acute stage, rapid recovery of function usually occurs. This type of natural history makes it difficult to estimate the value of cortisone therapy, but published cases and personal experience certainly suggest that it is of value in stopping deterioration and possibly in hastening recovery. Cortisone should be given as early as possible, and a suitable dosage for an adult is 300 mg. (in divided doses) daily for two days, 200 mg. for two days, and then 100 mg. daily for ten days, the dosage then being gradually tailed off. Prednisone is less likely to cause retention of salt, and for this reason is usually the preferable preparation to give (5 mg. of prednisone being equivalent to 25 mg. of cortisone).

Lead "Neuritis"

This condition is probably the result of a disturbance of resynthesis of phosphocreatine in the muscles affected rather than a disorder of the peripheral nerves. In the last few years the treatment of acute lead-poisoning, such as lead encephalopathy, has been greatly improved by the introduction of the chelating agent calcium disodium ethylenediamine tetra-acetic acid, or sodium calciumedetate ("calcium disodium versenate"), and though reports of its use in lead "neuritis" are scanty it would seem to be the rational method of removing the lead from the body in this condition. When this chelate is administered to a patient with lead-poisoning, the lead replaces calcium in the chelate, the resulting compound being excreted in the urine. The chelated lead compound does not ionize and so there is little danger of causing symptoms of acute lead-poisoning, which was always a risk with the older methods of removing lead from the body.

The preparation is supplied in 5-ml. ampoules containing 1 g. in a 20% aqueous solution; this should be diluted in 250–500 ml. of injection of sodium chloride B.P. or injection of dextrose B.P. and administered by intravenous drip over the course of an hour. For adults this dose should be given twice daily for five days; then there should be a rest for two days, followed by a further five days of treatment. Calcium disodium versenate can be given by mouth, but it is not yet clear how effective this method of administration is in removing lead from the body.

Diabetic Neuritis

The most essential treatment in the peripheral neuritis associated with diabetes mellitus is the adequate control of the disease by diet and in most cases insulin. Patients suffering much pain may sometimes have this alleviated by injections of vitamin B₁₂, a suitable dose being 1,000 µg. daily for a week. In the minority of cases where there is burning pain in the feet, calcium pantothenate 50 mg. three times a day by mouth is sometimes helpful.

Other Types of Polyneuritis

In the rare cases of polyneuritis due to ingestion of arsenic, dimercaprol (B.A.L.) is the best treatment. This