fluoresced when pretreated with six of eight sera, but nuclei in skin from a patient with D.L.E. (Case 4) showed fluorescence when pretreated with only three of the eight sera.

In view of the apparent diminution of affinity of L.E. serum for nuclei in L.E. tissue, skin obtained from another patient (Case 7) was examined. Pretreatment of this material with autologous serum gave an equivocal result, as shown in Fig. 7 on the Special Plate. Although diffuse fluorescence of epidermal cells was seen, selective fluorescence of nuclei did not occur. This finding cannot be explained by failure of autologous tissue and serum to react, as nuclear fluorescence was observed when tissue from Case 4 was pretreated with autologous serum. It was established by two experiments that the serum from Case 7 had not lost its potency. The serum was used to pretreat leucocytes from a healthy individual, and nuclear fluorescence was observed when the serum was diluted to 1:4,000. Undiluted serum produced nuclear fluorescence when applied to the cells for one minute.

The necessity for using fresh material in this type of investigation has been stressed by Holborow *et al.* (1957). The difficulty of obtaining regular supplies of tissue by biopsy is considerable, and it was decided to examine cells obtained by scraping the buccal mucosa. These could be taken as representative of fixed tissue cells as distinct from the circulating leucocytes. However, the nuclei of squamous cells from healthy individuals or patients with D.L.E. exposed to D.L.E. serum failed to show fluorescence on subsequent treatment with fluorescein-conjugated A.H.G. serum (Special Plate, Figs. 9 and 10). When, however, the time of exposure to D.L.E. serum was extended to 1½ hours nuclear fluorescence was observed in cells which had disrupted. As there was also evidence of diffuse bright fluorescence in structures thought to be of cytoplasmic origin, no conclusion can be drawn from these experiments. Further investigation with specific counter-stains for nuclear material is planned.

Discussion

These results confirm the findings of other workers that the nuclei of cells exposed to serum from patients suffering from D.L.E. adsorb a globulin detectable by application of fluorescein-conjugated A.H.G. serum. This property of L.E. serum is not apparently influenced by administration of steroid hormones.

It has been shown that the phenomenon is readily demonstrable in unfixed blood films. This technique, which avoids the need for preparing films from "buffy coats," would be useful in any large-scale investigation of the phenomenon in other diseases.

The observation that nuclear fluorescence is not present in tissue from cases of D.L.E. exposed directly to fluorescent A.H.G. serum is in agreement with the findings of Mellors *et al.* (1957) in the D.L.E. kidney, and supports the hypothesis that absorption of L.E. globulin on nuclei occurs only *in vitro*. The fact that tissues and leucocytes derived from patients with D.L.E. appear to be less susceptible to uptake of globulin from L.E. serum may be explained by the relatively small number of patients available for study, but the anomalous behaviour of the serum from Case 7 toward nuclei in autologous skin and the nuclei in the skin of another patient is of interest. Recent experiments, which are still incomplete, suggest that the time of incubation of tissue and L.E. serum may have a considerable influence on the intensity of nuclear fluorescence and that individual sera and tissues may differ appreciably in the rate of reaction.

The results of experiments with squamous epithelium suggest that intact tissue cells resist invasion of the specific globulin in L.E. serum. It has been noted in examination of thick sections of skin that nuclear fluorescence may be apparent in the superficial layer, but not in deeper layers. Leucocytes, however, seem to be extremely susceptible to the specific globulin in L.E. serum, and this susceptibility may well be associated with their natural property of phagocytosis.

Summary

With the use of Coons's fluorescent antibody technique it has been confirmed that sera from patients suffering from D.L.E. contain a globulin which combines with nuclei of tissue cells and leucocytes. In a study of sera from eight cases of D.L.E. it was observed that nuclei of leucocytes from healthy individuals showed the greatest affinity for the specific globulin. No evidence of uptake of globulin in tissue from D.L.E. was found *in vivo*, but *in vitro* such tissue had less affinity for the specific globulin than tissue from healthy subjects. Nuclei of intact squamous epithelial cells pretreated with D.L.E. serum did not fluoresce on exposure to fluorescein-conjugated anti-human-globulin serum.

During the period of this investigation the Rheumatic Unit was in receipt of grants from the Nuffield Foundation, the Medical Research Council, and Boots Pure Drug Company Ltd.

We wish to thank Mrs. Betty Greenhill, A.I.M.L.T., and Mrs. Ruth Wallace for their skilled technical assistance.

REFERENCES

- Alexander, W. R. M. (1958). *Immunology*, 1, 217.
 Coons, A. H., and Kaplan, M. H. (1950). *J. exp. Med.*, 91, 1.
 Friou, G. J. (1957). Communication to Ninth International Congress of Rheumatic Diseases, Toronto.
 — (1958). *J. Immunol.*, 80, 476.
 — Finch, S. C., and Detre, K. D. (1957). *Fed. Proc.*, 16, 413.
 — (1958). *J. Immunol.*, 80, 324.
 Holborow, E. J., Weir, D. M., and Johnson, G. D. (1957). *Brit. med. J.*, 2, 732.
 Holman, H. R., and Kunkel, H. G. (1957). *Science*, 126, 162.
 Mellors, R. C., Ortega, L. G., and Holman, H. R. (1957). *J. exp. Med.*, 106, 191.

LIQUID PARAFFIN AS A CAUSE OF OIL ASPIRATION PNEUMONIA

BY

GILBERT FORBES, M.D., B.Sc., F.R.C.S.Ed.
F.R.F.P.S.

Reader in Charge

AND

ANDREW BRADLEY, M.A., LL.B., M.B., B.S.
Lecturer

Department of Forensic Medicine, University of Sheffield

[WITH SPECIAL PLATE]

Liquid paraffin has, for various reasons, been under suspicion at different times since it began to be used to any extent medicinally at the beginning of this century. It has been charged with preventing intestinal absorption of food in general, and of vitamin A and its precursor, carotene, in particular, and on this latter count there is some justification for the charge. Since liquid paraffin is a mixture of liquid hydrocarbons obtained from petroleum, the possibility that it might contain carcinogenic fractions has been investigated (Boyd and Doll, 1954). At one time there was a practice of instilling liquid paraffin into the peritoneal cavity with the avowed object of preventing adhesions. The result has been to produce, in a number of cases, an oil granuloma of the peritoneum (Bennett and Collins, 1952). Probably, however, the most common danger arising from the medicinal use of liquid paraffin is the production of oil aspiration pneumonia.

The pneumonia which results from aspiration of oils and fats has been dealt with extensively in American literature, but has not received the same attention in Britain, although its incidence in this country may not be any less. The diagnosis of the condition in life is not

easy, since similar lesions may be produced by other agencies, and fats and oils are not often considered as aetiological agents, because of a fairly general lack of awareness of their ability to produce such lesions and a failure to elicit, in consequence, the relevant facts in the patient's history. In the cases described in this paper aspiration of oil was not considered in diagnosis during life. The diagnosis at necropsy also presents some difficulty.

Classification and Predisposing Factors

Ikeda (1937) classified oil aspiration pneumonia into two types, which he called "adult" and "infantile." The "infantile" type occurs in debilitated children and adults as a result of the administration of food and drugs containing animal oils. This type is essentially a diffuse low-grade pneumonia, and one of the main grounds of distinction from the "adult" type is that it may undergo resolution. The "adult" type occurs only among the older age groups as a result of habitual self-administration of liquid paraffin, either orally or intranasally, in large total amounts. Animal oil is of little importance in the "adult" type, where the lesion is a chronic granuloma.

Among the factors predisposing to aspiration of oil into the lungs are the feebleness associated with early infancy and old age, the disabilities arising from affections of the central nervous system, debility, recumbency, dysphagia, and lesions of the mouth, throat, oesophagus, and upper respiratory tract. The character of the oil also is an important factor. Liquid paraffin is implicated as the causal agent in the great majority of cases. Its blandness would seem to be the reason for its known capacity to enter the bronchial tree without exciting any reflex action. The type of case in which oil aspiration pneumonia is most likely to be found is the chronic sufferer from respiratory infection who for long periods uses oral or nasal sprays of which liquid paraffin is an ingredient, or the sufferer from chronic constipation who seeks relief in a regular oral dose of liquid paraffin over long periods (Freiman *et al.*, 1940).

Pathological Changes

There cannot be said to be a typical naked-eye appearance. There are irregular areas of consolidation in the lung, varying in size from 1 cm. or less in diameter to that of a whole lung segment. The lesions, being fibrous, have a tough or hard consistence, and are grey or yellow in colour. On naked-eye examination and radiologically (Siddons, 1958) they have been most often wrongly diagnosed as neoplastic infiltration or as caseous consolidation of tuberculosis. Microscopically, different fields may show different stages of reaction. In some areas an accumulation of phagocytes may be found occupying the alveoli. They are large, foamy, and vacuolated in varying degree. Strands of collagen fibres may be seen insinuating between these lipophages. Giant cells formed by the coalescence of the lipophages are commonly present. In yet other areas there is dense fibrosis with circular lacunae. Appropriate stains show these lacunae to be filled with fat.

As might be expected, there are a number of complications which will each add its quota to the appearances described. Recurrent bronchopneumonia, bronchiectasis, abscess formation, and empyema have been known to complicate the disease. Bronchogenic carcinoma has been suggested as a complication.

The crux of the diagnosis lies in the demonstration of fatty material in lacunae in the densely fibroid lung tissue. To distinguish between mineral oils on the one hand, and vegetable and animal oils on the other, use has been made of the reaction of osmium tetroxide, which is reduced by the animal and vegetable oils to give a black precipitate, whilst mineral oils have no such action. Some investigators have carried out chemical analysis of lung tissue (Jampolis *et al.*, 1953; Wagner *et al.*, 1955).

Case Histories

Case 1.—A woman aged 54 was admitted to hospital on January 17, 1956, with a diagnosis of bronchitis, bronchiectasis, and right-sided heart failure. There was a history of chronic cough since childhood. A radiological report at that time mentioned gross bilateral emphysema secondary to chronic bronchitis, with bilateral bronchiectasis in the lower lobes. Investigation for tuberculosis was negative.

On admission she had marked dyspnoea at rest. There were diffuse bronchospasm and expiratory and inspiratory rhonchi, with coarse crepitations at both lung bases. Jugular venous pressure was raised. There was oedema of the ankles and sacrum. Laboratory investigation of the sputum showed normal flora, and two specimens were negative for acid-fast bacilli. The urine was sterile on culture, and apart from a trace of albumin was not otherwise abnormal. Blood urea was 47 mg. per 100 ml. Haemoglobin 87% (Haldane). White cells numbered 10,400 per c.mm., with normal differential count. Chest x-ray report on January 26 was a repeat of previous findings, but in addition there was general enlargement of the heart, the appearances suggesting myocardial failure. The patient responded temporarily to treatment, but died on February 7, 1956.

At necropsy the main findings were as follows. The abdomen was distended and the ankles and legs were oedematous. The trachea and bronchi were filled with frothy mucus. Both lungs showed diffuse emphysema, with extensive areas of patchy consolidation, particularly in the lower lobes. There was marked prominence of the bronchi and bronchioles. The right ventricle was hypertrophied. The liver on cut surface showed a nutmeg pattern. The peritoneal cavity contained 2 pints (1.1 litres) of serous fluid.

Histological examination of the lungs showed the changes of chronic bronchitis and emphysema. In many areas the alveoli were filled with large, foamy, pink macrophages, many of which contained two or more nuclei. In other areas there was a picture of extreme disorganization characterized by the presence of giant cells of bizarre shape, suggestive of foreign material in the lung tissue. In yet other areas were large collections of lymphocytes and plasma cells. There were also extensive areas of fibrosis, in the meshes of which, on staining with Sudan IV, were revealed a number of oil globules of varying size. These globules were found not to reduce osmium tetroxide. It was therefore concluded that the findings disclosed the presence of an oil aspiration pneumonia, in both acute and chronic stages, due to liquid paraffin.

Subsequent inquiry yielded no information as to the source of the aspirated oil. It was surmised that it was probably an oral or nasal spray containing liquid paraffin.

*Case 2.**—A man aged 55 years, a universal grinder in a steel foundry, died suddenly on November 1, 1957. For several years he had been under the care of his doctor on and off because of pains in the chest, shortness of breath, and chronic cough; he had, in fact, been treated for bronchitis. Three weeks before his death he was seen by his doctor because of an attack of "influenza," and he had never fully recovered.

The main findings at necropsy related to the chest. There was a quantity of mucus in the trachea. In the lower lobe of the left lung were two dense fibrous masses, each about 1 in. (2.5 cm.) across. A similar mass was present just above the hilum of the right lung. There was no significant emphysema. The heart was substantially dilated and hypertrophied on the right side. Pulmonary fibrosis was present, and an important point was whether it was of industrial or non-industrial origin.

Histological examination revealed varying degrees of fibrosis throughout the lungs. In many areas the appearance was of a honeycomb of clear spaces, which by staining with Sudan IV were shown to be filled with fat (Special Plate,

*This and the following cases were the subject of necropsies undertaken on behalf of H.M. Coroners for the City of Sheffield and the County Borough of Rotherham.

Fig. 1). Osmium tetroxide was not reduced by this fat (Special Plate, Fig. 2). The framework of the honeycomb was of fibrous tissue, but this varied in appearance from area to area. In some areas this framework was of wide, dense, relatively acellular fibrous tissue, and here the clear spaces were largest in diameter. In other areas the fat-filled spaces were smaller and the fibrous-tissue framework was more slender and relatively cellular, the cells being a mixture of fibroblasts and vacuolated macrophages, also shown to be filled with fat. Occasional giant cells were to be seen in this fibrous framework. In some areas the normal alveolar pattern was to be seen, many of the alveoli containing numbers of the large macrophages in various stages of vacuolation, which we have learned to associate with lipid pneumonia (Special Plate, Fig. 3). Here and there throughout the lung fields were collections of lymphocytes and plasma cells. The basement membrane of the bronchi was thickened. No evidence of silicosis or tuberculosis was noted. There was dramatic narrowing of the branches of the pulmonary artery, with marked medial thickening. This was seen to a lesser extent in areas of lung tissue only slightly affected by fibrosis.

The cause of death was cor pulmonale due to pulmonary fibrosis following lipid pneumonia. In view of the findings an effort was made to trace the source of the aspirated oil, and it was found that the deceased had been in the habit of taking liquid paraffin as a purgative. This was the only ascertained source from which the lesion could have arisen.

Case 3.—A man aged 78 who had worked in various coal-mines all his life. Since the age of 70 he had had at least two strokes, from which he had made substantial recovery. For a number of years he had had shortness of breath, cough, and chest pains. By July, 1957, when he was admitted to hospital for a minor condition, he could do little more than totter about owing to senility. He was kept in hospital because of the difficulty of home nursing, and during his stay a chest radiograph was reported as showing nodular opacities in both lungs due to pneumoconiosis. On December 11, 1957, while sitting in the ward, he had sudden pain in the left leg, and examination elicited all the signs of arterial blockage in this limb. Gangrene of the limb developed, and he died on December 13.

Necropsy was undertaken to determine whether pneumoconiosis was the cause, or a contributory factor in the cause, of death. He was found to have died of bronchopneumonia due to gangrene of the left leg caused by femoral-artery occlusion and arteriosclerosis. An incidental finding was that in the lower part of the middle lobe of the right lung there was a fibrous honeycombed area, which was not pigmented and the appearance of which suggested that it might be an oil granuloma. Histological examination of this area showed that three stages of reaction could be distinguished (Special Plate, Fig. 4). In some fields the alveolar septa could be made out, and within the alveoli were collections of cells showing various degrees of vacuolation, most of them being, in fact, of signet-ring shape. Sudan-IV staining revealed them to be filled with fat. Their nuclei and cell membranes were in various stages of disintegration. Fine fibrous strands were just beginning to make their appearance, insinuating amongst these lipophages. In other fields the alveolar pattern was obliterated, and there were sheets of these signet-ring cells, distended in varying degree with fat, interspersed here and there with not very large clear spaces, which also proved to be filled with fat. In this type of field fibrous bands were more obvious and giant cells were also present. In yet other fields the picture was of a honeycomb of large clear spaces of varying size, shown to be filled with fat. Osmium tetroxide was not reduced by this fat. The framework of the honeycomb was dense fibrous tissue.

So far as the relatives knew, he had not been in the habit of taking liquid paraffin. He was described, however, as being always constipated and taking remedies for that condition. It may therefore very well be that he was in the habit of taking liquid paraffin unknown to his relatives.

Case 4.—A widower aged 79, retired pattern-maker, died suddenly on February 14, 1958. He had for some time been in failing health, and had been under the care of his doctor in the recent past, having had two attacks diagnosed as cerebral thrombosis, but at the time of his death had not been seen by his doctor for some weeks.

At necropsy he was found to have generalized arteriosclerosis, with two areas of old softening in the left cerebral hemisphere. The lungs showed bronchopneumonic change, and indeed the cause of death was bronchopneumonia complicating arteriosclerosis and cerebral thrombosis. In the lower lobe of the right lung a patch of consolidation was discovered, which was larger and had a much harder consistency than the surrounding patches of bronchopneumonic consolidation. Histological examination of this area revealed a picture similar in essentials to that described in Case 3. Sudan-IV staining showed the area to contain a large amount of fat (Special Plate, Fig. 5), which failed to reduce osmium tetroxide (Plate, Fig. 6).

Discussion

For practical purposes the agent responsible for oil aspiration pneumonia in adults is liquid paraffin. It gains access to the lungs most easily by its use as an ingredient of nasal or oral sprays. Some years ago it was commonly found in such sprays, and we cannot but feel that there must exist a considerable number of oil granulomata of the lung arising from their use. The evidence suggests that, once in the lung, liquid paraffin remains as an inert foreign body for many years, and probably permanently. From inquiries we have made it appears that the tendency now is for nose drops and sprays to be made up in an aqueous medium. A substantial proportion (about 20%) is, however, still being manufactured in a medium containing liquid paraffin, and we feel that their manufacture in an oily medium should be stopped. It is also clear that liquid paraffin gains access to the lungs as a result of oral administration, even in those who suffer no disability. A bland oil introduced by catheter into the pharynx of a healthy adult whilst asleep on his back was shown to be present in the lung the following morning (Pirie, 1932). It appears that there is still in Great Britain a steady sale for liquid paraffin as a laxative, and in our opinion its use for this purpose requires reappraisal.

Oil granuloma of the lung is largely preventable. Its presence is known to dispose to recurrent respiratory infection, and in these days when our lungs have to contend with so many other noxious substances this is one lesion which, however small, we can well afford to be without.

Summary

Four cases of oil aspiration pneumonia are described, two in which it was a contributory cause of death and two in which it was an incidental finding.

The pathogenesis and pathology of the condition are discussed.

The use of liquid paraffin as an ingredient of nasal and oral sprays, and also as a laxative, is criticized.

Our thanks are due to Dr. C. H. Hollins and Dr. T. V. Cooper, of Dorchester, for their assistance and for permission to publish the clinical details of Case 1. We are also grateful to A. P. Lockwood, Esq., H.M. Coroner for the City of Sheffield and for the Rotherham Division of the West Riding of Yorkshire, for permission to publish details of Cases 2 and 4; and to N. S. Robson, Esq., H.M. Coroner for the County Borough of Rotherham, with whose authority we publish details of Case 3. For technical assistance and the preparation of the illustrations we are indebted to Mr. S. J. Gray.

REFERENCES

- Bennett, H. S., and Collins, E. N. (1952). *Gastroenterology*, 20, 485.
- Boyd, J. T., and Doll, R. (1954). *Brit. J. Cancer*, 8, 231.
- Freiman, D. G., Engelberg, H., and Merritt, W. H. (1940). *Arch. intern. Med.*, 66, 11.
- Ikeda, K. (1937). *Arch. Path. (Chicago)*, 23, 470.
- Jampolis, R. W., McDonald, J. R., and Clagett, O. T. (1953). *Int. Abstr. Surg.*, 97, 105.
- Pirie, A. H. (1932). *Amer. J. Roentgenol.*, 27, 578.
- Siddons, A. H. M. (1958). *Brit. med. J.*, 1, 305.
- Wagner, J. C., Adler, D. I., and Fuller, D. N. (1955). *Thorax*, 10, 157.

G. FORBES AND A. BRADLEY: OIL ASPIRATION PNEUMONIA

FIG. 1

FIG. 2

FIG. 1.—Dense fibrosis of lung with honeycomb of fat-filled spaces, typical of a late stage of oil aspiration pneumonia. (Sudan IV. $\times 66$.)

FIG. 2.—Similar area to Fig. 1 stained with osmic acid and showing no reduction by fat. ($\times 66$.)

FIG. 3

FIG. 4

FIG. 3.—Typical vacuolated lipophages of oil aspiration pneumonia lying in alveolus. (H. and E. $\times 264$.)

FIG. 4.—Section of lung showing different stages in reaction to oil. On left are sheets of lipophages between which fibrous strands are beginning to insinuate. Farther to right, fat-filled spaces are enmeshed in fibrous tissue which is cellular in character. Still farther right, fibrous tissue is denser and much less cellular. (H. and E. $\times 66$.)

FIG. 5

FIG. 6

FIG. 5.—Section of lung showing oil aspiration pneumonia in early stage of development. (Sudan IV. $\times 66$.)

FIG. 6.—Section from same area of lung as Fig. 5 stained with osmic acid. ($\times 66$.)