THE TRADE IN TRANQUILLITY*

BY

D. R. LAURENCE, M.D., M.R.C.P.*Senior Lecturer in Applied Pharmacology, University College and University College Hospital Medical School, London*

AND

D. A. POND, M.D., M.R.C.P., D.P.M.*Physician, University College Hospital and Maudsley Hospital, London*

"In your ideas of the powers of remedies do not be too sanguine, for you are liable to continued disappointment" (Sir Anthony Carlisle, 1824).

A new group of drugs for which very great claims have been made has recently appeared in the field of therapeutics. These drugs are generally called "tranquillizers" or "ataractics" (from a Greek word meaning peace of mind). There is no doubt that "it would be difficult to refute the fact that there is a tremendous need for sedation at the national level" (SeEVERS, 1954), unwilling though some of us may be to recognize it. It is, of course, not a new problem, for in 1857 the *Lancet* felt it proper to express its editorial disapproval of the use of drugs as a prop on which to face life, in these words: "that man's nervous system must be in a very disordered state who resorts to a pipe or cigar in order to tranquillize himself and induce an aptitude for the business of the commencing day." A substantial proportion of the population are in just such a disordered state, and pending an alteration in our whole way of life, of which there is no sign at present, the individual physician is faced with as many as one-third of his patients whose real need is peace of mind.

Even if, by psychotherapy, peace of mind could be accorded to these people, there are too many people and too few psychotherapists. Doctors feel obliged to try to meet this need with drugs, if only because their patients ask for them. The demand is colossal and ageless, but the motives for canvassing the drugs are equally great, not only a desire to heal but also a desire for money. The field is wide open, and a fortune is to be made by any commercial organization which hits on an effective tranquillizer, or which can persuade the medical profession or the general public that it has done so. The profession is no longer in the commanding position that it evidently had when Dr. Faustus said:

Be a physician, Faustus; heap up gold,
And be eternis'd for some wondrous cure:

The opportunity for heaping up gold now lies with the commercial drug houses, who are not unconscious of this, although the physician may still manage to be "eternis'd" if he is clever or lucky.

An Example of Failure

In the field of neuroses tranquillizers have provided an example of dramatic failure to accomplish what was, and indeed still is, claimed for them by their manufacturers (the results in psychoses are better, but are not considered in this paper). There are at least three important reasons for this: (1) All new drugs are perforce developed in the first place by animal experiments that, as shown below, are at present irrelevant to the clinical use to which tranquillizers are put. (2) The lack of knowledge of the clinician of how these drugs are

*Based on a talk given by one of us (D. R. L.) to the Section of Psychiatry at the Annual Meeting of the British Medical Association, Newcastle upon Tyne, 1957.

developed in the pharmacological laboratory, so that when a new compound is offered for clinical use he is unaware of its inadequate scientific background and in no position to be critical, and is obliged to rely on the assertions of the producers, whose enthusiasm is inclined to outrun the evidence, so great will be the prize for success. (3) The performance of misleading and inadequate clinical trials, so that new drugs go into general use unjustifiably early.

Double-blind clinical trial—that is, therapeutic trial in which neither the patient nor the physician assessing his progress knows what drug the patient is taking—is essential for the measurement of the effect of a drug on the variable and largely subjective disorders of neurotics. It is harmful to both doctor and patient that a succession of new drugs which have not undergone this procedure should be promoted amongst, and widely used by, the profession. The result is that they are dropped when news filters through that each drug does not, after all, accomplish what was claimed; and the next new drug (which is often merely a variant of an existing successful drug that has aroused envy in the breasts of other firms) is tried as the next drug-firm representative calls.

The Animal Experimental Background

The readiness of some physicians to administer to their patients, under uncontrolled conditions, these potentially toxic compounds, which have been found to produce a miscellany of actions in animals, is a source of wonder to academic and profit to commercial pharmacologists. Boldness in reasoning from animals to man is essential, but it is equally essential that careful evaluation on man should precede widespread use under uncontrolled conditions, and this is not being done at present.

The pharmacologist, working on animals in his laboratory, suspects, or rather hopes for, a therapeutic action if a substance: (1) has any action, electrophysiological or biochemical, on any part of the central nervous system suspected to be related to behaviour; (2) antagonizes on any test preparation—for example, isolated intestine, intact snail—substances known to occur in the central nervous system such as 5-hydroxytryptamine, acetylcholine, or adrenaline; (3) antagonizes substances which produce abnormal behaviour, such as mescaline and lysergic acid diethylamide; or (4) modifies animal behaviour—for example, so-called "experimental neuroses" in animals, alterations in the shape of spiders' webs.

It is important to state that the study of phenomena mentioned above is not criticized, it is the too facile application of the results to man and the too widespread promotion of compounds that have not undergone controlled evaluation in man which are condemned.

Pharmaceutical firms are in a highly competitive market for new tranquillizers, and one of their prime needs is a simple cheap test by means of which hundreds of compounds may be screened for potential psychiatric value. This means that even tests which might at first seem superficially relevant to the human condition—for example, "experimental neurosis" in animals—are by no means often used, as they are expensive to perform, involving prolonged and tedious repetitions and observations. It is much easier to give the drugs to spiders of a sort which spin a web every day, photograph the web, and analyse the results at leisure. By such means numerous compounds can be tested with minimum effort.

The following are some tests in use for development of tranquillizers: (1) Prevention of epileptic fits in mice induced by sound of appropriate frequency (audiogenic seizures). (2) Potentiation of barbiturate-induced sleep. (3) Effects on conditioned responses in animals (rats, cats). Restoration of conditioned reflexes which have been exhausted by failure to provide a reward following the stimulus. (4) Effects on conflict or stress-induced behaviour in animals (rats, cats, monkeys). (5) Taming effects on monkeys fierce by nature or as a result of surgery. (6) Alterations in the form of spiders' webs. The creature may

become more assiduous in attending to the angles of its web, its geometrical sense may become distorted, or it may even omit portions. It is contended that some drugs produce consistent alterations, and that effects are specific on the nervous system and it is not simply that the spider is feeling unwell. (7) Introduction of compounds into jars containing Siamese fighting fish. They float at differing angles to the surface of the water, and may even become peaceable.

If drugs of proved clinical merit are examined by these tests no consistent results are obtained, and drugs which are active in one test may be inactive in another. Thus no adequate screening test for tranquillizers exists.

Most workers are quite aware of the limitations of the methods in present use; but unjustified assumptions are made, leading to the offering of unproved drugs for general clinical use. The capacity to modify animal behaviour for the better would seem at first sight least irrelevant to the problem. However, there are great disagreements on the significance of animal behaviour in relation to human behaviour. The symptomatic resemblances of disturbed behaviour are certainly similar—the sweating, tachycardia, affective reactions, disturbed sleep, changeability, moodiness, disturbed sexual activity, and so on that have been described in humans have all been seen separately and in combinations in monkeys, dogs, cats, rats, sheep, goats, and many other animals. The problem is whether these behavioural resemblances are really homologous—that is, due to similar underlying functional mechanisms—or merely analogous. As it has been put by a wit referring to Dr. Masserman's work on the production and therapy of behaviour disorders in cats,

Masserman, the cat man,
Makes felines neurotic;
Are cat and human
Really asymptotic?

Nevertheless Masserman (1943) found that alcohol removed the symptoms of stress—which coincides with human experience.

Desirable Clinical Effects

The fundamental clinical problem is not the control of all symptoms engendered by any stress or frustration, since these experiences are common to all people and the symptoms can have biological value in causing the organism to take remedial action; the problem concerns the difference between persons in the ease with which the neurotic patient produces symptoms with inadequate or irrelevant environmental stress. Some efforts are beginning to be made towards obtaining strains of animals which react excessively to environmental stresses, and to use these with controls of "normal" animals. It is an open question whether the differences between normal and neurotic cats are homologous to the differences between normal and neurotic humans.

From the psychiatric point of view, drugs may be evaluated (a) by subjective reports, (b) by objective psychological tests, or (c) by an independent account of a change in the patient's work capacity, social relations, or physical functions such as sleeping, eating, sexual activity.

There are three main groups of symptoms whose relief by drugs might be possible and would certainly be desirable.

1. *Excitement and Overactivity.*—This can of course be abolished by general anaesthesia, but the aim here is to produce a drug that causes little or no disturbance of consciousness. A more careful analysis of the phenomenology of the excited patient might give a lead here. For example, hyperactivity, especially in children, seems to be due to a distractibility. Stimuli are too stimulating; the child cannot fail to respond to them, and new stimuli command his attention for a few seconds until another stimulus comes. More pure manic states show a predominantly affective change, the patient in very severe cases paying no attention to the environment, but being consumed entirely by an inwardly produced excitation.

2. *Depression.*—This affective state is opposite to that of mania, but it is not the opposite to overactivity and

euphoria, as can be seen in the way in which amphetamine sometimes makes depressives worse. There are retardation, preoccupation, and also some striking somatic changes suggestive of pituitary underactivity.

3. *Tension.*—There is a muscular component, as shown by E.M.G. studies, and a visceral or autonomic component (changes in sweating, pulse rate, etc.).

If we return to possible animal experiments we can see that the undesirable human psychiatric effects are in fact very difficult to produce in animals, with the possible exception of tension. Likewise if we wanted to proceed below a symptomatological level to try to elucidate the fundamental psychopathological mechanisms we are faced with so far unsolved problems of the organization of behaviour. In the present state of knowledge it is very doubtful if any theoretically based animal test can be expected to include potentially useful drugs, nor is it certain that such tests might not cause promising drugs to be abandoned.

Some New Drugs

The principal new drugs at present advocated as tranquillizers are: chlorpromazine ("largactil"), reserpine ("serpasil"), meprobamate ("equanil," "mepavlon," "miltown"), benactyzine ("nutinal," "suavitil"), and methylpentynol ("oblivon," "somesin"), to mention only one representative of each chemical group. The classification "tranquillizers" at present includes substances with greatly differing pharmacological actions, and it is probable that, as knowledge grows, some members will be removed and perhaps others added until ultimately the group becomes homogeneous or else disappears as an entity.

Tranquillizers are not necessarily hypnotics, although they may promote sleep. Hypnotics, on the other hand, may be tranquillizers. It is probably unprofitable with our present knowledge to attempt too rigidly to distinguish between these groups. Experimental work is being done in man to distinguish any fundamental properties tranquillizers may have.

Meprobamate stems from the discovery of mephensin by Berger in 1946. It is said that mephensin was first made for testing as a possible preservative for penicillin. Its interneurone depressant effects were uncovered during routine toxicity tests when it was found that high doses produced reversible paralysis in animals without interfering with respiration. The discovery of *chlorpromazine* in France in 1950 arose from work on the therapeutic properties of phenothiazine compounds in malaria and as antihistamines, work which also produced promethazine ("phenegan"), diathazine ("diparcol"), and ethopropazine ("lysivane").

Rauwolfia can hardly be said to have been discovered in the same sense, having been used for insanity since ancient times in India, but only recently has it been investigated.

It is sad that the tranquillizer *benactyzine*, which might be claimed to have had the most "scientific" development, should prove to be of no greater therapeutic efficacy than any other of these controversial drugs. Benactyzine, which has a formula related to atropine, was synthesized in 1936 in a systematic search for compounds with peripheral parasympatholytic or anticholinergic effect. It was shown to be a mydriatic in 1938. In 1955 some workers considered that any drug opposing the action of acetylcholine at the periphery might also do so in the brain, might block "certain neuronal pathways," and affect behaviour. They therefore studied the effect of a group of anticholinergic compounds on experimentally induced conflict behaviour in cats and on stress-induced behaviour in rats. They found benactyzine to be the most effective of their series at normalizing the behaviour of these animals, and so the drug was introduced into clinical practice, with disappointing results. Incidentally, chlorpromazine failed to normalize the behaviour of the rats.

Methylpentynol is related chemically to amylene hydrate, long known to be a hypnotic.

Pharmacological Observations

A brief survey of these drugs will show the wide variation of their properties.

Of *chlorpromazine*, Brücke (1956) has remarked that it "could be said to be a substance which has nearly exclusively side-effects." Certainly it has an inconveniently wide range of activity, which is said to be why the name "larg-actil" was coined. It has antiadrenaline, local anaesthetic, atropine-like, antipyretic, weak ganglion-blocking, antihistamine, and quinidine-like actions. Allergy occurs, leading to dermatitis, jaundice, or agranulocytosis. On the C.N.S., low doses abolish Parkinsonian tremor and large doses induce a characteristic Parkinsonian syndrome. It abolishes spasticity and tetanus. It potentiates barbiturates powerfully. It abolishes conditioned reflexes, but not inborn reflexes (benactyzine, if anything, slightly revives conditioned reflexes, as do insulin and electric and leptazol convulsions). Chlorpromazine quietens aggressive rhesus monkeys and prevents hyperthyroidism in wild hares caught and exposed twice daily to barking dogs. Arousal reactions induced by stimulating the frontal hypothalamus or ascending reticular formation are suppressed. E.E.G. studies are compatible with this being the principal site of action. It is also a powerful anti-emetic.

Reserpine acts on the C.N.S. almost exclusively. Electrical stimulation of the anterior hypothalamus near the parasympathetic centre produces sleep, miosis, bradycardia, and decreased peripheral vascular resistance. Since reserpine produces all these effects it seems likely that it may exert its principal action in this neighbourhood. The mode of action is unknown, but it has been suggested that the demonstrated ability of reserpine to release 5-hydroxytryptamine in the brain may be relevant, although this cannot account for all the observed effects. Nor is it yet proved what part 5-hydroxytryptamine plays in mental abnormalities; but pharmacologists have seized upon the fact that a 5-hydroxytryptamine antagonist, lysergic acid diethylamide, induces psychotic states, thus linking it with reserpine. The tranquillizing effect of reserpine is marked in Siamese fighting fish. The clinical effects in man have given less grounds for satisfaction in neurotics, but good results in psychotic excitement. Other effects of reserpine are dyspnoea, nasal stuffiness, diarrhoea, and a Parkinsonian syndrome which is said to be relieved by atropine, although some might think it wiser to stop giving the drug. It is of interest that two chemically different tranquillizers (chlorpromazine and reserpine) should have this unusual toxic effect. Reserpine also may activate peptic ulcers.

Meprohamate has actions similar to mephensin, from which it has been developed. It depresses multisynaptic reflexes and the reticular formation. Its principal advantage over its parent is that it has prolonged action after oral administration. It also causes dermatitis, mental excitement, and diarrhoea, and addiction has been reported.

Benactyzine has peripheral actions similar to atropine, and also has feeble antihistaminic and quinidine-like effects. Human experiments on the mental effects are probably more complete for benactyzine than for other tranquillizers, although the mode of action remains hidden. It abolishes alpha rhythm in the E.E.G. in man. It has been described by various authors as inducing a devil-may-care state, of causing a lack of ability to make up one's mind, and of causing lack of attention, so that it has been suggested that patients taking more than 4-5 mg. per dose should not drive a car. Benactyzine also prolongs the reaction time and increases the frequency of errors in mental tests. A writer has advised that the drug should be used with caution, if at all, on active businessmen, for it has been found that subjects forget what they were going to say, and sometimes lapse into incessant laughter. Such an action would be difficult to predict from animal experiments.

Methylpentynol is a comparatively weak hypnotic and there is reason to doubt whether its qualitative effects differ significantly from those of hypnotics: such comparisons are

difficult because it is hard to choose equipotent hypnotic doses. It potentiates barbiturates and also alcohol, as was discovered by the man who, having read, in a newspaper, of its beneficial effects on anxiety, took some to allay his well-justified fear that the amount of gin he had consumed would lead to his being arrested for being drunk in charge of his car. He found that, although methylpentynol allayed his anxiety, it also made inevitable the event which he feared. Dermatitis occurs occasionally. A young lady has been reported as suffering a fate worse than death, due to the drug; and one or two people have actually died of overdose. On the whole, however, it seems to be a fairly safe drug.

Action of Tranquillizers

It is not possible to synthesize these observations into a coherent theory of drug action, but clearly it is desirable that both animal and human data should be collected on all these drugs until there is enough for both psychiatrist and pharmacologist to begin to know what they are really about. It is clear that the action of all tranquillizers depends greatly on the personality of the subject, a field in which little work has yet been done. At the present time it would seem better to try to test the tests by administering the few drugs known to be of benefit in psychiatry to animals, and in this way to find what animal investigations seem to give the best results. Phenobarbitone, amylobarbitone, barbiturate-amphetamine mixtures ("drinamyl," "dexylal," "desbutal"), and chlorpromazine are up to now the most effective drugs in psychiatry.

In the meantime it may be expected that the pharmaceutical industry will bombard the clinician with what are often unwanted and sometimes even dangerous drugs, with once in a while a really valuable therapeutic advance. We believe that an unnecessarily large number of the former are advertised for clinical use for every one of the latter produced. It is to be hoped that the discovery of a "complete" or "absolute" tranquillizer is an impossibility, for it is essential to distinguish between biologically useful and useless symptoms. The removal of fear, guilt, and anxiety under all circumstances would render man less than human. However, tranquillizers have their place as what H. G. Wells called "Doors in the Wall" to provide temporary refuge from acute stress, as all who have taken alcohol know.

Aldous Huxley (1954) has defined the "perfect" tranquillizer as a drug: "which will relieve and console our suffering species without doing more harm in the long run than it does good in the short. . . . It must be . . . less likely to produce undesirable social consequences than alcohol or the barbiturates, less inimical to heart and lungs than the tars and nicotine of cigarettes. . . . It should produce changes in consciousness more interesting, more intrinsically valuable than mere sedation or dreaminess, delusions of omnipotence or release from inhibition." Huxley concludes that we shall not escape "the need for frequent chemical vacations from intolerable selfhood and repulsive surroundings," because "most men and women lead lives at worst so painful, at best so monotonous, poor and limited that the urge to escape . . . is, and has always been, one of the principal appetites of the soul." Although this may be true, it might be wiser for the community to start trying to alter the conditions of life to render the "perfect" tranquillizer unnecessary, for it will certainly be long in coming, and may very well be a chimera and altogether unattainable.

Meanwhile doctors will do their best with the unsatisfactory drugs at their disposal to reduce the total of human misery.

REFERENCES

- Brücke, F. T. (1956). *20th International Physiological Congress, Brussels. Abstracts of Reviews*, p. 465.
 Carlisle, Sir A. (1952). Quoted in J. Langdon-Davies's *Westminster Hospital*. Murray, London.
 Huxley, A. (1954). *The Doors of Perception*. Chatto and Windus, London.
 Masserman, J. H. (1943). *Behavior and Neurosis*. University of Chicago Press.
 Seevers, M. H. (1954). In *Pharmacology in Medicine*, edited by V. A. Drill. McGraw Hill, New York.