Focal or Partial Epilepsies

The E.E.G. has taught us a great deal about these forms of epilepsy. Deductions on the site of origin based on clinical observation were confirmed and methods of spread over the cortex have been studied, but there are two ways in which we have gained original knowledge—the recognition of foci of discharge in the so-called silent areas of the frontal and temporal lobes, and the bilateral occurrence of so many focal discharges. We used to believe that major fits were of two kinds—those with a generalized onset and those with a focal onset. It is now evident that when a major fit begins with, say, twitching or tingling of a limb, what in fact has happened is that a focal fit has initiated a generalized convulsion—there have been two attacks, the focal fit having initiated the major fit much as electrical stimulation might. This fact, which is so evident in electroencephalographic records, was really evident from observation of the attacks too, because a major fit always has the tonic phase first, the clonus being caused by interruption of tonus, while a clonic focal fit consists of much more rapid jerks of part of the body, and the march of events is clonus, tonus, and then clonus—obviously two different kinds of fit succeeding each other.

Another fundamental fact that must be recognized to achieve a logical understanding of focal fits is this: that, however elaborate may be the movements or feelings in an epileptic attack in which consciousness is preserved, they are always due to focal disturbance of brain function. The only difference between an attack with a primitive muscle twitch and one with co-ordinated, apparently purposive disorder of social behaviour, is that the first is caused by disturbance of function in the relatively primitive motor cortex, and the second in a more highly elaborated integrative part of the cortex. Those attacks we most readily think of as “focal”—the clonus of a limb, local tingling or numbness, flashing light, a smell, a buzzing sound, and so on—all happen in the first cell station of reception of sensations into the brain or the last cell station of emergence of movement. They are simple phenomena, which were therefore understood early in the study of epilepsy. The “psychic fits” or psychomotor attacks, the elaborate hallucinations, dreamy states, emotional experiences, or visceral disorders of epilepsy are all just as truly caused by focal discharge in the brain as the experimental surgery of Penfield and others has demonstrated. That being so, focal fits can be classified according to their clinical pattern—motor, sensory, psychic, and so on—upon the basis of their known origin. The second is probably the more useful, because, as the list of epileptic events shown above indicates, there are too many forms of epileptic experience to make their division a profitable one. A classification based on the part of the brain first affected, with a statement of the clinical events, is acceptable in all countries.

Focal fits may be cortical or subcortical, but much of the cortex, like that of the island of Reil and the undersurface of the temporal lobe, is buried.

Cortical discharges may occur in *primitive areas* or *association areas*, or may spread from one to the other. They may, as so often in the temporal lobe, be *bilateral* and they may march over the cortex in a *Jacksonian* way. We now recognize that that Jacksonian march may include very complex experiences in which division into physical and psychological becomes meaningless.

REFERENCES

- Gastaut, H. (1954). *The Epilepsies. Electro-clinical Correlations*. Thomas, Springfield, Illinois.
Gibbs, F. A., and Gibbs, E. L. (1941). *Atlas of Electroencephalography*. Cambridge, Mass.
Penfield, W., and Jasper, H. H. (1954). *Epilepsy and the Functional Anatomy of the Human Brain*. Churchill, London.
— and Kristiansen, K. (1951). *Epileptic Seizure Patterns*. Thomas, Springfield, Illinois.
Symonds, C. (1954). *A.M.A. Arch. Neurol. Psychiat.*, 72, 631.
Williams, D. (1953). *Brain*, 76, 50.
— (1956). *Ibid.*, 79, 29.

VALUE OF THE E.E.G. IN DIAGNOSIS OF EPILEPSY*

BY

DENIS HILL, M.B., F.R.C.P., D.P.M.

Institute of Psychiatry, London

The value of clinical electroencephalography in the diagnosis of epilepsy depends on the accuracy of the technique in throwing light on two issues: (1) From which area or from which neuronal system does the epilepsy originate, and is there more than one such focus or system from which it originates? and (2) What is the nature of the pathological process, if one is present, of which the epilepsy is a symptom? Diagnosis is therefore both topographic and aetiological. Considerable advances have been made in the last few years in respect of the first issue, and careful repeated E.E.G. investigations planned with knowledge of the clinical problem involved will in most cases answer the question, From where in the brain does the epilepsy arise? Very often, of course, such investigations merely confirm the clinicians' expectations, demonstrating what could be anticipated from a knowledge of the aura and fit pattern which the patient exhibits. Only if neurosurgery is contemplated are such intensive investigations vitally necessary. With regard to the second issue, which is of such great practical importance, the E.E.G. technique is of more limited value in helping us to decide the nature of the pathological process of which the epilepsy is a symptom. Here again, however, some advance has been made. These two issues are the subject of this paper.

Improper Use of the E.E.G.

The E.E.G. is often used as a means of helping to decide whether a given patient's symptoms are epileptic or psychogenic or are due to some non-cerebral cause, such as syncope. I believe this is an improper use of the technique and that in fact the results of doing this can be misleading. A single routine normal E.E.G. is not evidence against epilepsy, nor indeed is the finding of epileptic discharges in the E.E.G. evidence that the patient's symptoms are necessarily epileptic in nature. Such discharges occur in the E.E.G.s of some non-epileptic persons, in many with degenerative brain disorders, in some psychotics, and in cases of behaviour disorder. The prescription of anti-convulsants on the finding of such discharges in the E.E.G. is unjustifiable and often illogical, and may lead the clinician to overlook more important issues for his patient.

The recognition of the epileptic nature of any symptom or group of symptoms is, I believe, a clinical task and must remain one. When the clinician requests an E.E.G. to confirm the diagnosis he is not so much interested in the question of whether the E.E.G. demonstrates focal discharge in the expected area—he is merely disappointed but does not alter his diagnosis if it does not—as in the question of whether the E.E.G. demonstrates the presence of a pathological process, either local lesion or a diffuse one, which is of such importance for the prognosis. However, when the question of surgical removal of a focal epileptogenic area arises both these questions are of equal importance.

Classification of E.E.G. Characters

Some 20 years ago Gibbs, Gibbs, and Lennox (1938) described the first classification of the E.E.G. characters to be found in epileptics. Their categories were described in terms of frequency-voltage patterns of discharge—the actual appearances of the rhythm—and they paid no attention to

*Read to the Section of Neurology at the Annual Meeting of the British Medical Association, Newcastle upon Tyne, 1957.

sites in or on the brain from which these discharges arose. Four different frequency-voltage patterns were related in this classification to three main clinical seizure types: paroxysmal spikes with grand mal, 3 per second or slower

alternating spikes and waves with petit mal, and 4 per second square-topped waves with psychomotor seizures. It was suggested that when one of these patterns was found in the E.E.G. the patient suffered from the corresponding type

of seizure. This classification soon proved unworkable, because it was found: (a) that non-epileptic patients occasionally showed such patterns in the E.E.G.; (b) the age of the patient was an important factor in determining the type of discharge; (c) there was no place in the classification for focal epilepsy; and (d) the type of E.E.G. discharge bore little relation to the type of clinical seizure or to the pathological processes which might be associated.

A more workable classification was proposed by Jasper (1941), and recent developments have largely stemmed from this. In this classification we can describe (a) epilepsy of cortical onset, having a focal character, (b) epilepsy primarily of subcortical onset, having no focal character, and (c) epilepsy arising from numerous areas both in the cerebral cortex and in subcortical structures, the latter being associated with a variety of diffuse pathological processes. To these three categories must be added a further rarity, first described by Tükel and Jasper (1952), in which epilepsy arises from a focal cortical area but spreads immediately to the subcortical system, giving rise in the E.E.G. to discharges of subcortical type and, in the event of a seizure, a fit-pattern suggesting a subcortical onset. This type of epilepsy is described as "secondary subcortical" in view of the primary and initiating cortical focus. These four categories of E.E.G. pattern are fairly distinctive, can be demonstrated repeatedly by careful study, have physiological meaning, and make, I believe, neurological sense. Moreover, their recognition has assisted in the second and important question of diagnosis—the question of the associated pathology that may be present. These four categories are discussed in turn.

The Pathological Process

Focal cortical epilepsy arising from a single area of cortex is recognized in the E.E.G. by the presence of a focus of spike or sharp-wave potentials which may occur singly or in groups. It may be associated with normal background electrical activity, with reduced or absent background activity, or with the presence of abnormal slow-wave activity in the area. When the focus is associated with a small superficial atrophic lesion such as an atrophic gyrus or cortical cicatrix, the background activity is often normal, particularly if the seizures are infrequent. When the focus is associated with a more extensive cortical or subcortical atrophy there is likely to be a reduction or absence of normal background activity, whereas, if the association is with a neoplasm, the background activity is likely to be abnormal in frequency, usually with slow waves of irregular frequency and form. The finding of a single easily located spike or sharp-wave focus, together with locally abnormal background activity in the region of the spike focus, is fair evidence in favour of a local pathological condition, but such a finding always requires confirmation by further study.

FIG. 1.—Case 1. Showing a normal E.E.G. with a symmetrical alpha rhythm which blocks on visual attention. No abnormalities seen in the frontal areas.

FIG. 2.—Case 1. The E.E.G. examined under intravenous thiopentone in a light stage of narcosis. The reduction in barbiturate-induced fast rhythm is evident throughout all areas of the left frontal region.

It must be emphasized that it is the combination of a single focal spike discharge seen repeatedly only in one area, together with abnormality in the rhythmic activity in the neighbourhood, which provides the clearest diagnostic evidence. Spike discharges are recorded from viable cortex which is adjacent to dead or diseased cortex. At times the sites for these discharges are at a considerable distance from the gross lesion. When there are multiple foci, which is often the case in large atrophic lesions, such as porencephalic cysts, it is impossible from the appearances and siting of the discharges to locate the lesion. This can be done only by a study of the background activity of the E.E.G., which will be reduced or abnormal over the lesion. Various methods for demonstrating abnormalities in the background activity are available, the most useful being the introduction of an intravenous barbiturate.

As Pampiglione (1952) first observed, areas of damaged or dead cortex fail to develop the fast rhythmic activity which the drug induces in normal cortex (Case 1: Figs. 1 and 2). This procedure also has the advantage that in many cases narcosis facilitates the firing at the epileptic focus. Epileptogenic cortical foci can, of course, occur in any area of the cerebral cortex. Difficulty, however, arises in identifying by the E.E.G. those which are situated on the orbital surface of the frontal lobes or on the medial surface of the hemisphere. In both instances the discharges may excite subcortical structures and give rise to widespread unilateral or bilateral discharges. In both instances, also, there is no means of studying the characters of the electrical activity of adjacent cortex.

Temporal-lobe epilepsy, which is a common form of epilepsy and possibly accounts for from a quarter to a third of all cases, must be classified as a focal cortical epilepsy, the site of origin being in the infero-medial surface of the temporal lobe and the buried grey matter of this area, the hippocampus, uncus, and amygdaloid nucleus. As Professor Penfield showed some years ago an atrophic condition in these structures and in the cortex over them is the commonest lesion associated. While the E.E.G.s in these cases often show spike discharges widespread over the anterior temporal cortex and Sylvian regions, in many cases this is not so, and the single E.E.G. recorded with the patient awake may be normal. As Gibbs *et al.* (1947) showed, discharges are often facilitated when the patient falls asleep or is narcotized with a barbiturate. The location of the discharge may, however, still be difficult, and the distinction between an orbital-frontal, temporal, or posterior temporal focus be impossible to make with scalp electrodes. For this reason my colleagues and I have for some years advocated the use of needle electrodes inserted laterally from below the zygomatic arch to lie beneath the middle fossa of the skull, just lateral to the foramen ovale.

After the needles are inserted the patient is given intravenous thiopentone, which excites the discharges if they are not present, gives a clear record free from muscle-potential artifact, and, most important of all, induces fast rhythm in all normal cortex. Here again the same principle to which I have referred applies. Atrophic cortex shows reduction or absence of the barbiturate-induced fast rhythm, and this is a surer guide to the site of maximal pathology than is the site of a spike discharge when a number of different such foci are evident. In temporal-lobe epilepsy independent spike discharges may be recorded from both anterior inferior areas of the temporal lobes by means of the sphenoidal needle electrodes, and indeed spike discharges in

these cases may also occur in other areas of the cortex. Dr. W. A. Kennedy and I (1958) have been able to show, however, that reduction of fast rhythm under barbiturate narcosis recorded beneath the temporal lobe of one side predicts Ammon's horn sclerosis on that side at the 1% level of confidence.

There is fairly good evidence now that any chronic local lesion, or any diffuse pathological process in the brain, which has given rise to focal epilepsy may in time give rise to other epileptogenic foci in the cortex. This is seen in the E.E.G. as multiple independent spike discharges, recorded from different areas. From the appearance of these discharges alone it is not possible to identify the primary focus—that is, the epileptogenic area from which the clinical seizure is initiated and which itself was set up by the lesion. Abnormal background activity may provide the answer, but not invariably. In this event it may be necessary to demonstrate the primary focus by activating a seizure under E.E.G. control. For this purpose intravenous 5% or 10% leptazol, and more recently 10% bemegride, given slowly, are used. By this means the seizure can be witnessed and the origin and spread of the discharge recorded in the E.E.G. When the aura and the pattern of the induced seizure coincide with the patient's habitual seizure, and the E.E.G. has recorded a focal origin of the seizure, one can feel confident that the origin has indeed been located. This, however, is a difficult procedure in many cases, and it is useless to attempt it if the habitual seizure is one in which a tonic-clonic convulsion occurs early in the march of events.

Subcortical Seizures

This description of the E.E.G. characters in focal cortical epilepsy must suggest that all such cases occur in patients whose epilepsy is the result of a brain lesion of some kind. The corollary of this is that the finding of a single spike focus, in repeated E.E.G.s, must always be regarded as good presumptive evidence of a local lesion. In the other large groups of epilepsy—those in which the discharges arise primarily in subcortical structures—this is not the case. In

FIG. 3.—Case 2. E.E.G. shows a spike focus at the left superior frontal electrode (channels 3 and 4), and also a burst of high voltage bilaterally synchronous 3-per-second spike-and-wave activity. In the lower record a classical attack of petit mal is recorded.

these groups are those epileptics who can be called "idiopathic," and those patients in whom the brain has been damaged by pathological processes of a diffuse character. The E.E.G.s in these groups show widespread complex discharges affecting both hemispheres, often in a symmetrical manner. The classical bilaterally synchronous 3-per-second spike-and-wave discharge occurring upon a background of normal cortical activity is the type of discharge—it seems at the present time the only type of E.E.G. discharge that can be associated with idiopathic epilepsy. Yet there are rare instances in which even this type of subcortical discharge is secondary to a cortical focus related to a local lesion. This is the case of "secondary subcortical epilepsy" (Case 2: Fig. 3). Three-per-second spike-and-wave discharges occur much more commonly in the E.E.G.s of young epileptics than in those of adult patients, and the meaning of this is not clear. Is idiopathic epilepsy largely a condition of childhood? Since epilepsy, particularly major seizures, can itself very probably damage the brain by anoxia and therefore be itself a cause of further epilepsy, the problem is a difficult one.

Widespread bilateral fast and slow forms of spike-and-wave discharge, episodes in which groups of spikes predominate over slow waves, and those in which slow waves predominate over spikes, all suggest a subcortical origin for the epilepsy and an association with some diffuse pathological process in the past—infective, toxic, or degenerative. The more severe the pathological process the longer the epilepsy has lasted, and the more severe it has been the greater the likelihood that, in addition to these subcortical discharges, there will be random cortical foci of discharge as well. These were cases called by Jasper (1949) the "diffuse dysrhythmias." They always suggest diffuse brain damage.

As in primary cortical epilepsy, so in the subcortical epilepsies the single routine E.E.G. may show no specific features and be of little diagnostic value. There may be excess slow activity related to recent seizures or to the post-ictal confused state of the patient, or, again, this pattern may be the result of medication. Focal irregular slow activity recorded in a single E.E.G. following a seizure may reflect the post-ictal state of the cortex near to the focus and *not* indicate, as it otherwise might, a cerebral tumour. It is therefore necessary to interpret the E.E.G. findings in the light of the patient's clinical state at the time the record is made. To elicit discharges, to observe their location and morphology, and so to observe which physiological systems are involved in the discharge, and further to study the background activity, it may be necessary to plan an E.E.G. study under conditions in which the different variables are known, if not controlled. Medication may have to be withdrawn, seizures may have to be elicited and witnessed, activating procedures known to stimulate cortical foci or subcortical systems may have to be used. For this reason I suggest that the value of the E.E.G. in the diagnosis of epilepsy depends upon how it is used and with what end in view. The practice of taking single E.E.G. records, unrelated to the clinical problem which the patient presents, and often without knowledge of that problem, is a procedure of little value in most cases and may be frankly misleading. E.E.G. investigations should perhaps be reserved for those patients in whom such investigations can answer certain specific questions. Such investigations have to be planned, and will perhaps involve several hours' work under varying conditions. They can, however, provide understanding of the nature of epilepsy which no other method can do.

Illustrative Cases

Case 1: Figs. 1 and 2.—Male, aged 52, with three months' history of convulsions with remembered aura of twitching of lower lip and three weeks' history of focal motor attacks arising in the right face and spreading to right arm and leg. Two weeks' progressive right hemiparesis with mental deterioration. Examination showed right hemiparesis affecting face and arm more than leg. X-ray film of skull and lumbar encephalogram, normal. Left carotid arteriogram showed middle cerebral artery depressed

and pathological circulation in the posterior frontal areas. Biopsy: glioblastoma multiforme.

Case 2: Fig. 3.—Male aged 18. Since the age of 13 has had a number of minor attacks without warning in which he appears dazed. A history of cyanosis after birth following a bad forceps delivery. Examination showed no abnormal physical signs. X-ray film of skull normal. Lumbar encephalogram showed mild left hemiatrophy mainly affecting the left middle cerebral artery territory.

REFERENCES

- Gibbs, F. A., Gibbs, E. L., and Fuster, B. (1947). *Trans. Amer. neurol. Ass.*, 72, 180.
 — and Lennox, W. G. (1938). *Arch. Neurol. Psychiat. (Chicago)*, 39, 298.
 Jasper, H. H. (1941). In Penfield and Erickson's *Epilepsy and Cerebral Localization*, p. 380. Thomas, Springfield, Illinois.
 — (1949). *Proc. 2nd Int. Congr. E.E.G., Paris, Electroenceph. clin. Neurophysiol.*, Suppl. No. 2.
 Kennedy, W. A., and Hill, D. (1958). *J. Neurol. Neurosurg. Psychiat.* In press.
 Pampiglione, G. (1952). *Electroenceph. clin. Neurophysiol.*, 4, 79.
 Tükel, K., and Jasper, H. H. (1952). *Ibid.*, 4, 481.

MEDICAL TREATMENT IN EPILEPSY*

BY

H. HOUSTON MERRITT, M.D.

Director, New York Neurological Institute

Various drugs were used by ancient and mediaeval physicians in the treatment of epilepsy, but the treatment methods of those times were mainly the invocation of supernatural powers and the administration of substances which had magical properties. The pharmacological approach to the treatment of epilepsy did not begin until the middle of the nineteenth century. On the occasion of the presentation of a paper on the treatment of epilepsy by E. H. Sieveking before the London Medico-Chirurgical Society on May 11, 1857, Sir Charles Locock, the presiding officer, remarked that, since the seizures in epilepsy were often related to hysteria or the menses, he had been led to try bromide of potassium. He had used bromides successfully in a case of epilepsy associated with sexual excitement after all other medications had failed. In 14 or 15 additional cases bromides had failed in only one.

Bromides were the standard therapy for epilepsy until 1912, when Hauptmann administered phenobarbitone to his patients as a sedative. He was greatly pleased to find that it had an anticonvulsant activity equal to or greater than that of the bromide. The next significant advance in the pharmacological therapy of epilepsy came with the introduction of diphenylhydantoin (phenytoin) (Merritt and Putnam, 1938). Since that time a large number of compounds have been tested in patients. Many of these have been discarded as valueless or too toxic, but a few have added greatly to our ability to manage patients with seizures.

General Principles of Therapy

The treatment of patients with convulsive seizures is divided into three parts: first, the removal of any factors, organic or psychological, that may be playing a part in the occurrence of seizures; second, regulation of the physical and mental hygiene; and, third, the administration of anticonvulsant drugs.

The surgical and psychological aspects of epilepsy are discussed by Dr. Penfield and Lord Cohen. My remarks are confined to the medical therapy.

The objective of all anticonvulsant therapy is to establish and maintain a reservoir of drug sufficient to control the

*Read to the Section of Neurology at the Annual Meeting of the British Medical Association, Newcastle upon Tyne, 1957.