These three viewpoints among many are representative of the difficulty of finding a satisfactory terminology.

The present field study supports the main contention of Shubert *et al.* that seronegative cases should be regarded as examples of a *different disease*. The characteristics which they selected as points of differentiation between seropositive and seronegative disease with their limited series of cases are not substantiated by the present study, but other characteristics appear to provide even more striking evidence. Thus, Shubert *et al.* considered that there were notable differences between the two disease states as regards the incidence of pyrexia, sore throat, and rashes, but this is not supported by the present study. Nor is the contention supported that in the seropositive disease lymphocytes exceed 50% of the total white cell count, whereas in the seronegative disease they tend to be less than 50% of the total count.

From the clinical standpoint, the really notable features have been set out in Table XV—namely, unequivocal differences in the age, sex, and seasonal incidence of the two diseases, and as regards laboratory findings the fact that the rate of disappearance of G.F. cells from the peripheral blood distinguishes the two diseases.

If the adjective "infectious" is employed, it is clear that the attribute of infectivity is a feeble characteristic of each disease state, but is more highly developed in the seronegative disease than in the seropositive.

Summary

Over a period of 30 months, 498 cases suspected on clinical grounds to be cases of glandular fever were found to comprise 242 cases Paul-Bunnell positive, 100 cases Paul-Bunnell negative and haematologically positive, and 156 cases due to other diseases.

On purely clinical grounds it has not been possible to distinguish seropositive cases from seronegative cases other than in retrospect: seropositive cases tend to be the more severe.

There were notable differences in the seasonal trends and in the age and sex incidence of the two disease states.

The infectivity of either disease state was low, but the seronegative disease appeared to be the more infectious.

Average data are presented about the duration of the positive Paul-Bunnell reaction and the associated presence of G.F. cells in the peripheral blood film.

A positive Paul-Bunnell reaction is found approximately twice as long as G.F. cells in the peripheral blood film.

A doubtful positive Paul-Bunnell reaction, too weak to titrate, has been found to be significant and associated either with an incipient phase or, more usually, a regressive phase of the seropositive disease.

In seropositive cases the G.F. cells are present in significant numbers in the peripheral blood film approximately twice as long as in seronegative cases.

We gratefully acknowledge the assistance of the members of the Glandular Fever Committee, of the hospital laboratory services, of the general practitioners, of the C.O. Military Hospital, Wheatley, and particularly of Dr. Rosemary Biggs, who collaborated in the collection of data for this field study.

REFERENCES

- Barrett, A. M. (1941). *J. Hyg. (Lond.)*, **41**, 330.
 Davidsohn, I. (1938). *Amer. J. clin. Path.*, **8**, Techn. Suppl., 1, 56.
 Downey, H., and McKinlay, C. A. (1923). *Arch. intern. Med.*, **32**, 82.
 Leibowitz, S. (1953). *Infectious Mononucleosis*. Grune and Stratton, New York.
 — (1956). *Ann. intern. Med.*, **44**, 717.
 Newell, K. W. (1957). *J. clin. Path.*, **10**, 20.
 Niederman, J. C. (1956). *Yale J. Biol. Med.*, **28**, 629.
 Pfeiffer, E. (1889). *Jb. Kinderheilk.*, **29**, 257.
 Shubert, S., Collee, J. G., and Smith, B. J. (1954). *Brit. med. J.*, **1**, 671.

ACUTE APPENDICITIS

BY

J. A. CAMPBELL, F.R.C.S.Ed.

AND

D. C. McPHAIL, F.R.C.S.Ed.

From the Royal Infirmary of Edinburgh

The number of deaths resulting from acute appendicitis in England and Wales was remarkably constant for the years 1935 to 1938, the successive annual totals being 3,178, 3,052, 3,039, and 3,027. By 1944 the number of deaths had fallen to 1,881, and since then the steady decline in mortality has been maintained, the comparable total for 1954 being 969 (Registrar-General, 1935-8, 1944, 1954). The major factor in bringing about this fall in mortality has been the advance in the treatment of peritonitis. It is generally accepted that further reduction in the mortality will be achieved only if earlier diagnosis allows surgical intervention before the onset of complications due to the spread of infection beyond the appendix.

Recent articles on appendicitis have focused attention on treatment, and we thought that a study with emphasis on diagnosis might be of value. Few problems of diagnosis present more difficulty to the general practitioner or surgeon, and one of the main difficulties is caused by the frequency with which the onset and clinical features are atypical, especially in the very young and very old patient. As with acute pancreatitis, the greatest aid to diagnosis of appendicitis is a constant awareness of its possibility.

Material

The 549 patients who form the material for this study were admitted to one surgical charge of the Royal Infirmary of Edinburgh over a period of three years (1953-6), and in all of them the diagnosis of acute appendicitis was proved at operation. Most of the patients comprising the series were operated on by one surgeon and the remainder were under his close supervision, so that the indications for operation, the classification of the operative findings, and the operative and post-operative treatment were uniform throughout. None of the patients was treated conservatively.

Classification of Patients.—We have adopted a modified form of the simple classification used by Gilmour and Lowdon (1952), Rees (1952), and Wolff and Hindman (1952) (Table I). In our series 80% of patients were operated on

TABLE I.—Classification

Group	Condition	No. of Patients
1	Acute appendicitis	441 (80.3%)
2	Acute perforated appendicitis with localized peritonitis	25 (4.5%)
3	Acute perforated appendicitis with abscess formation	39 (7.1%)
4	Acute perforated appendicitis with generalized peritonitis	44 (8.1%)
	Total	549 (100%)

before the infection had spread beyond the appendix. In the remaining 20% infection had spread to such an extent that drainage of the area was necessary.

Age and Sex Incidence (Table II).—None of the patients in this series was under the age of 12 years; 64% were under 30 years of age and only 5% were 60 or more. Table III shows that with advancing age the percentage of patients with simple acute appendicitis diminishes whereas the percentage with perforated appendicitis increases. For example, the ratio of complicated to simple appendicitis was 1:8 in the second decade, 1:2 in the fifth decade, and 3:2 in patients of 60 and over.

TABLE II.—Age Incidence

No. of Patients	Age Group in Years						Total
	12-19	20-29	30-39	40-49	50-59	60+	
Male ..	75	73	30	25	27	13	243
Female	111	93	35	29	22	16	306

TABLE III

Age	No. of Patients	Group 1	Groups 2, 3, 4
10-19 ..	186	164 (88%)	22 (12%)
20-29 ..	166	148 (89%)	18 (11%)
30-39 ..	65	52 (80%)	13 (20%)
40-49 ..	54	38 (70%)	16 (30%)
50-59 ..	49	28 (57%)	21 (43%)
60+ ..	29	11 (38%)	18 (62%)

Duration of Symptoms

Comparison of the average duration of symptoms in the different age groups is misleading because in the older patient a history of relatively mild symptoms for several days or even several weeks is common. To obtain a more accurate comparison and to avoid overestimation in the older age groups, the patients in the series with vague symptoms of long duration were assessed only according to the duration of their acute symptoms. Only 451 out of the 549 patients are included in the analysis of the duration of symptoms.

Table IV shows that 31.9% of the 451 patients were admitted to hospital within 12 hours of the onset of symptoms, 63.9% within 24 hours, and 91% within 48 hours.

TABLE IV.—Duration of Symptoms in 451 Cases

	12 Hours	24 Hours	48 Hours	Over 48 Hours
All groups { All ages	144 (31.9%)	288 (63.9%)	410 (91.0%)	41 (9%)
Under 50	132 (34.9%)	249 (65.9%)	351 (92.9%)	26 (6.9%)
50+	12 (16.4%)	39 (53.4%)	59 (80.9%)	15 (19.1%)
Group 1 { All ages	124 (36.9%)	246 (73.2%)	331 (98.5%)	5 (1.5%)
Under 50	117 (31.0%)	220 (58.0%)	299 (99.6%)	1 (0.4%)
50+	7 (19.4%)	26 (72.2%)	32 (88.8%)	4 (11.1%)
Groups 2, 3, 4 { All ages	20 (17.4%)	42 (36.5%)	79 (68.7%)	36 (31.3%)
Under 50	15 (19.4%)	29 (37.2%)	50 (65.4%)	27 (34.6%)
50+	5 (13.5%)	13 (33.1%)	29 (75.7%)	9 (24.3%)

In patients of 50 and over 19% had symptoms for longer than 48 hours and only 16% for less than 12 hours, while in those under 50 years of age the comparable figures are 8% and 35%.

Although perforation of the appendix is uncommon in the first 24 hours of illness, it occurred in 14% of the patients admitted within the first 12 hours and in a similar percentage admitted in the first 24 hours.

Clinical Features (Table V)

Abdominal Pain.—All the patients in the series complained of pain, although this was not always the first symptom. The classical textbook description of central pain moving to the right side of the abdomen was given by only 39% of all patients, more commonly by those in whom the infection had spread beyond the appendix (50%). This is to be expected if it is accepted that the localizing pain is due to peritoneal irritation, but of patients with spread of infection beyond the appendix 21.3% gave a history of central abdominal pain only.

Vomiting.—This was more common in patients in whom the appendix had perforated (74%).

History of Previous Attacks.—This was more common in patients in whom the appendix was intact, suggesting that a perforation is more apt to occur in the first attack of appendicitis.

Urinary Symptoms and Diarrhoea.—Twenty per cent. of all patients complained of urinary symptoms and 16% complained of diarrhoea.

TABLE V.—Symptomatology and Physical Signs

Clinical Feature	Group 1 (441 Cases)	Groups 2, 3, 4 (108 Cases)
Abdominal pain:		
Classical ..	161 (36.5%)	54 (50.0%)
Central only ..	82 (18.6%)	23 (21.3%)
R.I.F. ..	133 (30.0%)	25 (23.0%)
Loin ..	31 (7.0%)	6 (5.5%)
Vague ..	34 (7.7%)	—
Tenderness maximal in right lower quadrant	438 (99.3%)	108 (100.0%)
Vomiting ..	231 (52.4%)	80 (74.1%)
History of previous attacks ..	172 (39.0%)	28 (25.9%)
Urinary symptoms ..	85 (19.3%)	28 (25.9%)
Diarrhoea ..	62 (14.0%)	25 (23.1%)
Foul breath and furred tongue ..	284 (64.4%)	106 (98.1%)
Muscle-guarding ..	338 (76.6%)	102 (94.4%)
Rebound tenderness ..	228 (51.7%)	78 (71.6%)
Rovsing's sign ..	15 (3.4%)	5 (4.6%)

Foul Breath and Furred Tongue.—These very constant findings were found in 64% of the group I patients and in 98% of those in the other groups.

Muscle Resistance.—Involuntary resistance by the muscles to palpation of the right lower quadrant of the abdomen was elicited in 76% of the 441 patients comprising group 1. In all but six of the 108 patients in the other groups, muscle resistance was marked, and where there was generalized rigidity of the abdominal muscles from diffuse peritonitis the right lower quadrant could usually be identified as the site of maximum resistance.

Blumberg's Sign.—Rebound tenderness was elicited in approximately 70% of all cases: 338 out of the 441 patients in group 1 and 78 of the 108 patients in the other three groups.

Rovsing's Sign.—Pain in the right iliac fossa on deep palpation of the left iliac fossa was elicited in only 3.5% of all patients.

Temperature (Table VI).—We regarded an axillary temperature of 98.6° F. (37° C.) as being the upper limit of normal. In 70% of the patients the temperature was not

TABLE VI.—Temperature (Axillary)

Age	Groups	No. of Patients	98.6° F. (37° C.) or Less	—100° F. (—37.8° C.)	>100° F. (>37.8° C.)
Under 50 years	{ 1, 2, 3, 4	402	314 (78.1%)	71	17
50 and over	{ 1, 2, 3, 4	69	23 (33.3%)	31	15
	{ 1, 2, 3, 4	39	30 (76.9%)	8	1
	{ 1, 2, 3, 4	39	18 (46.2%)	11	10

raised: 78% in group 1 and only 38% in the other groups. Of the patients under 50 years of age in whom the infection was localized to the appendix, 78% showed no rise of temperature; and in those in whom the infection had spread, 33% showed no rise. However, of the patients aged 50 and over in whom the infection had spread beyond the appendix, no less than 46% presented with normal or subnormal temperatures.

Pulse Rate (Table VII).—We regarded a pulse rate of 90 a minute as being the upper limit of normal. In patients

TABLE VII.—Pulse Rate

Age	Group	No. of Patients	90 min. or Less	91-110 min.	>110 min.
Under 50 years	{ 1, 2, 3, 4	402	266 (66.2%)	106	30
50 and over	{ 1, 2, 3, 4	69	22 (31.9%)	29	18
	{ 1, 2, 3, 4	39	25 (64.1%)	13	1
	{ 1, 2, 3, 4	39	14 (35.9%)	21	4

of all ages there was no increase in pulse rate in 33% of those in whom the infection had already spread beyond the appendix.

Treatment

The operative treatment was standardized, and the surgical approach was usually through a muscle-splitting incision in the right iliac fossa which was converted into a muscle-cutting incision if access was inadequate. In a few cases where the diagnosis was in doubt a right para-

median incision was preferred. If drainage of the peritoneal cavity was considered necessary, the drain was brought out through the iliac incision except in cases of diffuse peritonitis, in which the drain was brought out through a separate midline stab incision. If diffuse peritonitis was present, intravenous fluid therapy and gastric aspiration were instituted as soon as the patient returned from the theatre. In the patients in whom there was gross contamination of the wound at operation, anti-gas-gangrene serum and anti-tetanic serum were given in addition to parenteral administration of penicillin and sulphonamides, or penicillin and streptomycin.

Mortality

There were six deaths in the series, an overall mortality of 1.08% (Table VIII). With one exception these were

TABLE VIII.—Deaths

Sex	Age	Group	Days of Post-operative Survival	Necropsy Report
M	52	1	8	Disseminated sclerosis and renal infection
F	42	4	5	Suppurative bronchopneumonia
M	56	4	10	Pulmonary embolus
M	61	4	8	Cerebral haemorrhage
M	69	4	17	Bronchopneumonia
M	79	4	6	Rheumatic heart disease and cardiac failure

patients in whom appendicitis had given rise to peritonitis. The mortality for group 1 patients was 0.23%. The five patients in group 4 who died had complicating medical conditions which contributed to the fatal outcome. No patient died from appendix abscess despite the fact that there were almost as many patients in group 3 as in group 4.

Discussion

Appendicitis must be suspected in all patients, irrespective of age, who present with abdominal pain. Acute pain in the right lower quadrant of the abdomen is an indication for urgent operation and denotes probably that the infection has already spread beyond the appendix. That right iliac fossa pain is not associated with central pain in over 25% of all patients may be explained by the threshold for intestinal pain felt in the umbilical region being higher than that for pain produced by irritation of the parietal peritoneum (Brown, 1951).

Vomiting is a characteristic feature of the obstructive type of appendicitis, and if a marked feature the probability is that the appendix has already ruptured. Looseness of the bowels is another indication that operation should not be delayed, and as a symptom it is particularly dangerous because it may be mistaken for enteritis. A dose of castor oil may not be so important in causing rupture of the appendix as in causing delay in diagnosis, with consequent rupture of the appendix (Brown, 1951). The incidence of bowel and urinary symptoms was higher in the patients in whom the infection had spread beyond the appendix.

The characteristic breath fetor and furred tongue we consider invaluable signs. If these are absent the diagnosis of appendicitis is in doubt.

Tenderness in the right lower quadrant of the abdomen, the point of maximal tenderness varying with the position of the appendix, makes the diagnosis almost certain, and operation should not be delayed. Marked muscular resistance is a late sign, and its degree is not reliably related to the pathology of the underlying appendix. It is negligible in the elderly and in the obese subject.

Mason Brown (1956) has pointed out the unreliability in children of Blumberg's sign of rebound tenderness, as the abrupt removal of the hand from the child's abdomen is in itself a cause for alarm. In adults, however, we consider this a most reliable diagnostic sign. Rovsing's sign is uncommon, and we agree with Davey (1956), who found it in 1.7% of patients, that it is of little diagnostic value.

Brown (1951) stressed the fact that temperature and pulse rate were completely unreliable in the early diagnosis of

acute appendicitis, and he stated that the majority of patients who presented with a raised pulse rate had perforated appendices at operation. However, we found that increase in pulse rate was more common in simple acute appendicitis than in appendicitis with spread of infection. Thirty per cent. of our patients had an axillary temperature exceeding 98.6° F. (37° C.), and 40% had a pulse rate greater than 100 a minute. In the group of unperforated appendicitis, 78% had no rise of temperature and 66% no increase in pulse rate. In the patients in whom the infection had spread beyond the appendix, 38% had no rise of temperature and 33% no increase in pulse rate. In patients aged 50 and over we found that 70% of those with pyrexia had perforated appendices at operation. We agree with Wakeley and Childs (1950) that "absence of pyrexia has no significance but when slight pyrexia is present operation is urgent," and, we would add, particularly in the elderly patient, in whom elevation of temperature is usually associated (70%) with the complication of spread of infection beyond the appendix.

The difficulty of early diagnosis of appendicitis remains. Ascroft (1954) has shown that the error tends to be in the direction of over-diagnosis, other diseases being diagnosed as appendicitis five times as often as appendicitis is diagnosed as something else. This mistake is three times more common in females than in males. Since early diagnosis depends on the assessment of slight clinical findings, there will be unnecessary admissions to hospital and the sacrifice of many normal appendices. However, this is no error of judgment for which the doctor should be taken to task, but rather "should be regarded as an insurance premium which is paid lest some serious catastrophe arise" (Mason Brown, 1956).

The proportion of atypical cases is invariably highest in patients in whom the appendix is perforated, and "it may be that gangrene, rupture, or abscess occurs because the patient has presented atypically" (Boyce, 1954). That the disease presents atypically at the extremes of age is well known, but it is not so well recognized that it may present atypically in the middle-aged.

Appendicitis appears to be a more fulminating process in the elderly than in the young, possibly because vascular occlusion occurs more readily, thus increasing the liability to early rupture (Maes *et al.*, 1934). The higher incidence of perforated appendicitis in the older patient—47.4% of patients aged 50 and over as compared with 16.5% of those under 50—is probably attributable to the more fulminating disease and not to delay in hospitalization, although it would appear that the diagnosis tends to be delayed in the older patient (Table IV).

Perforation of the appendix had already occurred in 62% of patients aged 60 and over. This figure corresponds closely to that given by Wolff and Hindman (1952), who found perforation in 58% of patients over 60, and by Officer Brown (1934), who found perforation in 54% of patients over 50. In young children, Abel and Allen (1950) and Mason Brown (1956) estimated the incidence of complicated appendicitis to be in the region of 40%. It would appear, therefore, that the liability to perforation is greater at the extremes of age.

That it is a dangerous policy to observe a patient while one "makes up one's mind" is apparent from the fact that the appendix had already perforated in 14% of patients admitted within 12 hours of the onset of symptoms and in a similar percentage of those admitted in the first 24 hours. Delay of even a few hours is more perilous in the elderly, as is shown by the fact that, although a smaller percentage (53) of patients aged 50 and over arrived in hospital on the first day of illness, as compared with 66% of those under 50, the incidence of perforation was twice as common in the older age group (18% as against 8%). Although patients of 50 and over made up only 13.5% of the admissions on the first day of illness, they accounted for 31% of the perforations occurring within 24 hours of the onset of symptoms.

The overall mortality in this series of 1.08% is little different from the figure quoted by Gilmour and Lowdon (1952) for a comparable series of 535 patients (1.1%). All

the deaths were in patients over 40 years of age, and the Registrar-General's figures for 1954 showed that 75% of the deaths from appendicitis occurred in patients over 40. In patients in whom the infection was confined to the appendix there were no deaths attributable to the appendicitis in the series of Gilmour and Lowdon (1952) or in that of Rees (1952), and the death in our series was due to ascending renal infection complicating disseminated sclerosis. We may conclude, therefore, that, with present-day accepted treatment based on established surgical principles, the mortality from uncomplicated acute appendicitis is at the irreducible minimum for any surgical operation.

The avoidable mortality is in patients in whom the infection has spread beyond the appendix. The combined mortality rate for our groups 2, 3, and 4 was 4.6%, which is similar to the 4.5% given by Gilmour and Lowdon and by Rees. Five of the six deaths in our series occurred in group 4, giving a mortality for diffuse peritonitis from appendicitis of 11.4%. This figure corresponds to that of Gilmour and Lowdon of 11.4% and to that of Rees of 12%, and represents a great improvement over the 30-40% mortality in similar patients 30-40 years ago.

When a diagnosis of appendicitis is made the treatment of choice is early operation. In this manner only will an increasing number of patients be operated on while the infection is still limited to the appendix.

All our patients with appendix abscess were treated by early operation. Rees found the greatest mortality in this group (7-10%), and stated that this was the hard core perpetuating the high mortality in appendicitis. With modern antibiotics we believe that it is safer to drain appendix abscesses. If the appendix presents easily at operation, we believe that it should be removed; if not, a large drain is inserted and appendicectomy carried out three months later. Furthermore, the diagnosis of perforation of the appendix is notoriously inaccurate, and without operation it is impossible to be certain whether a localized mass in the right iliac fossa is an abscess or omentum wrapped round an intact appendix, and whether the disease is spreading or localizing (Boyce, 1951). Rees has called the appendix abscess "a veritable surgical wolf in sheep's clothing."

We consider that there is a real danger that antibiotics may lead to a lesser sense of urgency in diagnosis and to the abandonment of established principles in the treatment of appendicitis. Pulaski *et al.* (1948) expressed the fear that there was a tendency to institute chemotherapy and antibiotic therapy without regard to the fundamental principles of surgical management. In early uncomplicated appendicitis there is no doubt that antibiotics are no substitute for immediate appendicectomy. With regard to complicated appendicitis, Boyce (1951) stated that it is "against all reason to argue that it is safer to permit the escape of infection from a perforated viscus to continue than it is to carry out a properly planned and skilfully executed surgical procedure," and we believe that operative treatment should be carried out provided hospital facilities are available.

If sound surgical principles are followed, details of surgical technique are of little importance, except perhaps for the question of drainage of the peritoneal cavity. Shipley and Bailey (1932) advised strongly against the indiscriminate use of drainage tubes in the peritoneal cavity because they felt that their presence promoted the formation of adhesions and increased the liability to intestinal obstruction. Bancroft (1920) reported an incidence of incisional hernia of 15% after drainage, as compared with 1.9% after wound closure without drainage. However, the complications of intestinal obstruction and incisional herniae may well be the results of the initial infection and may not be due entirely to the presence of a drainage tube.

In diffuse peritonitis from appendicitis argument over the question of drainage still arises. Wakeley and Childs (1950) advocated cleansing the peritoneal cavity by suction and then draining the superficial wound only after closure of the peritoneum. We have drained the peritoneal cavity through the gridiron incision or sometimes through a separate stab incision in the midline, the drain being

shortened on the third and removed on the fifth post-operative day. We have no doubt that the immediate institution of intravenous drip therapy and gastric suction in every patient with peritonitis is an invaluable procedure. This anticipates the development of paralytic ileus and prevents distension of the bowel, thus promoting the recovery of normal peristalsis.

Summary

A review is given of a series of 549 patients with appendicitis: 64% were under 30 years of age and 5% were over 60.

No patient was treated conservatively. In 80% the operation was carried out before the infection had spread beyond the appendix, but in the remaining 20% drainage of the area was necessary. It was found that the incidence of complicated appendicitis increased in the elderly patient, and that this was not solely due to delay in diagnosis but also the more fulminating process in the elderly.

The classical textbook description of appendicitis was not often met with, and the frequency of atypical presentation is stressed. Emphasis is made on the importance of suspecting appendicitis in all patients who present with abdominal pain irrespective of age.

The characteristic breath fetor and furred tongue are important signs, and if associated with much vomiting or diarrhoea suggest that the infection has already spread beyond the appendix.

Local tenderness over the inflamed appendix, usually in the right quadrant of the abdomen, is the most important sign; it may even be the only sign, but, on the other hand, it may be absent in the very obese or elderly patient.

No reliance should be placed on the pulse rate or temperature for the early diagnosis of appendicitis. If there is a rise of temperature in the elderly patient, the infection has almost certainly spread beyond the appendix.

If operation is carried out while the infection is confined to the appendix the mortality is at the irreducible minimum for any surgical procedure. In cases in which appendicitis has progressed to abscess formation, we believe it is safer to operate and to drain the abscess. If the appendix presents, it is removed, but if appendicectomy is difficult and would require some dissection the abscess is drained.

It is emphasized that early diagnosis and prompt surgical treatment are still the most important principles in dealing with this, the most common surgical emergency.

We are indebted to Mr. W. A. D. Adamson, surgeon-in-charge, the Royal Infirmary, Edinburgh, for allowing us to treat these patients and for his help with this paper.

REFERENCES

- Abel, W. G., and Allen, P. D. (1950). *Ann. Surg.*, 132, 1093.
- Ascroft, P. B. (1954). *Arch. Midx Hosp.*, 4, 182.
- Bancroft, F. W. (1920). *J. Amer. med. Ass.*, 75, 1635.
- Boyce, F. F. (1951). *Ann. Surg.*, 133, 428.
- (1954). *Ann. Intern. Med.*, 40, 669.
- Brown, C. J. Officer (1934). *Med. J. Aust.*, 2, 407.
- Brown, F. R. (1951). *Brit. med. J.*, 1, 136.
- Brown, J. J. Mason (1956). *J. roy. Coll. Surg. Edinb.*, 1, 268.
- Davey, W. W. (1956). *Brit. med. J.*, 2, 28.
- Gilmour, I. E. W., and Lowdon, A. G. R. (1952). *Edinb. med. J.*, 59, 361.
- Maes, U., Boyce, F. F., and McPetridge, E. M. (1934). *Surg. Gynec. Obstet.*, 58, 32.
- Pulaski, E. J., Spicer, F. W., and Johnson, M. J. (1948). *Ann. Surg.*, 128, 46.
- Rees, W. S. (1952). *Lancet*, 2, 1199.
- Registrar-General (1935-8, 1944, 1954). *Statistical Review of England and Wales*.
- Shipley, A. M., and Bailey, H. A. (1932). *Ann. Surg.*, 96, 537.
- Wakeley, C., and Childs, P. (1950). *Brit. med. J.*, 2, 1347.
- Wolff, W. I., and Hindman, R. (1952). *Surg. Gynec. Obstet.*, 94, 239.