An attempt was therefore made to determine whether the chest radiographs of the five patients (seen between 1952 and 1956) were considered normal by those accustomed to interpreting chest radiographs. This investigation is reported in detail elsewhere (Somner and Sklaroff, 1958). The five radiographs were presented to 18 observers along with those from three similar cases encountered during 1957 (not included in the general analysis in this paper) and that of a ninth patient who presented with a paralysed vocal cord but without haemoptysis. The observers were asked to review these nine radiographs along with 60 others (in random order) from patients also with haemoptysis who had been examined by bronchoscopy to confirm or exclude a bronchogenic carcinoma.

Although we classified fewer hilar shadows as enlarged than the panel of observers as a whole, there was no evidence that our classification of "normal" hilar shadows was different from that of the majority of observers. Moreover, since the 18 observers did not need to make clinical decisions, they may have been inclined to be more positive in their assessment of abnormality. It is therefore concluded that our standard of interpretation of a normal hilar shadow was similar to that found among the group of observers.

The final diagnosis in the 93 cases was as follows: bronchogenic carcinoma, 5; hypertension, 7; bronchitis, 20; bronchiectasis, 9; mitral stenosis, 1; foreign body, 1; no cause, 50.

In view of the relative frequency of bronchogenic carcinoma in this group of cases we consider it advisable that all patients with haemoptysis, especially males over the age of 40, should be examined by bronchoscopy to exclude bronchogenic carcinoma even when the chest radiograph is normal, unless there is some pre-existing respiratory or cardiovascular disease which could possibly account for it.

Summary

In an attempt to clarify the indications for bronchoscopy, in particular where there is a possibility of bronchogenic carcinoma, an analysis was made of 1,109 patients examined during 1952 to 1956.

In 276 patients with a bronchogenic carcinoma the results of bronchoscopic examination and bronchial biopsy were analysed in relation to the radiographic appearances and the presence or absence of haemoptysis. A visible abnormality at bronchoscopy was found in 189 (68%) and histological evidence of the disease was obtained by bronchial biopsy in 168 (61%). The value of "blind" biopsy in obtaining histological evidence of carcinoma when the bronchoscopic appearances were normal is stressed. The diagnostic value of bronchoscopy was found to be related both to the type of radiographic abnormality and to its anatomical situation. The finding of an abnormality at bronchoscopy was not in general related to the presence or absence of haemoptysis, but in six patients with tumours and a normal chest radiograph haemoptysis was the presenting feature in five.

In the 118 cases of bronchiectasis reviewed it was concluded that bronchoscopy was not an essential part of the investigation unless surgical treatment was contemplated. Out of 132 cases of pulmonary tuberculosis examined by bronchoscopy an active endobronchial lesion was found in six patients with primary tuberculosis and a bronchogenic carcinoma in three.

Our thanks are due to Professor John Crofton and Dr. Christopher Clayson for permission to include those patients under their care and for help in the preparation of the paper, and to Miss Winifred A. M. Tait for secretarial assistance.

REFERENCES

- Penman, A. C. (1957). *Brit. med. J.*, 1, 401.
Sakula, A. (1955). *Ibid.*, 1, 759.
Somner, A. R., and Sklaroff, S. A. (1958). To be published.

TATTOOING IN THE SERVICE OF SURGERY

BY

NORMAN C. LAKE, M.D., M.S., D.Sc., F.R.C.S.

Honorary Consulting Surgeon, Charing Cross Hospital,
London

The practice of tattooing (Polynesian, *ta tau*, to mark) is undoubtedly of very ancient, prehistoric origin. In caves in Western Europe hollowed stones for grinding coloured clays are to be found; such colours were, and indeed in some regions still are, used for surface decoration (cp. woad in Britain), but it seems likely that, perhaps as the result of an accident, it was discovered that the implantation of the clay, or of charcoal from a fire, into a cut or puncture produced a permanent pattern which avoided frequent renewals. An early historical reference to the practice is to be found under the Mosaic Law in Leviticus, xix. 28, "Ye shall not make any cuttings in your flesh . . . nor print any marks upon you."

Early Tattooing

Although at first probably purely decorative, tattooing later acquired religious, tribal, and social significance, or in some cases was merely of horrific intent. The cult

FIG. 1.—Keloid scar tattooing on an Australian aboriginal produced by the method described in the text.

varied greatly. Thus in Polynesia it was a mark of puberty; among Arabs, infants were tattooed as a simple ornament or means of recognition; for the American Indians and the Australians it was a sign of the tribe or family; while the Kaffirs employed it as a decoration for bravery. The most extensive tattooing is found on the Laos of Indo-China, where the division into "black paunch" and "white paunch" areas depends upon the tattooing of the abdominal wall, but, in addition, the thighs, legs, and breasts are often completely covered with fantastic animal figures. In

Polynesia, too, large areas of the body were covered, whereas the Maoris concentrated on the face. Among dark-skinned races where ordinary pigment tattooing would be almost invisible the tendency is to produce pale keloid scarring by rubbing clay into the cuts forming the pattern (Fig. 1).

In Japan the practice became a high art, and the most varied and delicate colours were produced; there its original function was neither ceremonial nor symbolic, but in lieu of clothing. Apart from decorative and symbolic uses tattooing has had utilitarian employment to identify slaves and criminals; in the British Army until 1879 the letter D was tattooed on a deserter and B C on a bad character. Much more recently the suggestion has been made, and I believe carried into practice, that medical information, probably in code, might be tattooed on patients liable to attacks of coma—for example, in hypoglycaemia—or post-operatively when some unexpected or additional structure had been dealt with—for example, an appendicectomy in an operation for gastric ulcer.

There is no doubt that in this country tattooing, even of soldiers and sailors, among whom it was very popular not so long ago, is on the decline, but I was surprised to learn, by personal acquaintance with one, that the "young bloods" of late Victorian times were often extensively and expensively tattooed. The scene in this case, in a greyish-green monotone, was a classical one, reminiscent of a Leighton painting, extending all over the torso. It is the only tattoo in which I have seen shading and shadow most realistically represented. The tattooist evidently deserved the 250 guineas which he was paid in the 1890's for work occupying over three weeks. Other ambitious tattoos include a hunting scene in full colour around the torso of a woman (for tattooing is not exclusively restricted to males) excellently executed, a picture on the back of a doctor representing a tiger stalking through the jungle, which was a real work of art, and a large tropical butterfly on the chest of an officer, a most exquisite piece of work (Japanese) in which colours rarely seen, such as purples, blues, greens, yellows, etc., were beautifully combined. I have come across a large number of quite attractive tattoos of rather less extent which nevertheless displayed a good deal of artistry and a fairly wide range of colouring from the common red through the spectrum to the rare pure blue and purple.

But most tattoos are much less skilful: we are all well acquainted with the entwined hearts, the clasped hands, the initials of the loved one or the announcement "I love Kate," the regimental and other crests, flags, ships, the female form, and so on. Many of these are in the two common colours blue-black (Indian ink) and red (cinnabar); the materials used by the professional tattooists for the other colours are kept a close secret, and so it has been necessary to experiment with some of the modern synthetic dyes in an attempt, not entirely successful, to discover what is available.

In the surgical as opposed to the ornamental uses of tattooing the main object is, of course, a cosmetic one, and it can be employed as an adjunct to or a substitute for plastic surgery. It is, however, sometimes used therapeutically as a treatment for certain skin diseases, especially pruritus (Turell, 1941). Several papers have been published, chiefly in the United States (Cantor, 1943), on this aspect of tattooing, but the present article is concerned only with its cosmetic use.

Colours

Of the colours already noted, the ones most needed for this purpose are black and brown for hair, red for lips, pink and brown for skin, dark blue for beard stubbling, and white for tattooing down other colours.

The material used for "black" tattooing is some form of carbon, lamp black, Indian ink, but, as is well known, owing to the selective absorption of incident light by the overlying tissues this really gives a deep blue-black result, which is a poor match for black hair. Remembering that the intensely black patches of a melanoma are seen under the microscope to be produced by a dark umber pigment, I have used a combination of the dye "golden umber" with a little carbon and thus achieved a better approximation to a true black. The common red pigment is cinnabar (mercuric sulphide), sometimes called Chinese red; this has the virtue of being permanent, whereas some of the synthetic dyes tend to become faint and even to

FIG. 2.—Photomicrograph of section of skin tattooed with cinnabar. In a black-and-white reproduction the refringent pigment particles are difficult to distinguish; some are indicated by arrows.

FIG. 3.—Higher magnification of Fig. 2 to show perivascular collection of pigment.

disappear entirely in the course of time. It is, however, stated that mercury poisoning has followed the tattooing of large areas, but I have no record of this. In personal trials it was found that after 15 years "irgalite red 36" and "lake fast rubine" had disappeared, while "red oxide R. 93A" and "irgalite orange P" remained, although somewhat reduced in intensity. Of the brown pigments, both "iron oxide" and "golden umber" were reasonably permanent. The most satisfactory and permanent of several "whites" tested proved to be titanium oxide when used for tattooing out and reducing colour. It is worth noting that the chromium greens, yellows, and reds all caused rather violent immediate reactions; the "red chrome R 5" actually sloughed out, but this was possibly due to a personal idiosyncrasy.

All tattoos tend to decrease in intensity with time. For the first few weeks this is fairly rapid and may be due to transference of some of the pigment to neighbouring lymph nodes, to loss of pigment which reached no further than the epidermis—as pointed out by Matthews (1947)—and possibly to a certain amount of local fibrosis. Later there is a slower decrease accompanied by some local diffusion of the pigment, so that the margins of the tattooed area become less distinct; as mentioned above, with some materials the dye itself seems to be slowly absorbed or decolorized.

The depth to which the pigment is carried varies with the technique and the instrument used. The permanent colour is always in the dermis or subdermis, as is shown in the photomicrograph (Figs. 2 and 3) of a section in which the pigment from a red (cinnabar) tattooed area appeared as refringent particles in both these situations, the deeper masses being aggregated around the vessels. Without colour reproduction it is difficult to distinguish the particles, some of which have therefore been marked with arrows. Dr. Shore has kindly reported upon this section as follows: "There are many pigment granules in the dermis, all apparently within the cytoplasm of phagocytic cells. These pigment-containing cells are scattered fairly evenly in the middle dermis, but denser collections are present in relation to the small arterioles. No pigment is seen in the dermal papillae or in relation to the secretory portion of the sweat glands." This tattoo was of many years' duration; the particular technique used is unknown.

Antiseptic Precautions

Although some of the professional tattooists nowadays make a perfunctory application of some antiseptic to the skin and the needles, for the most part, especially in seaports abroad, no sort of antiseptic precautions are taken. The surprising thing is that so few seem to go grossly septic, but when they do the result can be both serious to the subject and disastrous to the tattoo. In one case seen many years ago the heart-rending message "I LOVE KATE" was outlined in pustules down the forearm; some of the pigment particles came away in the dressings which were applied, and the final result was the persistent message in a bizarre mixture of white scar and patchy pigment. It is well known that syphilis, tubercle, and other diseases have been inoculated by the tattooing needles. In surgical practice the usual aseptic technique will of course be applied.

Small areas can be tattooed in relatively insensitive parts without any anaesthetic, the discomfort being minimal; in surgical work, however, for large sensitive areas either a local or a general anaesthetic is necessary. If the former it is well to massage any excess of fluid away before the needles are used, otherwise it is difficult to judge depth and intensity. Many of the decorative tattoos are done, with or without consent, while the subject is under the influence of alcohol—evidently a good anaesthetic for this purpose.

Originally all tattooing was, of course, done by hand, employing a small bundle of bone (later metal) needles or points. This hand method can still be employed by the surgeon if the area to be covered is small; usually, however, a vibratory machine is used. There are several patterns—electromagnetic, solenoid, synchronous, or even mechanical clockwork. The needle points, 6 to 10 in number, are usually protected by an adjustable guard which enables the depth of penetration to be varied according to the thickness of the dermis and epidermis at varying sites. In some types of machine the actual needle stroke is controllable. About 2 to 3 mm. penetration is an average. This part of the instrument must be sterilizable. The powdered pigment is mixed with a little water to form a paste in a container which can be placed in the ordinary instrument sterilizer. I have not found this alters the colour of the pigments I have used, but Matthews (1947), for fear of such a change, sterilizes his pigments with absolute alcohol. The addition

of a small quantity of penicillin to the paste before use may be an additional precaution, unless the patient is found to be sensitive to it.

Technique

The area to be tattooed is prepared and covered with the paste through which the needles work. Judging the intensity of the coloration is a great difficulty; there are probably few surgeons who get enough experience to be able to do this without a lot of trial and error. From time to time all the paste, to which a certain amount of blood is now added, is wiped off and oozing of blood arrested so that the depth of colour can be estimated. Even so, it is not at all easy to be certain of the final result; for the multiple punctures and hyperaemia of the skin tend to obscure the pigmentation. Furthermore, allowance must be made for the early fading noted above, a factor which is always rather unpredictable. In some cases it may be considered wise to call in the help of a professional tattooist, whose greater experience will enable him to achieve a nearer colour match. I am told that some professionals are now turning their attention to cosmetic as distinct from decorative tattooing. It is certainly better to underdo rather than overdo the process, for at a subsequent session more pigment can be tattooed in, but it is much more difficult, although not always impossible, to reduce the intensity.

During the tattooing, the skin should be kept stretched and steady, but care must be taken not to distort it. The area to be covered must be marked out, and as such markings may be obscured when the paste is applied it is sometimes of value to use a template. This is particularly so when a definite line has to be followed—for example, an eyebrow or the red lip margin. Fig. 4 shows an easily made template for tattooing the "cupid's bow" on the lip; it is of sheet metal; if the subject indicates her wishes with a lipstick the template can be fashioned to suit (Fig. 5). A second backing sheet (with a notch for the frenum if this is present) fits into the bucco-labial recess to give even counter-pressure. Eyebrows, too, can be marked with grease-paint to the patient's satisfaction and the curve copied

FIG. 4.—Sheet-metal templates used for tattooing red border of lip.

FIG. 5.—Showing use of templates. The notch in the upper margin is to clear the columella if the lip is very shallow.

on a template to guarantee accuracy. Fig. 6 is a print from a colour photograph of a case where an ugly irregular scarred lip with practically no red margin followed a hare-lip operation in infancy. The result shown, although the reproduction unfortunately loses considerably from lack of colour, was achieved by using a template of the sort described. The

exact tinting of skin grafts in conspicuous positions to match their surroundings is difficult; although improvement is usual, perfection is rare.

Many other possible indications for tattooing will present themselves from time to time. Thus beard-stubbling can be imitated in dark-haired men with pallid facial grafts or scars; nails (painted or otherwise) can be simulated after amputation of the ends of digits, usually fingers but nowadays also toes; while, following plastic reconstruction of the breast, if the nipple has not been transplanted, a nipple and areola tattooed in may mollify the patient's embarrassment. "Some patients are extremely sensitive about sacrificing the umbilical pit" (Max Thorek); this surgeon trans-

FIG. 6.—Print from colour photograph of a case of hare-lip described in the text in which templates were employed. Before tattooing practically no red margin was visible. The colour original shows an excellent colour match (without make-up). The pigment used was a mixture of "red oxide R.93A" and "irgalite orange P."

plants the umbilicus when performing plastic reduction of the abdominal wall, but this contingency could alternatively be dealt with by tattooing. An unusual demand came some years ago from a Chinese lady who, as the result of a motor accident, was left with a very conspicuous wide scar running right through the hair margin some distance into the scalp. This was an aesthetic calamity, for the framing of the face in a perfect unbroken square of intensely black hair is, so I was

Removal of Tattoo Marks

As already mentioned, complete tattooing out is almost impossible, but a reduction of colour intensity may be accomplished by over-tattooing with titanium oxide. This intensely white pigment appears from our tests to be remarkably permanent and has greater "covering" value than other materials tried.

Removal of tattoo marks is often demanded, sometimes on account of a change in social status, sometimes because the tattooing was accidental and unwelcome—for example, after road and mining accidents with tar or coal-dust inclusion, or bomb incidents, which produced some rather fantastic examples, such as the pattern, in negative, of an ornamental iron railing on the bare back of a lady who was leaning against it at the time of the explosion. Then there is the embarrassment which a transfer of affection to "George" occasions when the tattoo declares that the lady "Loves Fred" (an actual example that required complete excision and skin-grafting to eliminate).

A linear accidentally tattooed scar can obviously be excised completely, but in most cases the area involved is too extensive to be dealt with in this way. It will be remembered that the pigment particles lie in the dermis and subdermal tissues, so that no superficial removal such as is

produced by scarification (Kromayer, 1905), whether by sand-paper or the wire-brush technique (Eller and Eller, 1956), can be expected to be entirely satisfactory, however good such methods may be for surface lesions such as acne scars. Dermabrasion (Pegum, 1957) of this type would have to be carried into the deep layers of the dermis, allowing epithelization only from the sparsely scattered hair roots and skin glands, a very slow process which not only leaves a very conspicuous scarred area but often results in ugly keloid scars. The only satisfactory method is by complete excision followed either by sliding flaps or skin-grafting; in many areas a split-skin Thiersch graft is acceptable, but in others a full-thickness or pedicle graft is required. The moral is that, once a tattoo is in, the subject can never more be "without mark or blemish."

Conclusion

Apart from the chance of transferring diseases, the risks of tattooing are those of infection, avoidable in surgical practice; of local reaction to the pigments used, sometimes idiosyncratic, as in the case of chromic salts producing sloughing; of general poisoning, as recorded very rarely with mercuric salts; and, equally infrequently, of a much-delayed reaction. These late cases may occur many years after an uneventful tattoo (Bonnell, 1956)—especially, apparently, with the green chromic salts, if the subject comes into intimate contact with other chromic-containing material, such as paints, to which he appears to have become sensitive.

All these risks are, however, quite remote and should not debar the proper employment of an excellent method of overcoming many disfigurements.

Summary

A brief account of the historical development and decorative application of the common forms of tattooing is given and certain ambitious tattoos are described.

The pigments which may be employed in the surgical uses of tattooing are discussed, as is also their suitability for this purpose and their distribution in the skin.

The technique of tattooing by hand and mechanical methods is described, with an indication of possible surgical applications and of the assistance afforded in some instances by the use of templates.

The complications and risks of tattooing, both immediate and remote, are considered, and the long-term changes which occur are described.

The methods which may be employed in the removal of unwanted tattooing are discussed and their limitations indicated.

I am indebted to Mr. Clifford Evans, F.R.C.S., whose chemical knowledge was of great assistance; to Dr. J. H. Shore and the department of pathology, Charing Cross Hospital Medical School, for the pathological report and the photomicrographs; and to Miss Turnbull, of the photographic department of the hospital, for the photographs.

REFERENCES

- Bonnell, J. A. (1956). *Proc. Roy. Soc. Med.*, 49, 823.
- Cantor, A. J. (1943). *Amer. J. Dis. Skin*, 10, 254.
- Eller, J. J., and Eller, W. D. (1956). *Med. Tms (N.Y.)*, 84, 158.
- Kromayer, (1905). *Derm. Z.*, 12, 26.
- Mathews, D. N. (1947). *Proc. Roy. Soc. Med.*, 40, 861.
- Pegum, J. S. (1957). *Ibid.*, 50, 607.
- Thorek, Max (1942). *Plastic Surgery of the Breast and Abdominal Wall*.
- Turell, R. (1941). *Amer. J. Obstet. Gynec.*, 42, 290.

The Sir George Earle Trophy for a notable achievement in the cause of industrial accident prevention has been awarded this year to Imperial Chemical Industries Ltd. This is the third time the trophy has been awarded. In 1956 the recipients were H.M. Inspectors of Factories, and last year it was awarded to the Birmingham and District Industrial Safety Group.