Discussion

Maintenance of the circulation during cardiac arrest (together with adequate administration of oxygen) is important for two reasons. A heart which is anoxic is unlikely to resume its normal rhythm and the dysrhythmia itself is often the result of anoxia. Anoxia from cessation of the circulation of more than three minutes' duration may be followed by irreversible damage to the central nervous system. Thus the question whether cardiac massage has a beneficial effect because it provides a local mechanical stimulus to the heart or because it is a means of maintaining the circulation is not an academic one. If it can be shown that it serves the latter purpose then its importance in this respect far outweighs the former, and the operator must not be content with merely gaining a limited access by which to apply some mechanical stimulus to the heart. His exposure must enable him to achieve as efficient a pumping action as possible.

Our tracings offer evidence that cardiac massage in man is an effective means of maintaining some peripheral circulation. In one case the peripheral blood pressure during this manoeuvre was 75/30 and in the other it was 55/20 even with the pericardium closed. The blood pressure depends not only on the cardiac output but also on the peripheral resistance, so that the recording of such pressures does not provide an exact indication of the amount of the circulation. It is possible that over a short period of time intense peripheral vasoconstriction could result in such pressures being achieved with only a small cardiac output. But this consideration does not detract from the possible value of the procedure, for it is generally believed that such vasoconstriction does not involve the cerebral and coronary circulation, and is in fact a means of achieving a favourable distribution of blood to these organs. Thus it is reasonable to suppose that the recording of such pressures offers evidence that the flow of blood to the brain and heart achieved by cardiac massage is of considerable value. Reports of recovery without serious cerebral damage, following cardiac massage for several minutes (for example, Coyne, 1952; Touroff and Adelman, 1949), also support this supposition.

The shape and amplitude of the peripheral pulse depend on the volume, rapidity, and duration of cardiac ejection and the elasticity and state of contraction of the peripheral vessels. In both cases we recorded a slow rising and sustained pulse; this characteristic may be connected with the difficulty reported by others in detecting the pulse by palpation or auscultation and measuring the blood pressure by means of a sphygmomanometer (McMillan and Cockett, 1952).

Summary and Conclusions

In two cases a record of the peripheral pulse has been obtained by means of an intra-arterial catheter and an electronic manometer during the performance of cardiac massage. The records obtained offer evidence that cardiac massage is capable of maintaining some circulation. It follows that for cardiac massage to be properly performed the surgical approach must enable the heart to be firmly grasped.

We wish to thank Mr. R. C. Brock for his encouragement to prepare this report.

REFERENCES

- Coyne, F. (1952). *British Medical Journal*, 2, 1046.
 Johnson, J., and Kirby, C. K. (1950). *J. int. Chir.*, 10, 157.
 McMillan, I. K. R., and Cockett, F. B. (1952). *British Medical Journal*, 2, 1308.
 — and Styles, P. (1952). *Thorax*, 7, 205.
 Peterson, L. H., Dripps, R. D., and Risman, G. C. (1949). *Amer. Heart J.*, 37, 771.
 Touroff, A. S. W., and Adelman, M. H. (1949). *J. Amer. med. Ass.*, 139, 844.

DRUG RESISTANCE IN MALARIA

BY

T. C. HALLINAN, L.M.S.S.A.

Medical Officer, West Indies Sugar Co., Frome Estates,
Jamaica, B.W.I.

The resistance of many cases of malaria to drug treatment has long been a matter of concern to those who have to treat this disease, and so far no satisfactory solution has been found. The evidence submitted here suggests that infestation with hookworm, which is endemic in all malarious areas, or with other blood-sucking or blood-infesting worms, is a cause of malaria being resistant to drug treatment.

A chance observation in a very resistant case of malaria (*Plasmodium falciparum*) in which the parasites persisted in the blood in spite of repeated courses of treatment with proguanil, mepacrine (by intramuscular injection), and quinine, used either singly or in combination, showed that the patient was excreting hookworm in the faeces. It was observed that by the day following treatment of the patient for hookworm the malaria parasites, fever, and general symptoms had disappeared. During the next 8-9 months 12 similarly resistant cases were studied. All were shown to have hookworm infestation, and all failed to respond to anti-malarial treatment until hookworm treatment was given, when they immediately responded.

In January, 1949, a large-scale study was begun. All cases of malaria were thenceforward treated for hookworm and observations made to see whether, compared with previous years, the relapse rate fell and continued at a low level. The standard of "relapse" adopted was that any case returning with malaria within six months of the original treatment was to be regarded as a relapse and not as a new infection, unless the parasite found was of a different variety from the previous one. The drugs and methods of treatment used were the same in all the years mentioned in Tables I and II, except that in 1948 hookworm treatment was not given.

TABLE I

	1948		1949		1950		1951	
	New Cases	Re-lapses	New Cases	Re-lapses	New Cases	Re-lapses	New Cases	Re-lapses
<i>Plasmodium</i> :								
Vivax ..	8	26	3	1	5	2	4	1
Ma'ariae ..	94	217	94	24	14	6	65	7
Falciparum ..	219	712	127	37	70	9	307	44
Total ..	321	955	224	62	89	17	376	52
% relapses to new cases ..	297.5		27.7		19.1		13.86	

Results

Table I refers to proved cases of malaria in which parasites had been demonstrated. In Table II are presented the number of cases suspected of malaria clinically in which no parasites were found but other evidence—for example, malaria debris, monocytosis, etc.—was usually present. The results given in both tables show a great reduction in the malaria relapse rate and have a similar trend, though the relapse figures for 1950 in Table II are slightly higher than might be expected; this may be due to a period of four and a half months' leave, during which the diagnosis was in the hands of a locumtenent with rather less experience of malaria.

The figures for 1947 were in general similar to those for 1948.

TABLE II

	1948		1949		1950		1951	
	New Cases	Re-lapses	New Cases	Re-lapses	New Cases	Re-lapses	New Cases	Re-lapses
Cases of clinical malaria ..	472	1,302	1,175	429	1,465	977	1,486	175
% relapses to new cases ..	275.8		36.5		66.7		11.8	
No. attending at clinic ..	14,279		19,196		26,586		19,504	
No. of blood films examined ..	2,459		1,791		1,756		2,879	

Discussion

Hookworm has been especially mentioned, as it is believed to be the main cause, but whipworm has been present in most of the infested cases, and occasionally roundworm; tapeworm not at all. The hookworm infestation rate in this area is in the region of 80%.

When attached to the gut wall the hookworm is known to inject an anticoagulant into the blood stream to prevent clotting while obtaining blood. It is possible that either this anticoagulant or some other unknown substance injected at the same time has the effect of preventing the action of antimalarial drugs. It seems unlikely that the hookworm acts by absorbing the drugs while they are still in the intestine, because mepacrine and quinine when given by injection have also often failed to act until the hookworm treatment has been given.

It is possible that the hookworm treatment causes the hookworms to release their hold on the intestine, and their products may be destroyed or be eliminated from the body. The antimalarial drugs may then exert their effect before the surviving hookworms are able to regain their position and activity.

Medical Memorandum

Generalized Actinomycosis with Possible Cardiac Involvement

Actinomycosis is not uncommon in the classical sites, but general dissemination is infrequent and endocardial involvement rare. Cornell and Shookoff (1944) found only 18 reported cases with endocardial ulceration capable of causing pyaemia. In only two were valves infected; the remainder arose by direct spread from the mediastinum (11) or by embolic involvement of the myocardium (5). One of their personal cases, a clinical mitral lesion, violates any pre-mortem diagnosis of an actinomycotic valve infection: the valves were normal at necropsy, but granulation tissue had penetrated the auricle from the mediastinum.

Localized lesions usually yield *Actinomyces bovis*, a non-acid-fast anaerobe. The acid-fast aerobic organisms are pathologically less common, but more often cause generalized spread (Cope, 1938). Cases recently reported incriminate members of this group (Custis *et al.*, 1944; MacNeal *et al.*, 1946; Stokes *et al.*, 1951).

A case is reported below in which general dissemination was due to *A. bovis* of probable endocardial origin. Prolonged penicillin therapy resulted in apparent cure.

CASE REPORT

An apprentice joiner aged 18 was admitted on December 20, 1950. Eight months previously he had a left facial palsy following an injury to the homolateral mastoid region, the site of mastoidectomy at the age of 6. Lassitude, weight loss, and stabbing pains deep to the xiphisternum on bending forward and local pressure then started. Three-day periods of discomfort in the same situation were accompanied by dyspnoea on exertion. Later, "cramps" in his calves on walking preceded the appearance of tender swellings.

Examination revealed a harsh aortic systolic murmur with a split second sound and accentuated pulmonary second sound. Discrete, tender, subcutaneous swellings 2 to 4 cm. in diameter were present in the calves and thighs—some fluctuant, others firm. The feet were plantar-flexed, and dorsiflexion caused pain. Knee movements were similarly limited. Diagnosis was not aided by numerous investigations. A blood culture yielded a spore-bearing organism rejected as a contaminant. An x-ray film of the chest showed equivocal opacities at the right base; the posterior mediastinum was seen to be clear. Biopsy revealed non-specific lymphocyte infiltration and fibroblastic reaction.

The temperature ranged from 98 to 100° F. (36.7 to 37.8° C.), and the pulse 100 to 110. Bacterial endocarditis (organism not known) was diagnosed and penicillin, 250,000 units four-hourly, was administered for seven weeks. The temperature settled in six days, but the tachycardia persisted until the sixth week, when the spleen was palpable for a few days. Pain, tenderness, and the abnormal heart sounds disappeared, though some nodules persisted. For the next six weeks he remained well, but new lesions developed on the legs and forearms.

In April, 1951, the patient was admitted to the Manchester Royal Infirmary. Both firm and fluctuant nodules were present on the arms and legs; the calves felt woody. There was no fever, splenomegaly, or cardiac abnormality. *A. bovis* was isolated from a fluctuant lesion, and microscopy of the faeces showed many morphologically similar organisms. An x-ray film of the chest remained equivocal. Penicillin ("distaquaine"), 600,000 units eight-hourly, was given for 16 days, and then once daily for two months. All lesions gradually subsided. In September, 1952, he was well; an x-ray film of the chest showed no change, and there were no actinomycetes in the faeces.

DISCUSSION

The varying cardiac bruits suggested the endocardium as the focus for pyaemic spread. Whether it was on the valves, or whether it was on the mural endocardium as described by Cornell and Shookoff, could not be decided. The lungs were an unlikely focus for dissemination. The faecal actinomycetes and retrosternal pain indicated a primary oesophageal lesion, possibly spreading to involve the endocardium. The absence of mastoid inflammation made it an unlikely focus for a blood-borne endocardial infection. The fresh lesions after the disappearance of cardiac signs may have been due to the slow growth of deposits already in the tissues and not to the persistence of the cardiac focus.

Penicillin therapy should be prolonged in systemic actinomycetes infections; with small dosage, up to nine months has been necessary (MacNeal *et al.*). Chloramphenicol, however, has proved successful after four weeks in an *Actinomyces muris* infection (Stokes *et al.*). The present case suggests that with the current large doses of penicillin at least three months' treatment is necessary, though one daily injection of a long-acting product may be enough after the first month.

Finally, in a case of doubtful pyaemia an unusual spore-bearing organism from blood culture should not be lightly disregarded as a contaminant.

Our thanks are due to Professor T. H. Oliver for his kind co-operation and to the Department of Clinical Pathology, Manchester Royal Infirmary.

RONALD S. SAVIDGE, M.D., M.R.C.P.,
Senior Registrar, Manchester Royal Infirmary.

DONALD M. DAVIES, M.D., M.R.C.P.,
Physician, Bury and Rossendale Hospitals.

REFERENCES

- Cope, Z. (1938). *Actinomycosis*. Oxford Univ. Press, London.
Cornell, A., and Shookoff, H. B. (1944). *Arch. Intern. Med.*, 74, 11.
Custis, D. L., Halley, H., and Bacon, C. M. (1944). *Arch. Path.*, 38, 332.
MacNeal, W. J., Blevins, A., and Duryee, A. W. (1946). *Amer. Heart J.*, 31, 668.
Stokes, J. F., Gray, I. R., and Stokes, E. J. (1951). *Brit. Heart J.*, 13, 247.