WEIGHTS OF NEWBORN INFANTS SEVENTEENTH AND EIGHTEENTH CENTURIES COMPARED WITH THE MID-TWENTIETH CENTURY

BY

PETER E. SYLVESTER, M.R.C.S., L.R.C.P., D.C.H.

*From the Child Study Centre, Institute of Child Health,
University of London*

Clothes, suits of armour, doorways, and sometimes furniture seen in museums and ancestral homes often give the impression, perhaps incorrectly, that our ancestors were smaller than ourselves. While working on a longitudinal growth study at the Child Study Centre an opportunity presented itself to put this intriguing impression to the test, at least so far as newborn infants were concerned.

It was conceived that if a number of suitable infants' period garments could be found and measured then it might be possible to ascertain the size of the wearers

A pair of eighteenth-century muslin christening mitts. The fingers were exposed beyond the palm but covered at the back by a flap. (Victoria and Albert Museum. Crown Copyright.)

from statistics of modern infants. At the same time if it were possible to know the age of the wearers then an even more accurate and interesting comparison could be made. During a search amongst various items of infants' clothing, which included bonnets, vests, bootees, robes, and gloves, a number of christening mitts (see Fig.) were discovered. These offered the best medium for study because it was possible to date their origin—the seventeenth and eighteenth centuries—and to assess the ages of the wearers reasonably accurately from church registers of the period. The mitts had the advantage of being made of inelastic unshrinkable materials such as linen or fine muslin, which had resisted the passage of time very well.

This paper describes briefly what measurements were needed and how they were used, and raises some points for discussion before drawing conclusions.

Experiment

Eight pairs of seventeenth-century and 16 pairs of eighteenth-century mitts were measured on a flat rigid surface across the knuckle. Twice this dimension was taken as the circumference.

Most baptisms 200–300 years ago took place within the first month of life. It was possible to narrow this margin to the fifteenth day (mean 15.3 days), the median of a random sample of baptisms obtained from the eighteenth-

century parish registers of St. James in Clerkenwell* and La Patente in Spitalfields.†

The size of infants who would fit mitts of known knuckle circumference was determined by first measuring, with a cotton thread and mark, the knuckle girth of 74 newborn infants in the lying-in wards of the West Middlesex Hospital and the Chiswick Maternity Hospital. This girth was correlated with the weight at the time of measurement. Regression equations were used from these data to estimate the weights of the wearers of the mitts.

So that the final comparison could be made, it was necessary to know the weights of present-day infants on the fifteenth day of life. Information from 95 infants at the Child Study Centre made this possible. Their birth weights were correlated with their weights at 4 weeks—an age at which routine measurements were made at the centre. A relationship being assured, a graph was composed in which the mean weights at birth and at 4 weeks were plotted against the age in days. The mean weight for the sample on the fifteenth day was taken at the point where a line joining the two weights intersected that day.

(a) Mean circumference of christening mitts=9.2 cm. (range, 7.6 to 10.6 cm.).

(b) Mean hand circumference of 74 newborn infants=8.5 cm. (range, 5.8 to 9.6 cm.).

The circumference was correlated with the weights, $r=0.79$ (significant beyond the 0.001 level).

(c) Mean of the estimated weights of infants which would fit the mitts=8 lb. 1 oz. (3.657 kg.) (standard deviation, 1 lb. 5 oz. (0.595 kg.); range, 5 lb. 5 oz. (2.401 kg.) to 10 lb. 7 oz. (4.729 kg.)).

(d) Mean birth weight of 95 children=7 lb. 6 oz. (3.340 kg.) (standard deviation, 1 lb. 3 oz. (0.539 kg.)).

Mean weight of same children at 4 weeks=8 lb. 10 oz. (3.921 kg.) (standard deviation, 1 lb. 4 oz. (0.576 kg.)).

The weight at birth was correlated with that at 4 weeks, $r=0.85$ (significant beyond the 0.001 level).

Mean weight at 15 days deduced from a graph=8 lb. 1 oz. (3.657 kg.) (weights to the nearest ounce).

Result.—The estimated mean weight of 24 mitt-wearers on the fifteenth day of life was equal to the mean weight of 95 newborn infants at the same age—namely, 8 lb. 1 oz. (3.657 kg.).

Discussion

Two weaknesses in this study will arise in the mind of the reader. The first one concerns the snugness of fit of the mitts. There were no means of securing the mitts by buttons or tapes, and it was assumed that if they were to be worn safely then they must fit snugly. However, it is possible that an infant, irrespective of snugness of fit, could grasp the palmar aspect of the mitt in its normally flexed fingers, thus preventing it from falling off; indeed, this may be the reason for the design of the mitts.

The second weakness is also related to the snugness of fit, because it was thought that the garments were probably made to measure. If the mitts were part of the layette, then they must have been made to a predesigned pattern, modelled on another infant, or guessed. It was not possible to verify these points, but in order to test the theory of guesswork 29 experienced midwives were asked to draw an estimated pattern for an infant; their results were measured, and a mean mitt circumference of 10.4 cm., with a range of 6.7 to 14.5 cm., was obtained. These results are larger than the mitts, and indicate that perhaps the seventeenth- and eighteenth-century seamstresses were better guessers than modern experienced midwives. If these figures be accepted then it means there is an approximate error of overestimation of 12% (10.4 cm. being 1.2 cm. more than the presumed hand circumference of 9.2 cm. of the 95 newborn infants, whose mean weights equalled those of the mitt wearers). Therefore, if an allowance of 12% be made

*The Publications of the Harleian Society, Registers, vol. 10, A true register of all the christenings of the Parishes of St. James, Clerkenwell, 1701–1754.

†The Publications of the Huguenot Society of London, vol. 11, Registers of the Church of La Patente, Spitalfields, 1898.

for the mitt wearers then their mean weights have been overestimated by 15 oz. (0.419 kg.).

The authorities consulted (curators, keepers, and owners of costume collections) agreed that the mitts came from well-to-do families, as in the case of most preserved garments; a lace mitt—the only one in the series—from the London Museum was thought to have been used at the baptism of George III. Thus the comparison made is between infants of the upper social level in the pre- and early Industrial Revolution era and infants of a predominantly working-class population from the mid-twentieth century.

Conclusions

Newborn infants of the seventeenth and eighteenth centuries, by the methods used, are estimated to have been equal in size to infants of the mid-twentieth century. It is possible that the present-day infants may even be larger, but they are certainly not smaller than those of 200 to 300 years ago.

I wish to thank Professor Alan Moncrieff for his encouragement, Dr. Colin Hindley, who helped with the statistics, and Mr. Donald King and Miss J. Macnab, of the Victoria and Albert Museum, for assistance and access to their materials. Dr. M. Dynski-Klein and Dr. A. D. Barlow allowed me to use children under their care, for which I am grateful. In addition my thanks are due to the curators and staffs of the following museums: the London Museum; the Gallery of English Costume, Platt Hall, Manchester; the National Museum of Ireland, Dublin; the City Museum and Art Gallery, Birmingham; the Castle Museum, Norwich; not forgetting those who searched in vain.

Reports of Societies

PREVENTION AND MANAGEMENT OF TETANUS

A joint meeting of the Sections of Neurology and Experimental Medicine and Therapeutics of the Royal Society of Medicine was held on May 13 to discuss "Tetanus." Professor J. V. DACIE was in the chair.

"Tetanos Post-sérique"

Professor G. PAYLING WRIGHT (London) gave an account of some of the limitations in the common practice of passive prophylaxis against tetanus with antitoxic serum (A.T.S.) and discussed the immunological features that may underlie the occurrence of cases of *tetanos post-sérique*. He pointed out that the modern techniques of labelling foreign proteins with radioactive isotopes and fluorescent dyes now enabled their fate after inoculation to be followed with much greater precision than formerly. As a result of such studies, it was apparent that foreign proteins such as tetanus antitoxic horse globulins were removed from the circulation and destroyed in the tissues much more rapidly on their second or subsequent administration. Whereas a first dose of 1,500 units of A.T.S. might be expected to confer protection for two to three weeks afterwards, a second dose would be lost in a much shorter time. It was this accelerated destruction of antitoxin, well within the usual incubation period of the disease, that probably accounted for many of the cases of *tetanos post-sérique*.

As an instance of a failure to appreciate the phenomenon of "immunological education," with its more rapid destruction of the foreign antigen, Professor Payling Wright referred briefly to a case of tetanus which had occurred after the usual prophylactic dose of A.T.S. and which had been reported in the *British Medical Journal*.^{*} He urged that once a person had received an injection of any horse-serum antitoxin, irrespective of the disease for which it might have been given, that person should be subsequently immunized

against diphtheria and tetanus with the appropriate toxoid. Once a person had been injected with horse-serum proteases no reliance could be placed upon his protection by A.T.S. should he be subsequently exposed to risk of tetanus.

Drugs for Relieving Spasm

Dr. D. R. LAURENCE (London) discussed the need for an experimental approach in judging the value of drugs in the treatment of tetanus, stressing the meaninglessness of reporting successful results achieved by polypharmacy with drugs of unknown potency in single cases. He described an ingenious technique for producing local tetanus in rabbits injected with toxin. Their response to a standard afferent stimulus allowed him to carry out quantitative electromyographic comparisons of the potency of different drugs. The ideal drug should invariably abolish spasms without interfering with respiration or abolishing consciousness. Such a drug was not yet available, but he had found that the phenothiazine derivatives, such as chlorpromazine, were by far the best in abolishing spasms and were superior to other centrally acting muscle relaxants such as mephenesin. A new compound, acetylpromazine, was superior to chlorpromazine, but had not yet undergone clinical trial. The phenomenon of tachyphylaxis—failure to respond to a previously satisfactory dose—occurred if phenothiazine drugs were given more often than about every two hours, and very large doses might even produce excitation. In a patient whose spasms had been abolished by thiopentone, reactivation could be repeatedly brought about by stressful conversation, and the spasms disappeared again when the subject was reassured. This underlined the importance of relief of anxiety in the general management of the patient with tetanus.

Professor E. B. ADAMS (Durban) reported the results of a carefully controlled clinical trial of chlorpromazine, barbiturates, and chloral hydrate in the treatment of nearly 200 patients with tetanus admitted during the past two years to the King Edward VIII Hospital, Durban. His impression was that a mixture of barbiturates and chlorpromazine was better than either drug alone, and moreover smaller doses of each could be given. In neonatal tetanus, where the mortality is notoriously high, there was a slight bias in favour of barbiturates. Chlorpromazine could be given intravenously if necessary and was very dramatic in controlling laryngeal spasm. No side-effects had been encountered, but the total daily dose should not exceed 1 g. in adults. Supportive therapy in the form of A.T.S., penicillin, and oxygen was given, and oral feeding was used if possible.

In trying to assess the value of any new form of treatment in tetanus it was important to take into account prognostic criteria, such as the period of onset and the intensity of the spasms, he continued. The earlier the onset the higher the mortality, whereas in patients with little or no spasm death was unusual. Many of the reports in the literature were difficult to assess because these factors had not received sufficient attention. Another point that needed more study was the immediate cause of death; very often nothing obvious was found at necropsy, and it might be that some patients, especially in the younger age groups, died as a result of over-treatment. In some cases death could be attributed to respiratory complications, hyperpyrexia, or circulatory failure; exhaustion from spasms seemed an unlikely cause.

Artificial Respiration

Dr. A. CRAMPTON SMITH (Oxford) began by pointing out that new methods of treatment have really had little dramatic effect on the overall downward trend in mortality rate from tetanus. Any new therapy must therefore be put forward cautiously, and its value assessed against the known dangers of the disease. For this reason the Respiration Unit at Oxford had selected seven very severe cases whose

^{*}Littlewood, A. H. M., Mant, A. K., and Wright, G. Payling, *Brit. med. J.*, 1954, 2, 444.