mechanical origin. Indeed, attempts at a more accurate diagnosis, if they lead to delay, are misplaced, as urgent surgical relief is required in all these conditions except acute pancreatitis—and even this, after Polya gastrectomy, needs confirmation by laparotomy, because marked elevation of the serum amylase is not diagnostic. High levels may occur with duodenal leakage and with closed obstruction of the afferent loop.

Treatment

Urgent evacuation of the distended afferent loop is essential. The passage of a nasal tube might possibly achieve it, but, failing this, operation is indicated after shock has been treated with appropriate vigour. In Case 1 there was much improvement after about six hours, but operation was unfortunately deferred until further deterioration had occurred. A stomach tube was then passed by the anaesthetist and guided by the surgeon into the afferent loop, which was emptied. A nasal tube was subsequently left in the afferent limb and an enterostomy was made in the efferent limb. An alternative would have been an entero-anastomosis between the two loops, a procedure still recommended as a routine prophylactic measure by Simon (1956). Although such a stoma must divert the alkaline pancreatic juice and bile to greater or less degree, he doubts on theoretical and practical grounds whether this predisposes to stomal ulceration. While this view may not be generally acceptable, routine nasal aspiration of the afferent loop after Polya gastrectomy can scarcely be criticized.

C. F. W. Illingworth (1957, personal communication) leaves a 7-mm. nasal tube in the afferent limb in two types of case: where the closure of the duodenal stump seems insecure and where there is reason to fear temporary obstruction at the anastomosis; for example, after a difficult gastrectomy for stomal ulcer. The wisdom of this practice is evident, but duodenal leakage by no means always follows a difficult closure, nor was afferent-loop distension in the two cases described associated with any technical doubts or difficulty. There is then a case for routine aspiration of the afferent loop until emptying is proved, although it may not prevent these complications absolutely. Wangenstein (1941) described a forked catheter, one limb of which was left in each limb of the anastomosis. He used it routinely and considered aspiration of the afferent loop the more important. The need for the surgeon to number afferent-loop distension among the early complications of Polya gastrectomy is indisputable. It may be necessary to reopen the abdomen urgently on slender evidence, but, once the decision is made, treatment is simple and success would seem almost certain.

Summary

Two fatal cases of acute distension of the afferent loop after Polya gastrectomy are reported.

The course in Case 1 suggests that severe shock can develop from the distension alone and shows that there may be no early physical signs whatsoever on examination of the abdomen.

The essential clinical feature in Case 2 was persistent abdominal distension in a patient who remained in good general condition until the onset, on the eighth post-operative day, of diffuse peritonitis from gangrene from the third part of the duodenum.

In both cases the absence is stressed of an evident mechanical factor to account for the distension: atony is suspected, and may have been due to an unusually extensive incidental division of vagal fibres.

Acute afferent-loop distension may be indistinguishable clinically from the other early abdominal complications of Polya gastrectomy. The treatment is rapid resuscitation, when necessary, and urgent surgical relief of the distension.

Acute afferent-loop distension is commonly suspected to be a cause of leakage from the duodenal stump, and the possibility of avoiding both these complications by routine post-operative aspiration of the afferent loop is discussed.

I thank Professor Milnes Walker and Mr. W. M. Capper for permission to report these cases.

REFERENCES

- Burge, H. (1956). *Brit. med. J.*, 2, 1428.
 Dragstedt, C. A., Lang, V. F., and Millet, R. F. (1929). *Arch. Surg. (Chicago)*, 18, 2257.
 Eisberg, H. B. (1925). *Ann. Surg.*, 81, 926.
 Henson, G. F. (1955). *Lancet*, 1, 595.
 Hoyte, F. C., Jayne, W. H. W., and Pallister, W. K. (1957). *Ibid.*, 1, 193.
 Quinn, W. F., and Gifford, J. H. (1950). *Calif. Med.*, 72, 18.
 Roux, G., Pédoussaut, R., and Marchal, G. (1950). *Lyon chir.*, 45, 773.
 Simon, M. M. (1956). *Amer. J. Surg.*, 91, 423.
 Taylor, N. B., Weld, C. B., and Harrison, G. K. (1953). *Canad. med. Ass. J.*, 29, 227.
 Wangenstein, O. H. (1941). *Surg. Gynec. Obstet.*, 72, 257.
 West, J. P. (1953). *Surgery*, 34, 98.

FAECAL OCCULT BLOOD TESTS WITHOUT DIETARY RESTRICTIONS

BY

R. L. SMITH, B.Sc.

Biochemist, Chase Farm Hospital, Enfield, Middlesex

The most widely used tests for occult blood in faeces involve the use of benzidine. This substance is now difficult to obtain because its carcinogenic properties make its manufacture dangerous, and it seems likely that it will soon be completely unobtainable. It will therefore be necessary to use methods for detecting faecal occult blood which do not involve the use of benzidine. The orthotolidine method of Kohn and O'Kelly (1955) appears to be a suitable alternative, but when this method is used the patients must be put on a meat-free diet if false-positive results are to be avoided. This is not always convenient; special diets have to be ordered (and adhered to) and there is a delay of four days before the result is known. As Kohn and O'Kelly point out, it is quite easy to alter the sensitivity of the test, and we have tried to find a level of sensitivity such that negative results are obtained from patients who are on a normal diet and who are not bleeding into the gastro-intestinal tract.

Methods

The ages of the 49 hospital in-patients selected (27 males and 22 females) ranged from 16 to 85 years. They were on a normal hospital diet (excluding liver) and had no signs or symptoms of gastro-intestinal or other disease which might cause bleeding into the alimentary tract. Three specimens of faeces were collected from each patient, on consecutive days where possible. Each specimen was tested for occult blood by the test-tube method of Kohn and O'Kelly (1955), using five different levels of sensitivity.

Details of Method.—A piece of faeces about the size of a pea was mixed with 5 ml. of water, and the resulting suspension was boiled and cooled. A stock solution was prepared containing 4% orthotolidine "AnalaR" in absolute industrial alcohol. Five dilutions of this stock solution with 50% v/v acetic acid were made—namely, 33%, 20%, 10%, 5%, and 2%. These solutions were stored in the refrigerator and renewed monthly. Four drops of each dilution were treated with one drop of 20-volume hydrogen peroxide, and left to stand for one minute. If no blue or green colour developed, a drop of the faecal suspension was added.

The results were recorded thus :

Strongly positive (+++) Blue or green colour within 5 seconds
Positive (++) .. " " " " " 5-20 "
Weakly positive (+) .. " " " " " 20-60 "
Negative (-) .. " " " " " No blue or green colour within 60 "

Fourteen patients who showed at least a weak positive with the 10% dilution in more than one specimen were investigated as follows: (1) The three specimens were re-tested; these results were in good agreement with the first results. (2) The patients' case histories were examined for any indication of possible bleeding into the alimentary tract. (3) In two cases the patient was put on a meat-free diet and further specimens were collected. In both cases positive results were obtained, using the 20% dilution of the stock orthotolidine solution.

On the basis of these investigations the two patients mentioned, and two others in whom gastro-intestinal bleeding seemed likely, were excluded from the series. The results obtained in the remaining 45 patients (135 specimens) are shown in Table I.

TABLE I.—Results in 135 Specimens

Reaction	Dilution of Stock Solution				
	33%	20%	10%	5%	2%
Negative	4 (3%)	36 (27%)	97 (72%)	122 (90%)	131 (96%)
Weakly positive	6 (4%)	30 (22%)	26 (19%)	7 (5%)	1 (1%)
Positive	84 (63%)	65 (48%)	11 (8%)	5 (4%)	2 (2%)
Strongly positive	41 (30%)	4 (3%)	1 (1%)	1 (1%)	1 (1%)

Sensitivity of Methods

Specimens of faeces were collected from a normal subject who was on a meat-free diet. These were diluted with water to give a semi-solid specimen. Out-of-date transfusion blood (haemoglobin 13.3 g./100 ml.) was added to the sample to give various concentrations (blood expressed in millilitres and faeces in grammes). These dilutions of blood in faeces were tested for occult blood by several methods. In each case two separate sets of dilutions of blood in faeces were prepared and two tests carried out on each dilution. The tests used were:

(1) The orthotolidine tube test of Kohn and O'Kelly (1955), using five different dilutions of the stock solution.

(2) The "occultest" and "hematest" tablet tests. The occultest tablets are intended for the detection of blood in urine, and the hematest tablets for the detection of blood in faeces. In both cases it was found that the result was easier to read, and that more consistent results were obtained, if the tablet was placed at the edge of the faecal smear instead of in the centre. The method used was as follows. A thin smear of faeces was made on one of the test papers and an occultest or hematest tablet was placed half on and half off the smear; two drops of water were placed on the tablet so that the water flowed off on to the paper. The results were recorded as follows, depending on the time of appearance of a blue colour on the paper:

Strongly positive (+++) .. Colour within 15 seconds
Positive (++) .. " " " " " 15-60 "
Weakly positive (+) .. " " " " " 60-120 "
Negative (-) .. " " " " " No colour within 120 seconds

(3) Gregersen's slide test as described by Harrison (1957). "Benzidine for blood testing" supplied by British Drug Houses Ltd. was used. The results were recorded thus:

Positive (++) .. Colour developing within 30 seconds
Weakly positive (+) .. " " " " " 30-60 "
Negative (-) .. No colour developing within 60 seconds

The results obtained with these tests are shown in Table II.

Discussion

In this series of tests a large number of positive results were obtained with the 33% and 20% dilutions of the stock orthotolidine solution. It is obvious that neither of these

TABLE II.—Results with three Tests

Dilution of Blood in Faeces 1 in:	Haemoglobin (mg./ 100 g. Faeces)	Orthotolidine Tube Test					Tablet Tests		Grega- sen Slide Test
		Dilution of Stock Solution					Occul- test	Hema- test	
		33%	20%	10%	5%	2%			
5,000	3	—	—	—	—	—	—	—	—
2,500	5	+	—	—	—	—	—	—	—
1,000	13	+	—	—	—	—	—	—	—
500	27	++	+	—	—	—	—	—	—
250	53	++	+	—	—	—	—	—	+
100	130	+++	++	+	—	—	+	+	++
50	270	+++	+++	++	+	—	+	+	+++
25	530	+++	+++	+++	++	+	++	+	+++
10	1,300	+++	+++	+++	+++	++	+++	++	+++
5	2,700	+++	+++	+++	+++	+++	+++	+++	+++

dilutions can be used to test specimens from patients on a normal diet. At the other end of the scale, four positive results were obtained even with the 2% dilution of the stock solution. These were from two patients, in neither of whom was there any evidence of gastro-intestinal bleeding; one was suffering from pneumonia, and the other had fractured her femur about two months previously.

These results suggested that a certain number of false-positive results must be expected in patients on a normal diet unless very insensitive tests are used. It should be pointed out, however, that the two patients who showed positive occult blood results when on a meat-free diet had no clinical evidence of bleeding. One was convalescing after the amputation of a leg, and the other was a diabetic with peripheral vascular disease. It is quite possible that some of the patients in this series did have gastro-intestinal bleeding, and the results given in Table I should be taken as showing the *maximum* number of false-positive results which may be expected with the different dilutions of the stock orthotolidine solution when specimens from patients on normal diets are tested. It would be more satisfactory to test specimens of faeces from healthy subjects, but this was found to be impracticable.

On the basis of these results, the 5% dilution of the stock orthotolidine solution was chosen for testing specimens from patients who are on a normal diet, provided foods containing particularly large amounts of blood—for example, liver—are excluded. This dilution was chosen as representing a reasonable compromise between a high incidence of false positives and a low sensitivity. With this dilution, a strongly positive result is highly significant, but false-positive or weakly positive results may occur in up to 10% of cases. The reduced sensitivity of this test when compared with the sensitivity of the test using a 33% dilution of the stock solution, as recommended by Kohn and O'Kelly (1955), is, of course, inevitable. In cases of doubt a meat-free diet should be instituted and the more sensitive test used.

The use of the 5% dilution of the stock solution gives a rather less sensitive test than the Gregersen slide test, which is also intended for use without dietary restrictions but which involves the use of benzidine. However, Thornton and Illingworth (1955) have noted the occurrence of positive reactions in the Gregersen slide test with specimens from normal subjects on unrestricted diets.

If it is desired to carry out tests for faecal occult blood in the wards, the tablet tests have the advantages of convenience and simplicity. One criticism of these tests is that false-positive results may occur because the faeces are not boiled to destroy enzymes. A series of tests carried out in this laboratory suggests that this source of error is not of very great importance. A comparison was made of the results obtained with the occultest tablets and those obtained with the benzidine method described by Harrison (1957). Out of the 15 specimens which gave a negative result with the benzidine test, four gave weakly positive results with the occultest tablets, while none gave positive or strongly positive reactions.

The hematest tablet sometimes turns pale blue when the water is placed on it, even when the test is negative. This

may cause confusion, particularly with inexperienced staff, so we prefer the occultest tablet test for ward use, in spite of its greater sensitivity, which will presumably produce a higher incidence of false-positive results. If this test is used on specimens from patients on a normal diet, weakly positive results are of doubtful significance, but positive results are probably significant, and strongly positive results are highly significant.

Summary

A modification of the orthotolidine test for faecal occult blood is described. It may be used when the patient is on a normal diet, provided foods containing particularly large amounts of blood—for example, liver—are excluded. The test is much less sensitive than the standard orthotolidine test. Strongly positive results are significant, but false-positive and weakly positive results may be expected in up to about 10% of cases.

The occultest tablet test is suggested as a suitable alternative, particularly for ward use.

My thanks are due to Dr. H. Loewenthal, senior pathologist, for his advice and encouragement; to the medical and nursing staff of the Enfield Hospital Group who assisted in the collection of the specimens; and to Messrs. Ames (London) Ltd. for the supply of hematest tablets.

REFERENCES

- Harrison, G. A. (1957). *Chemical Methods in Clinical Medicine*, 4th ed. Churchill, London.
Kohn, J., and O'Kelly, T. (1955). *J. clin. Path.*, 8, 249.
Thornton, G. H. M., and Illingworth, D. G. (1955). *Gastroenterology*, 28, 593.

INDICATIONS FOR USE OF IPRONIAZID IN PSYCHIATRIC PRACTICE

BY

P. J. DALLY, M.B., M.R.C.P., D.P.M.

*Registrar, Department of Psychological Medicine,
St. Thomas's Hospital, London*

Iproniazid ("marsilid"; 1-isonicotinyl-2-isopropylhydrazine phosphate) is a derivative of isoniazid and was originally introduced because of its similar activity on tubercle bacilli. In the early clinical trials of iproniazid in pulmonary tuberculosis, using doses of 300 mg. daily, increased appetite associated with large gains in weight was noticed, accompanied in a proportion of patients by mood changes which sometimes developed into frank psychosis. These changes were reversible on stopping the drug (Kamman *et al.*, 1953; Crane, 1956).

The mechanism of this reaction is at present not fully understood, but it is thought to arise from the action of iproniazid as an inhibitor of monoamine oxidase. Iproniazid in animal experiments has been shown to produce an elevation of the 5-hydroxytryptamine (serotonin) level in the brain, and, if injected into rabbits over two to three days, to lead to a state of excitement.

These findings have aroused interest in the effect of iproniazid on depressive and allied affective states in man, and a number of papers have been published and a symposium was held in New York in November, 1957, on the use of iproniazid in psychiatric diseases. It seems to be generally agreed that iproniazid is of value in the treatment of depressive illness, but claims have been made that as many as 75% of patients do as well, if not better, with it than with electric convulsion therapy

(Kline, 1957). Electric convulsion therapy is widely used in the treatment of depression at the present time, and such claims, if confirmed, would obviously have far-reaching effects. A pilot trial has therefore been undertaken in this department to determine its value in depressive and allied psychiatric disorders.

Method

So far 131 cases have been treated with iproniazid. They comprise 99 depressions, 24 anxiety states, 6 obsessional states, and 2 cases of anorexia nervosa. Cases were at first unselected, but as the indications for the use of iproniazid became known the tendency was to choose those most likely to be helped. Both in-patients and out-patients were treated, but the majority were out-patients, these being seen once a week for the first month and then at progressively longer intervals.

Each patient was started on 150 mg. (three tablets) daily, in three divided doses, but was warned to reduce the drug to 100 mg. daily if side-effects developed, and to stop altogether if these continued. This daily dosage was continued for four weeks, and then reduced by 50 mg. at intervals of a month, until either the patient was off the drug completely or symptoms of depression again began to be experienced. If this happened the daily dosage of iproniazid was increased by 50 mg. weekly until the patient was symptom-free, and maintained for a further month before again being reduced.

The results are shown in the Table.

Results of Treatment

Type of Patient	No.	Re- covered	Im- proved	No Effect	Worse (Agitation and Tension Increased)
Depression with weight loss	67	15	22	21	9
" without weight loss	32	—	1	13	18
Anxiety state	24	—	—	—	24
Obsessional states	6	—	—	6	—
Anorexia nervosa	2	—	—	2	—

Discussion

Iproniazid was found to be of value in the treatment of certain patients showing symptoms of mild depression associated with loss in weight. There was a time-lag of between 7 and 10 days after starting the treatment before improvement was noticed, the patient then reporting that he had an increased appetite, a sense of "couldn't care less," and more energy. Some of these mild depressive illnesses had continued for several years, adversely affecting work capacity and family happiness. Amphetamine and its derivatives had had an immediate but short-lived uplifting effect. E.C.T. had caused temporary or partial improvement only. The natural course of the illness is to progress to full recovery, and this makes the interpretation of any treatment of depressive illness particularly difficult.

Many of the patients in this trial had been ill for over a year, and may therefore have been remitting naturally. None the less, iproniazid undoubtedly helped some to improve, and this was confirmed by the return of symptoms that sometimes occurred when the drug was reduced or stopped. The advantage of iproniazid over amphetamine or E.C.T. in this type of depression is that improvement appears to be maintained as long as the drug is given in adequate dosage. It thus seems possible that the course of some of these mild but obstinate depressive illnesses associated with loss of weight, which in these trials included both manic-depressive and involutional groups, may be shortened by treatment with iproniazid. Severe depressions, however, were not helped and depressive states associated with agitation were often made worse, and E.C.T. was usually