has by no means been emptied even by the spontaneous vomiting which has occurred during the stage of obstreperous behaviour.

Certainly the dangers of "silent regurgitation" and glottic spasm are every bit as great in certain types of emergency surgical cases—particularly cases of intestinal obstruction. It is for that very reason that many anaesthetists in these cases avoid a thiopentone induction at all costs and induce anaesthesia by a slower, safer inhalation technique, at least until the airway has been safeguarded by some means such as cuffed endotracheal intubation. If in such cases thiopentone induction is used, then it is established practice among many anaesthetists to lessen the risk of regurgitation by placing the patient in a slight reverse Trendelenburg tilt prior to injection and, in any case, by performing the induction with a sucker ready at hand for immediate use.

In conclusion, I feel that if thiopentone were widely used as a means of quieting unruly drunks, desirable though this end may be, without the facilities of an anaesthetic-room available, there would be introduced a far graver hazard to life than if intramuscular paraldehyde or apomorphine were substituted.—I am, etc.,

London, W.2.

PETER NOBES.

Phenylbutazone

SIR,—The toxic reactions after phenylbutazone ("butazolidin") therapy given for various arthritic conditions are well known, and include anorexia, nausea, vomiting, diarrhoea, dermatitis, palpitations, and, finally, agranulocytosis and sometimes complete aplasia of the bone marrow. The importance of these reactions has been stressed in many recent articles and in particular by Drs. J. R. Nassim and Thomas Pilkington¹ and by Dr. J. C. Leonard.²

I have taken 28 consecutive cases of rheumatoid arthritis, 20 of osteoarthritis, and two of gout which have been treated with this drug. The majority had 0.2 g. three times a day, reducing the dose in many instances to 0.2 g. once a day. Seven received intramuscular injections of 1 g. twice a week, reduced to 1 g. once a week after one month. The majority of the patients have been treated in this way for nine months and none for less than three months. All those with severe symptoms were admitted into hospital and the effect of the initial doses of phenylbutazone carefully assessed. After discharge from the wards they joined the others attending the out-patient department at intervals varying between one and three weeks. The possibility of toxic symptoms occurring was explained to each one, and they were told to stop the drug immediately if this happened and to report back to the clinic the following week.

All the toxic reactions which I have experienced have been mild and have either responded to a reduction in the dose or disappeared spontaneously without any interference. There was only one exception. Severe vomiting in this instance caused the drug to be stopped. The toxic symptoms encountered in this series of 50 people were: palpitations 1, conjunctivitis 1, nausea 9, vomiting 5, swelling of eyes 2, pruritus 1, diarrhoea 2, dermatitis 4, ulcerated gums (without agranulocytosis) 2.

Twenty-seven of the 50 patients, therefore, experienced some unpleasant symptoms, and this was higher than reported by Nassim and Pilkington (46 out of 109), is about the same as that reported by Cudkowicz and Jacobs,³ but higher than the average of 22% quoted by Leonard after a review of the literature. My results have differed from those of Nassim and Pilkington in one important respect: it was only necessary to stop the drug once, whereas they had to withdraw the drug in 46 out of 109 patients.

Out of the 28 patients suffering from rheumatoid arthritis, 21 had a very good result, three a moderate result, and one was made worse. Out of the 20 cases suffering from osteoarthritis, a very good result was obtained in 13, moderate in three, and complete failure in four. Both those suffering from gout were almost cured. Forty-two out of the 50 cases have derived considerable benefit from this drug: this figure is much higher than that quoted by Nassim and Pilkington (59 were greatly improved out of a total of 109).

The only other treatment for rheumatoid arthritis which has gained any popularity is that with injection of gold salts, and favourable results with this therapy have been few and troublesome complications due to dermatitis, liver and kidney damage, etc., have been frequent. Furthermore, there is no other drug known which will effectively alleviate the pain in osteoarthritis. The treatment of gout with

colchicum, cinchopen, or salicylates is uncertain. These facts must be taken into account before too much emphasis is placed on the toxic side-reactions of phenylbutazone.—I am, etc.,

Birmingham, 18.

R. O. GILLHESPY.

REFERENCES

- ¹ *British Medical Journal*, 1953, 1, 1310.
- ² *Ibid.*, 1953, 1, 1311.
- ³ *Lancet*, 1953, 1, 223.

Bornholm Disease

SIR,—The numerous excellent descriptions of the 1951 epidemic of Bornholm disease, or, as our Norwegian colleagues prefer, epidemic pleurodynia (*Journal*, June 20, p. 1345), and the letters from readers indicate a widespread interest among British physicians in the events of that late summer. Laboratory studies provide evidence of the responsibility of Group B Coxsackie viruses, as pointed out in your leading article (p. 1379). Type B3 apparently was prevalent. It also accounted for epidemics in the U.S.A. in 1950 and 1951, and was identified in Australia the same year. It may have been responsible for the epidemic on the Continent as well. We must wait for more information from the laboratories. Meanwhile it is worth noting that while B3 has now been repeatedly implicated, another type, B1, which has not appeared for some years, was the aetiological agent of the American cases in 1948 which led Dr. Curnen to suggest that Coxsackie viruses might be the cause of Bornholm disease.¹

The virologists can be relied upon to solve the question of how many types cause Bornholm disease by comparing their strains and identifying them in terms of a common terminology. Meantime a second point which may have more general interest seems to have attracted little attention in Great Britain. It is whether poliomyelitis and Bornholm disease ever occur simultaneously. The records indicate that 1951 was a relatively light poliomyelitis year in England and Wales, but I find no comment regarding the concurrence or competition of the two diseases. This has attracted attention in the past.

The first statement to come to my attention was that of Gsell,² who described a 1940 epidemic of Bornholm disease in Canton Appenzell and mentioned that an epidemic of poliomyelitis did not occur in 1940 and that "not a single case of infantile paralysis appeared during that summer and fall in Gais and Teufen." Linberg,³ I have since found, made a similar comment regarding a Norwegian outbreak in 1934. There are others. Lazarus,⁴ describing the epidemic in the State of Washington in 1950, the first association of type B3 with Bornholm disease, wrote, "... only one case of paralytic poliomyelitis occurred in Hoquiam during the 1950-1 season, when the pleurodynia epidemic took place. The incidence of poliomyelitis in Grays Harbor County during the same season was about 15 per 100,000 population. Preliminary data indicate that the same thing is happening this year, with cases of pleurodynia again appearing and paralytic poliomyelitis being at a low level." Gabinus, Gard, Johnsson, and Pöldre⁵ investigated an outbreak in south central Sweden in 1950. They mention, "No cases of paralytic poliomyelitis appeared in this region."

During 1951 the disease was widespread in southern Germany. Gärtner⁶ wrote, "In no place where Bornholm disease occurred was a single case of paralytic poliomyelitis recognized." And from Stuttgart, Windorfer and Born⁷ reported, "Interestingly in the same period only one definite case of paralytic poliomyelitis occurred in Stuttgart."

Dr. R. L. Huebner tells me there was not a single case of poliomyelitis in the Texas counties in which pleurodynia was prevalent during 1951, but there was little or no poliomyelitis the following summer as well. He established a type B3 virus as responsible for the outbreak. On the other hand, a type B3 virus appears to have caused widespread disease in Sydney, Australia, late in 1951, and Dr. Stanley⁸ was greatly impressed by the coincidental disappearance of poliomyelitis. It had been prevalent the previous year and returned in force in 1952. Most recently I find a description of an epidemic in Reykjavik, Iceland,⁹ and the comment, "In this period no cases of anterior acute poliomyelitis or aseptic meningitis were reported."

I mention these observations in the hope that readers of the *Journal* may have evidence that bears on the point.

Proof of an interference between the two in nature may be difficult to establish. Certainly all we have so far is merely suggestive. But the evident interest of your readers in the epidemic leads me to think they may have confirmation, and I would be grateful to them for their comment.—I am, etc.,

Albany, New York.

GILBERT DALLDORF.

REFERENCES

- ¹ *Bull. N.Y. Acad. Med.*, 1950, 26, 335.
- ² *Schweiz. med. Wschr.*, 1941, 71, 83.
- ³ *Acta paediat.*, 1937, 19, 1.
- ⁴ *Amer. J. publ. Hlth.*, 1952, 42, 20.
- ⁵ *Arch. ges. Virusforsch.*, 1952, 5, 1.
- ⁶ *Dtsch. med. Wschr.*, 1952, 77, 151.
- ⁷ *Ibid.*, 1952, 77, 1012.
- ⁸ *Aust. J. exp. Biol. med. Sci.*, 1953, 31, 31.
- ⁹ *J. Amer. med. Ass.*, 1953, 152, 814.

Prognosis in Diabetic Pregnancy

SIR,—In his interesting article (*Journal*, June 27, p. 1413) on the prognosis in diabetic pregnancy Dr. Wilfrid Oakley states that his results fail to confirm the existence of any direct relationship between age of onset of maternal diabetes and foetal loss. As mentioned in the article, this relationship was found by me,¹ but in a series of babies weighing 2,500 g. or more from patients without adiposity. Dr. Oakley uses for his study of this point his total King's College Hospital series (267 of 275 babies), and this may be the reason why the relationship was not found by him.

To get a clearer concept of the causes of the high foetal mortality and of what might be accomplished by a certain treatment, one has to know something of the "natural" prognosis in different groups of patients. The two most important factors seem to be parity of the mother and age of onset of diabetes,^{1,2} which cannot be influenced by any treatment. To use a total material for analysis without classifying may obscure everything but the gross mortality.

In his section on foetal hypoglycaemia, Dr. Oakley concludes that this problem clearly needs further study. I am inclined to draw the opposite conclusion. From what Dr. Oakley tells us it is evident that no relationship exists between foetal hypoglycaemia and neonatal death. In my rather extensive work³ on this topic I came to the same result. Since 1946 the newborn infants have not received glucose at the lying-in department B of Rigshospitalet, Copenhagen. It is a characteristic feature of the newborn young of mammals that they can stand a low blood sugar.—I am, etc.,

Copenhagen.

JØRGEN PEDERSEN.

REFERENCES

- ¹ *Trans. XIIIth Brit. Congr. Obstet. Gynaec.*, 1949, p. 186. Austral Press, London.
- ² *Ugeskr. Læg.*, 1951, 113, 1771.
- ³ *Diabetes and Pregnancy. Blood Sugar of Newborn Infants*, 1952, p. 230. Danish Science Press, Copenhagen.

Tropical Diseases in Britain

SIR,—May I be allowed to draw attention to the following points raised in your annotation, "Tropical Diseases in Britain" (*Journal*, May 16, p. 1096)? I entirely agree with Professor W. Melville Arnott's insistence in his letter (*Journal*, May 30, p. 1219) that time must not be lost in starting therapy in cases of malignant tertian malaria, and that in the seriously ill patient intravenous quinine is the most effective treatment.

The writer of the annotation claims that "true hookworm disease occurs only in debilitated populations suffering from heavy infestations superimposed on malnutrition and the avitaminoses; it is not seen in the well-nourished." In the last year I have seen many Europeans carrying ankylostoma ova who were not suffering from "true hookworm disease," but I have also attended two senior officers who were suffering from ankylostomiasis, the clinical features including the rash, pulmonary symptoms and signs, and a heavy intestinal load of parasites, resulting in both cases in severe debility. Both these officers were harassed, frustrated, and overworked, but neither was, or had been, malnourished nor presented any feature of the "avitaminoses."

The other statement in the annotation with which I cannot agree is that "emetine . . . should find no place in the diagnosis" of amoebic dysentery. In my experience the patient who suffers from dysentery which does not respond to phenylsulphathiazole in adequate dosage, and in whom *Entamoeba histolytica* or its cysts cannot be found on examination of several stool specimens, very frequently produces amoebae or cysts in the stools after one dose of 1 gr. (65 mg.) emetine. Further, the case of "pyrexia of unknown origin" in the Tropics may be due to amoebiasis, and if it has been impossible to find a cause for the illness, it is worth trying a course of emetine as a diagnostic measure.

In this part of the world a doctor or a laboratory technician is not expected to be an "experienced pathologist" in order to be able to detect the presence of *E. histolytica* or its cysts in a specimen of faeces, or to be able to differentiate such protozoa from the free forms or cysts of entamoebae which are at present considered to be of little pathological importance.—I am, etc.,

Kaduna, Northern Nigeria.

BRUCE NICOL.

Medical Auxiliaries

SIR,—The letter from Dr. A. P. Cawadias (*Journal*, July 11, p. 96) appeals on somewhat emotional grounds for "equal rights" under the N.H.S. for the three associations of physiotherapists he mentions. The Ministry's only interest in this matter should be in the standard to which medical ancillary bodies shall be required to attain before their members are accepted for employment in the N.H.S., and the absence of any reference to such standards in the Ministry's recent proposed regulations is a cause of considerable concern to many members of the medical profession.

Most of us have in the past relied upon members of the Chartered Society of Physiotherapy, because we know that they have undergone a full-time three-year training at one of the recognized hospital training schools. The published report of the evidence given before the Cope Committee gives little evidence of similar standards of training being required by the other associations mentioned by Dr. Cawadias. It was possibly for this reason that the Cope Committee felt it could not recommend the acceptance of the certificates of these associations for employment under the N.H.S. It may be, however, that further information is now available regarding the training syllabus of these bodies, and where it is carried out; if so this would be of considerable interest in view of Dr. Cawadias's plea.—I am, etc.,

London, W.1.

W. S. C. COPEMAN.

POINTS FROM LETTERS

Cord Traction

DR. JAMES T. BURROWES (Jamaica) writes: May I be allowed to record my support of Dr. F. C. R. Picton (*Journal*, April 18, p. 887) and the Brandt-Andrews method of placental extraction? This method gives such good results that the routine use of ergometrine intravenously before the commencement of the third stage would appear to be unnecessary.

Treatment of Herpes Simplex

DR. GORDON J. C. HERD (Elstree, Herts) writes: During the past year I have treated a number of cases of herpes simplex by the application of 10% chloramphenicol in propylene glycol. The eruption in most cases was confined to the lips, the first manifestation having presented some 12 to 24 hours previously. The area involved is washed with soap and water and then painted with the solution, which is left to "dry in." Treatment is repeated twice daily (morning and evening) for about three days. Following the initial application, local discomfort is alleviated, no further extension of the lesions occurring. Within 36 hours there is a marked diminution in the size of the vesicles. These generally break down by the third or fourth day, leaving small areas of granulations which subsequently heal. The only disadvantage I have observed is the unpleasant taste of the preparation should it find its way into the mouth. This effect, however, is evanescent.