PROSTATIC DYSPEPSIA

BY

B. E. CRAWFORD STANLEY, M.S., F.R.C.S.

Late Assistant to the Surgical Unit, St. Mary's Hospital, Paddington; Consulting Surgeon, St. Bartholomew's Hospital, Rochester

Although "silent prostatism" is a well-recognized clinical condition, it is surprising how few references to it occur in the current literature, and, with one or two possible exceptions, mention of the condition is conspicuous by its absence from the standard undergraduate and postgraduate textbooks of surgery and urology. Ewert and Summons (1951) have drawn attention to the not uncommon insidious onset of prostatic obstruction. They have emphasized that this condition may present itself with symptoms unrelated to the genito-urinary tract, and state that these symptoms are due to varying degrees of renal insufficiency.

Silent prostatism is more common than is generally realized and may present as a gastro-intestinal disturbance, with a history suggestive of carcinoma of the stomach or early hepatic cirrhosis in the absence of urinary symptoms and before there is biochemical evidence of renal insufficiency or azotaemia. Eighteen cases of this type were encountered during the last five years in a series of just over 200 patients who had undergone prostatectomy.

The presenting symptoms of this syndrome, which may well be called "prostatic dyspepsia," are upper abdominal discomfort or pain, nausea, and anorexia. Urinary symptoms are absent, but the residual urine is invariably much increased and the cystoscopic appearances confirm the diagnosis. The following case histories are typical.

Three Case Reports

Case 1.—A man aged 73 was admitted to a medical ward for investigation. He had a four-months history of dyspepsia, anorexia, and vomiting with loss of weight. Barium meal examination and fractional test meal and liver-function tests showed nothing abnormal. Blood urea was 27 mg. per 100 ml. An intravenous pyelogram showed normal function of both kidneys, but a large filling defect at the base of the bladder and considerable vesical retention in the post-micturition film. Cystoscopy was attempted, but the instrument was not long enough to get through the prostate, which could be felt above the symphysis pubis when the bladder was empty. A one-stage prostatectomy was carried out and a gland weighing 340 g. was removed. Three months after operation the patient declared himself to be in perfect health with complete restoration of his digestive powers.

Case 2.—This patient, aged 65, was referred by his general practitioner with a history of four months' anorexia, pain, and nausea. He had lost 2 stone (12.7 kg.) in weight. He would admit to no urinary symptoms. Barium meal, test meal, and liver-function tests revealed no abnormality. Blood urea was 34 mg. per 100 ml.; haemoglobin, 76%. The history was so typical of gastric carcinoma that laparotomy was advised. This was performed, but no lesion of the stomach or intestines was detected. When the hand was palpating the pelvis the bladder was felt to contain several ounces of urine. Subsequent investigations showed a residual urine of 8 oz. (230 ml.), a normal intravenous pyelogram, and a moderately trabeculated bladder. The man did not rise at night to pass his water and had complete control with no delay or urgency. Removal of a small prostate (24 g.) cured his digestive symptoms, and his weight was back to normal three months after operation.

Case 3.—A man aged 68 attended his doctor complaining of feeling out of sorts with loss of appetite and constipation. He had no urinary symptoms, sometimes rose once at night to micturate, but passed urine with a good stream. On examination he had a bladder palpable 3 in. (7.5 cm.) above the symphysis pubis. The prostate was moderately enlarged but benign. The blood urea was 36 mg. per 100 ml., and the urine was uninfected. An intravenous pyelogram showed normal concentration in the kidneys with no hydronephrosis. The bladder shadow was enlarged. Cystoscopy showed moderate intravesical projection of the prostate and trabeculation of the bladder, with 30 oz. (850 ml.) residual urine. This examination precipitated an acute retention of urine, so a closed transvesical prostatectomy was performed 24 hours later. Convalescence was delayed by a femoral thrombosis, but when seen three months after operation, although he still had a little swelling of the leg, he said he was feeling fitter than for some time.

Remainder of Series

Fifteen other cases of the condition have been seen, a summary of whose clinical histories are tabulated below.

Details of Cases

Age	Predominant Symptom	Initial Provisional Diagnosis	Blood Urea mg./100 ml.	Urine	Residual Urine	
					oz.	ml.
74	Anorexia	Carcinoma of stomach	34	Sterile	8	230
72	Epigastric discomfort	" "	39	"	6	170
58	" "	" "	31	"	14	400
60	Nausea	Gastric ulcer	44	Infected	6	170
64	Epigastric discomfort	Duodenal ulcer	32	Sterile	18	510
70	Anorexia	Cirrhosis	37	"	6	170
78	Nausea	" "	40	"	14	400
61	Anorexia	Carcinoma of stomach	28	"	20	570
74	Nausea	Gastric ulcer	31	Infected	8	230
72	Epigastric discomfort	" "	42	Sterile	23	655
82	" "	Duodenal ulcer	36	"	8	230
76	Nausea	Carcinoma of stomach	45	"	6	170
72	Epigastric discomfort	Gastric ulcer	29	"	15	425
70	Anorexia	Carcinoma of stomach	31	"	10	285
54	"	Cirrhosis	38	"	22	625

Comment

It is a well-recognized fact that patients who have had a prostatectomy frequently volunteer the information after the operation that they feel fitter than they have felt for some time. The observation is not limited to those in whom there has been some degree of uraemia.

Several of the patients in this series were submitted to detailed investigations, including urea concentration and clearance tests, blood alkali reserve, etc., and in none has there been evidence of renal impairment.

The cause of the syndrome of dyspepsia, nausea, and anorexia, associated with prostatic obstruction which has not gone on to the stage of renal impairment, is not clear, but is in my submission a definite entity which is not generally recognized until uraemic symptoms supervene. It would seem that the importance of these cases, which in my experience account for a little over 5% of all patients requiring prostatectomy, lies in the fact that, unless the clinician concerned, usually not a surgeon or urologist, is alive to the possibilities of prostatic obstruction being the underlying cause, the patient may be sent away after a negative gastro-enterological examination or submitted to an unnecessary laparotomy, when the catheter to measure the residual urine or the cystoscope will make the diagnosis clear.

Summary

A syndrome of epigastric pain, nausea, and anorexia is a not uncommon initial complaint in patients with prostatic obstruction without urinary symptoms.

A series of eighteen patients are described with this syndrome in whom there was no real evidence of renal impairment.

The importance of the condition lies in the fact that after a negative gastro-enterological investigation it can be, and often is, overlooked until uraemic symptoms supervene.

REFERENCE

Ewert, E. E., and Summons, H. J. (1951). *Surg. Clin. N. Amer.*, 31, 659.

HYPERPARATHYROIDISM WITH NORMAL SERUM CALCIUM

BY

H. GORDON MATHER, M.B., M.R.C.P.

Medical Registrar, King's College Hospital, London

Osteitis fibrosa cystica was first described by von Recklinghausen in 1891, but the first clear demonstration that a parathyroid adenoma was the cause came in 1925, when Mandl cured a man crippled with the disease by the removal of such a tumour. Since then our knowledge of the disease, its clinical, radiographic, and biochemical features, has steadily increased, though as recently as 1950 Rienhoff estimated that there were only 597 proved cases in the world's literature or known to him personally. Thus the disease is rare enough to warrant a record of every case, especially when there are any unusual findings, and since treatment is so effective.

Case Report

A spinster tea-packer aged 33 first consulted her doctor in February, 1951, because of "rheumatism" in the legs and back for five months. Before this her health had been good; she had lived an active life and had played tennis during the previous summer. The aching pain had started in her right thigh rather suddenly in September, 1950, and had spread in several weeks to involve the lower back, thighs, and knees. The pain was made worse by movement, but did not prevent sleep. She noticed increasing weakness in the legs, so that walking, especially starting off, became difficult: a waddling gait developed. These disabilities caused her to stop her work at Christmas, 1950. The symptoms became increasingly severe; some aching and weakness developed in the right shoulder, and a painful lump appeared below the manubrium sterni. She had noticed some frequency of micturition, which disturbed her twice each night.

She was referred to the medical out-patient department, where the most striking abnormality was her gait as she entered the room. It resembled the waddling of a duck, and was thought to be due to muscular weakness, especially of the hip abductors. She was of stocky build, and had a marked thoracic kyphosis, which, though present since childhood, she thought had become more pronounced. The bones were not tender, and passive movement of the limbs was full and pain-free. There was definite muscular weakness of the abductors, flexors, and extensors of the hip-joints, but no wasting and no reflex or sensory changes. Examination of the neck, lungs, heart, and abdomen was normal. The blood pressure was 130/90 and weight 8 st. 9 lb. (54.9 kg.). There was a tender bony lump in the sternum between the third

costal cartilages. X-ray examination of the pelvis revealed a generalized osteoporosis, so she was admitted for investigation with a tentative diagnosis of hyperparathyroidism.

Investigations.—In order to confirm the clinical diagnosis and to exclude in particular multiple myelomatosis and polyostotic fibrous dysplasia (Albright's disease) extensive investigations were made. Skeletal x-ray films revealed generalized osteoporosis, with a "moth-eaten" appearance in the skull (Fig. 1), and early fibrocystic areas, especially in the pelvis and right upper humerus. There was a fracture of the body of the sternum, and a loss of the lamina dura of the teeth. No calcification was seen in the renal tract. Four pre-operative serum calcium estimations (10.8, 10.32, 9.8, and 9.7 mg. per 100 ml.) during three months were within the normal range: the method used was that described by Trevan and Bainbridge (1926). Two out of the three plasma phosphorus estimations (1.7, 2.2, and 1.8 mg. per 100 ml.) were, however, below normal limits. The serum alkaline phosphatase was 92.4, 94.4, and 98.0 King-Armstrong units per 100 ml. at monthly intervals. The blood urea (34 mg. per 100 ml.) and urea clearance were normal, as were the serum proteins (total, 6.7 g.; albumin, 4.16 g. per 100 ml.). An eight-day calcium balance showed that on an intake of 97 mg. daily her average urinary excretion was 322 mg. and faecal excretion 162 mg., a daily negative balance of 387 mg. The combination of low plasma phosphorus and a negative calcium balance, even in the presence of normal serum calcium, made the diagnosis almost certain.

A sternal bone biopsy showed osteoclastic destruction of bone trabeculae associated with proliferation of fibrous tissue. There was very little evidence of new bone formation and there was no mosaic pattern in the islets. The appearances were those of osteitis fibrosa cystica.

A blood count showed: red cells, 4,300,000 per c.mm.; Hb, 11.9 g. (80%); colour index, 0.9; E.S.R. 4 mm. (Wintrobe corrected); white-cell count, normal. There were no casts, cells, or albumin in the urine.

Operation.—On June 11, 1951, Mr. E. G. Muir explored the neck through the usual collar incision. A normal parathyroid gland was identified on the left side, but a small parathyroid adenoma was found attached to the right lower pole of the thyroid. It was removed by dividing the short pedicle. The adenoma was deep red in colour with a smooth surface; it measured 1.8 by 1 by 0.8 cm. and weighed 0.9 g. Microscopical examination showed parathyroid tissue composed almost entirely of "wasserhelle" cells.

Subsequent Course.—The patient was given 2 pints (1,140 ml.) of milk, 60 gr. (4 g.) of calcium lactate, and 10,000 units of calciferol daily from three days before operation until six months afterwards. The bone pain vanished immediately after operation, and muscle strength returned within a few weeks. Five days after operation latent tetany was detected by a positive Chvostek's sign; at that time the serum calcium was 7.9 mg. per 100 ml. The dose of calcium lactate was increased to 240 gr.

FIG. 1

FIG. 2

FIG. 1.—Skull. Moth-eaten osteoporosis. FIG. 2.—Skull. Six months after operation.