carried out. But local conditions may complicate in a varying degree the solution of these problems. In most countries of Latin America, and possibly in some other regions of the world, an excess of students and economical factors complicate the picture. An erroneous concept of democracy has frequently opened the medical schools to all those who merely wish to pursue a medical education. The result has been the admittance of a number of students entirely incompatible with a high efficiency in the training, and this situation has brought about all the evils of mass education, in which the personal and highly useful contact between the teacher and the student is practically suppressed. The limitation of funds available for teaching and research makes difficult the introduction of the necessary reforms and the proper use of men adequately trained. In addition, geographical and social conditions are still responsible for large areas of inadequate medical and sanitary attention, which results in an uneven distribution of the graduates, who seek the large populated centres in which they find a better scientific environment for their professional activities.

Fortunately, facing these handicaps there is, in the medical school authorities and members, and in the medical profession in general, a sincere desire and a determined intention to raise the educational standards to the level found in those countries which have marked out the pathways in this field. These sentiments explain the presence in this meeting of men who have come from our distant continent. Allow me to conclude by expressing my confidence in the ultimate result of these deliberations, and my best wishes, participated in. I am sure, by all of you, for the success of this conference, which represents a united effort to raise the educational standards of our profession.

PREVENTIVE AND SOCIAL MEDICINE*

BY

RENE SAND, M.D.

*Emeritus Professor of Social Medicine, University of
Brussels*

In the hospital as well as in private practice the physician deals essentially with sick people, so that the codes of ethics drawn up by the medical profession in the various countries define the doctor's duties towards his patient, not in a more general sense towards the person who consults him in health or disease. So had Hippocrates in the *Oath*, although elsewhere in his work he maintains that the first duty of the physician is to protect and cultivate health.

When, about a century ago, sanitation was revived and put mainly in the hands of the engineer, and when medical health officers were appointed, the doctor still served the cause of health by giving advice as family physician. Unfortunately, the advent of specialization has nowadays reduced the part played by the general practitioner.

Preventive medicine in the strict sense of the term—that is, preventive medical procedures—appeared with Jenner and his first immunization, followed a century later by the preventive examinations of pregnant women, infants, and school-children. Occupational medicine was launched by Ramazzini in 1700, medico-social surveys and statistics by Villermé in 1828.

In the century and a quarter which has elapsed since these pioneer days, preventive and social medicine—the terms are often used as synonyms for reasons of con-

venience—have proved their case, and if weighed against curative medicine would no doubt be found more effective and less costly. However, they occupy only a small place in medical teaching; and they come late in the curriculum, at a time when the student's mind is already monopolized by the clinical and laboratory aspects of disease. Besides, having had no instruction in the social sciences, he feels lost in a strange country.

Signs of Improvement

Here, as in so many other fields, institutions have lagged behind needs. Hospital and medical practice are still turned mainly towards cure. The Faculty is still made up of a large group of clinicians, a large group of laboratory workers, and a solitary professor of public health. Yet signs of change are appearing. The importance given to preventive medicine by the Report of the B.M.A. (*The Training of a Doctor*), and by the Goodenough Report, as well as by similar documents elsewhere; the creation of departments, institutes, and chairs of preventive and social medicine in various countries; the part assigned to these disciplines by W.H.O. and other international organizations—all point in the same direction.

If the social elements in many medical schools are still comparatively neglected it is because the training takes place in the hospital, from which these elements are absent, and because few members of the Faculty have studied social sciences and acquired experience in social work. However, a growing number of hospitals have almoners, and use them to demonstrate the circumstances and needs of the patients. Besides, the idea is spreading that a much greater part of the training should be given in the out-patient department and in the home itself, either by associating the students with the daily work of general practitioners, as Professor Crew does in Edinburgh, or by sending the students to observe domiciliary cases.

Need for Change

The lack of training in the social sciences calls for a radical change: in his first university year the student is taught mathematics, physics, chemistry, and biology, which are required for the understanding of health and disease; he should study also psychology and sociology, not only because he will need this initiation to play his full part as a doctor but because everyone, and more especially a university graduate, should understand himself and the world in which he lives. Biology teaches us to understand our body; psychology to understand our mind and our behaviour; sociology to understand the structure and life of the groups, small and large, of which we are members. So long as psychology and sociology remained mainly speculative their practical use was limited. More realistic to-day, they have become indispensable to the statesman, the employer, and the parent, to the judge, the Army officer, and the doctor.

On the background of biology, psychology, and sociology, acquired in the first year, the principles of preventive and social medicine should be taught at the same time as the basic sciences, anatomy, and physiology. The aim of medicine should be defined as the maintenance and culture of physical, mental, and social health, with the repair of damaged health as the second consideration only. The nation-wide and permanent achievements of preventive medicine and public health should be put in parallel with the remarkable but individual and temporary successes of curative medicine.

Later, when the student enters the hospital, the part played by social, economic, and occupational factors should be explained by the internist, the surgeon, the obstetrician, the paediatrician, the psychiatrist, with the help of the almoner, and preferably also of the professor of preventive or social medicine; and this teaching should take place in the clinical lessons, in the daily rounds of the wards, in the out-patient department, in the dispensaries and clinics, and in the home itself.

*This paper was received for publication just before Professor Sand underwent an operation which prevented him from coming to London as one of the Vice-Presidents of the First World Conference on Medical Education. The news of his death was received as this issue of the *Journal* went to press.

Then preventive and social medicine will pervade medical training and practice as did bacteriology, serology, and immunology when I was a medical student more than fifty years ago. However, bacteriology was not as foreign to men trained in laboratory methods and the use of the microscope as are the social sciences and social work to those who never approached these fields.

There are other difficulties—the crowded curriculum, the natural resistance of men and institutions to any change, and the trouble and expense connected with a fundamental move.

But we do not want to make the curriculum still heavier : what we ask for is a broader point of view in the existing lectures rather than additional courses. Besides, there remains from former schemes of training much that is out of date and could be abridged or omitted. Specialization also should be curbed in the student's programme. It is impossible to teach everything in a limited number of years. We can give the background, the method, the example, the essentials of medical science and practice : above all, we must try to cultivate the spirit of scientific observation and the spirit of service ; the rest will have to be acquired by and by.

The lack of medical personnel trained in the social sciences will gradually be overcome. Facilities have been given in this direction by the Rockefeller and other American Foundations, the Nuffield Provincial Hospitals Trust, and W.H.O. The chairs recently created in various countries—whether called chairs of social medicine, or preventive medicine, or human ecology as in Cambridge—will breed a new staff and spread a new spirit.

Besides these general points, there are others of special interest, among them the place of occupational health in medicine. This deals with a specialized field ; it has developed its own methods in occupational physiology, pathology, and psychology ; and has grown to a full-sized science with its own associations, periodicals, and congresses.

Other fields, such as epidemiology, demography and vital statistics, genetics, nutrition, and housing, are allotted by some to public health, by others to preventive and social medicine.

To what extent can the student health service be used in medical education ? Experience shows great differences, according to the amount of discretion which national psychology demands in personal matters. Fruitful developments may nevertheless be expected in this direction.

Pioneers of Social Medicine

The term "social medicine" was coined in 1848 by Dr. Jules Guérin, of Paris. The teaching of social medicine began in 1902 with a few extracurricular lessons given at the Berlin Medical School by Professor Martin Kirchner. The first fully fledged professors of social medicine were Edoardo Bonardi in Milan (1908), Johan Marcus Baart de la Faille in Utrecht (1918), and Alfred Grotjahn in Berlin (1920), who had taught the subject, as lecturer, since 1912. Giuseppe Tropeano, in Naples, began his optional lessons in 1910. At Nancy, when taking his chair (1920), Jacques Parisot had it styled "chair of hygiene and social medicine." In the New World, the first department of preventive medicine was created at Harvard University in 1909.

Contributions to preventive and social medicine have come from various fields : the social sciences and social work, forensic medicine, public health, and clinical medicine. These various approaches have led some professors of forensic medicine, in France especially, to teach also preventive and social medicine ; professors of public health have done the same in various countries ; clinicians are increasingly following this example. The merits of these various schemes will be weighed against those of an autonomous chair of preventive and social medicine. That these disciplines should be included in the medical curriculum no one, I think, will deny.

PHYSIQUE AND CHOICE OF FACULTY

BY

R. W. PARNELL, D.M., M.R.C.P.

Nuffield Research Physician in the Constitutional Aspects of Psychiatric Medicine, the Warneford Hospital, Oxford

In the report of the 1951 Conference of Home Universities the Vice-Chancellor of Leeds, Mr. C. R. Morris, is quoted as saying : "There is no question that, in the main medical schools, the bottom candidates are very far indeed below the bottom candidates in the special schools of, say, physics and chemistry. They are, in fact, a different order of being academically. . . . I think that these people who are very low academically may be extremely good candidates for the medical profession. I hope you will not take that as a joke, because it may very well be true. Some very good doctors with long experience think that some of these people will fill, say, 40 or 50% of the medical profession with better advantage to the nation than others with very high academic qualifications."

In the same discussion on selection of university students, Dr. C. F. Harris, Dean of St. Bartholomew's Hospital Medical College, remarked : "With one or two exceptions, if one follows the careers of winners of our open scholarships it is apparent that the method of always allowing the open scholarship to be won by examination alone fails. The scholarship winner may have matured as an examination fiend much too early, and later this ability rapidly fails, or, apart from his examination brilliance, he may have qualities most undesirable in a doctor."

These engaging comments, coming from a leader in the world of education and the dean of a medical school, command attention, and any objective evidence which may be brought forward to support or refute these contentions is therefore of interest.

Temperament and Physique

In 1951 I reported on physical characteristics and differences between men who shine at short- and long-distance running. Association of physical characteristics and physical prowess is hardly surprising. More recently, however, in the course of a long-term study of physique and university performance, it has become apparent that qualities of temperament may be closely enough associated with physique to influence academic performance (Parnell, 1952). These findings it is hoped to report in full when the men who were measured have all completed their final examinations. Their choice of faculty, however, is known, and the association between this and certain physical characteristics can be described now.

Choice of faculty is partly a matter of temperament, and since temperament has been shown by Sheldon and Stevens (1942) to correlate fairly closely with physique it was thought worth while to plot the somatotype distribution of university students according to the faculty of their choice. It so happened that Sheldonian somatotype ratings were available for some 400 Oxford men undergraduates, and that various anthropometric measurements enabling a reasonably accurate estimate of somatotype to be made were taken during routine health examination of students entering Birmingham University in October, 1952. Because of the great variety of reasons governing choice of faculty it is perhaps surprising that any correlation with physical features should be found at all ; but the results are interesting. They are given in the series of figures illustrating somatotype distribution in the