CLINICAL PATHOLOGY IN GENERAL PRACTICE

VENEPUNCTURE

BY

I. A. B. CATHIE, M.D., M.R.C.P.

Clinical Pathologist, The Hospital for Sick Children, Great Ormond Street, London

There are nine and sixty ways of constructing tribal lays,

And every-single-one-of-them-is-right !—KIPLING.

As venepuncture is so commonly performed, it is probably the subject of more likes and dislikes and individual variations in technique than any other minor medical procedure. No author can do more than state one series of preferences, many of which may be anathema to those practising a different approach. What follows, therefore, is clearly open to limitless minor variations and cannot please everyone.

Venepuncture in the adult, where the vein to be entered is usually large and visible, is quite different from that in the infant, where the chosen vein is not seen but must be entered by the observation of anatomical landmarks. The two will be considered separately; in both it will be assumed that there is available a sterile syringe with airtight plunger, together with an unblunted needle not smaller than size B.W.G. Smaller needles are not materially less painful for the patient, and withdrawing of blood through them is more difficult and more liable to yield a haemolysed specimen.

As bacteria may interfere with so many tests the aim of every venepuncture should be to obtain a sterile sample of blood to be transferred into a sterile container.

After the hands have been thoroughly scrubbed and dried the instruments are examined. Whatever the size of the patient an essential preliminary to any venepuncture is to assemble the syringe and ensure that the plunger moves freely and the needle has not become blocked. The needle should be placed on the syringe nozzle with sterile forceps—if it was not in position before sterilization. No part of the fingers should touch the needle or the surface of the plunger, either at this stage or when the vein is being entered. In choosing a syringe one larger than apparently necessary is always to be preferred to one the size of the blood sample needed, as so often air bubbles replace much of the space which should contain blood. A 10 ml. syringe is suitable for most purposes. It should be ensured that the cover of the container into which the blood is to be put is free so that it can be removed with one hand while the other expels the blood into the bottle or tube (this is particularly important with screw-capped bottles).

Adults

Both for ease and convenience the antecubital fossa veins are nearly always used (see Fig. 1), with the arm lying on a table of convenient height. Either the central or lateral veins are used; in spite of the way the lateral accessory cephalic vein stands out the cephalic or median cubital veins are bigger and easier to penetrate. The lateral vein is more mobile and therefore more likely to be pushed aside, and incidentally it is more painful to have punctured.

1. Inspect and palpate the antecubital fossa and select the vein to be used. If necessary compress the upper arm with the free hand to make the veins stand out; if the patient "makes a fist" a few times this will help.

2. Sterilize the skin of the antecubital fossa with whatever antiseptic is preferred. Weak iodine solution, cetrinide, spirit, or ether are commonly used; a coloured antiseptic is useful, as the painted area can be seen.

3. Obstruct the venous return by compressing the upper arm. This can be done with a piece of rubber tubing, tied as in Fig. 1 for quick release, with a sphygmomanometer cuff inflated to a pressure of 90–100 mm. Hg, or by getting the patient to grip his upper arm with his free hand so that the veins are compressed beneath the belly of the biceps against the humerus. This can be demonstrated when selecting the vein, and is often advantageous in the nervous patient as it takes his mind off the coming prick and makes him feel he is being useful.

4. Set the arm supine on the table and resterilize the skin.

5. Get the patient to make a few more fists, then tell him to keep quite still and enter the selected vein. The bevel of the needle should face the skin, to minimize the risk of transfixing the vein. The vein may be approached directly through the skin over it or by penetrating the skin at the side of it and then attempting to enter the vein from its side. The first procedure is much the easier, but large and simple-looking veins are sometimes pushed aside by the needle.

6. Make slight suction on the syringe with the free hand when it is thought that the needle is in the vein. If no blood enters the syringe push it gently forward in the line of the vein, keeping up slight suction. If about two-thirds of the needle have gone into the arm and no blood has been obtained it is legitimate with the free hand to palpate the skin to feel where the needle point is. Then, making gentle suction again, slowly withdraw the needle to but not out of the skin, stopping immediately, of course, should blood be obtained. If still unsuccessful the process should be repeated, more than once if necessary, until the blood is obtained.

If swelling in the antecubital fossa, blue discoloration of the skin, or oozing of blood round the needle should appear, indicating haematoma formation, the pressure on the upper arm is released immediately, a swab with antiseptic on it is placed gently over the needle, the needle is withdrawn sharply—at the same time pressing the swab down firmly—and the patient is told to bend his arm double tightly. Haematomas are obviously caused by damage to the vein, usually by exploring too deeply and through its deep wall.

7. When blood is entering the syringe freely, slacken off the pressure on the upper arm. In fat arms, where the veins can be palpated but not seen, and sometimes when they cannot even be palpated, it may be necessary to maintain the obstructive pressure until the syringe is full. If blood starts to enter the syringe and then stops it can usually be got flowing again by moving the point of the needle slightly; often merely elevating the point is effective. Too great suction should not be made on the plunger, as this tends to drag air into the barrel, giving a smaller yield of blood than is required.

8. When the requisite blood is in the syringe, make sure that all pressure is released from the upper arm, hold a swab over the site of puncture, withdraw the needle sharply, at the same time pressing the swab down firmly. Tell the patient to bend his arm double and keep still. Hold the syringe with the needle slightly upwards; if it is held pointing downwards blood may drip out, while if held too high the plunger has been known to fall out of a full syringe.

9. Deal with the blood sample, distributing it into the appropriate containers.

10. Wash out the syringe and needle with water, and dismantle the syringe.

11. By this time any bleeding from the patient's arm should have stopped. Straighten his arm gently, at the same time keep-

FIG. 1.—Veins of the arm, with quick-release tourniquet in position.

ing light pressure on the swab for a few further seconds. Remove the swab and bathe the antecubital fossa with more antiseptic. Any tendency to ooze can be controlled by further pressure for a minute or two, or the arm may be held vertically for half a minute while moderate pressure is maintained on the puncture.

12. Nothing further should be necessary, although some like to put a small piece of adhesive plaster over the puncture.

The above is essentially a "no-touch" technique, and, provided the needle and the patient's sterilized skin are not touched, should yield a sterile blood sample. Nothing can be deprecated more than a guiding finger lying along the needle, a practice all too often seen. The technique is perfectly simple to carry out single-handed, although more efficacious hand-washing drill can be practised if there is someone to assist by passing swabs and holding blood containers.

Adult venepuncture may be unsuccessful in the most expert hands for many reasons. The two commonest are either simply failure to enter the vein or damage to the vein with haematoma formation. In the former case the arm should be re-sterilized and another vein attempted. When a haematoma has been caused no further obstructive pressure should be made on the arm, which would lead to further blood extravasation, and after ensuring by firm pressure that any bleeding and oozing has stopped an attempt should be made on the other arm.

Children

By the age of 5 most children have veins well enough developed for the above technique to be used, and often as young as 3 a successful venepuncture can be performed. As distinct from an adult, who usually wants to get the business over and is therefore co-operative, children will generally struggle, and assistance is imperative if an effective puncture is to be carried out. Even when the arm is skilfully held above and below the elbow it is amazing how much movement the average child can still achieve. The operator must remember to be most flexible and prepared to allow his hand, the syringe, and particularly the point of the needle to move with the wriggling that the child's arm may produce. Movement is minimized if the arm is abducted and externally rotated so that the back of the wrist lies flat on the couch. The hand of the assistant round the biceps acts as a tourniquet, but often with smaller arms the assistant's second hand at the child's wrist seems enormously in the way. A rehearsal without the syringe will usually enable the operator and assistant to dispose themselves most conveniently.

Infants

From birth to the age of about 3 years visible or palpable veins cannot be expected, and some other approach must be made. The most popular, particularly where quantities of blood up to 10 ml. are required, is the internal jugular vein, while the external jugular, the femoral, or scalp veins are sometimes used. The technical preliminaries regarding apparatus and sterilization are exactly the same as for adult venepuncture.

Internal Jugular Puncture.—The essence of this simple and safe procedure is in the position of the baby. It is first wrapped tightly in a blanket which comes up to the shoulders and retains the baby's arms at its sides. The baby is then held on its back with its head just overhanging the edge of a couch or cot. The head is rotated away from the side to be punctured, so that the sternomastoid muscle is put on the stretch. After sterilizing the skin, the operator inserts the needle at the posterior border of the sternomastoid, halfway between the mastoid process and the sternal notch (Fig. 2). Then, maintaining gentle suction on the plunger, the needle is pushed slowly beneath the muscle in a direction at right angles to the long axis of the baby. The underside of the sternomastoid is easily felt with the tip of the needle, and the aim is to keep as closely under the muscle as possible. In the majority of cases the blood will enter the syringe almost immediately and the required amount can be withdrawn. The position of the internal

jugular vein varies from child to child; it may be necessary to push the needle across the neck as far as three-quarters of an inch (2 cm.) or more before blood is obtained. If no blood flows the needle is slowly withdrawn, maintaining suction all the time, until the tip of the needle can be seen under the skin at the point of entry. The traverse beneath the sternomastoid can then be repeated until the vein is entered. After blood is withdrawn moderate pressure with a sterile swab is maintained over the site of puncture for about half a minute. If the attempt at venepuncture is unsuccessful on one side of the neck the same procedure should be carried out on the other. The secret of the technique seems to be the full extension and rotation of the child's head, and blood is got much more readily from a crying child than from one which none of this handling causes to cry. A variation of this method of internal jugular puncture is to enter the skin at the mid-point of the posterior border of the sternomastoid, the needle then being directed towards the sternal notch. This oblique approach has nothing to commend it over the direct approach, while anatomically it is potentially more dangerous.

FIG. 2.—Position for internal jugular puncture, showing the course of the external jugular vein.

External Jugular Puncture.—The external jugular vein runs obviously and temptingly across the sternomastoid (Fig. 2), and puncture of it is often unsuccessfully attempted. The reason is that the vein is relatively unfixed, and is so mobile that it is more likely to be pushed aside than penetrated by the needle. On occasions the vein can be successfully punctured, and it makes no difference whether the direction of the needle is towards the head or the heart.

Femoral Vein.—This lies on the medial side of the femoral artery, and can be entered just distal to the inguinal ligament. The baby is held flat on its back with a sandbag under its buttocks, and after the position of the vein has been ascertained by palpation of the artery the needle is either directed towards the heart or else thrust vertically into and through the vein, blood being sucked into the syringe after the point of the needle has been slightly withdrawn. Large amounts of blood may be got from the femoral vein, which has been used with increasing frequency in recent years (Fig. 3).

FIG. 3.—Femoral vein puncture.

Scalp Veins.—Small amounts of blood may be obtained from these. The wrapped baby is held with its head on one side, while digital pressure is made in front of the pinna of the ear. With the scalp over the temple and above the ear shaved a choice of three or four small veins will be seen lying between the top of the head and the front of the ear. They are brought into greater prominence if percussed lightly. The importance of the scalp veins is not so much as a source of blood as a channel through which small intravenous injections may be given, particularly in the newborn.

It will be appreciated that when it is desirable that all of a substance should be given intravenously with none in the surrounding tissues none of the other methods of venepuncture in infants is satisfactory, as the exact position of the needle point cannot be seen. The infant's scalp veins can be seen, withdrawal of a little blood will ensure that the needle is truly in the vein, and as the veins are so superficial any of the injection fluid not entering the vein can be seen and the injection stopped at once.

Fontanelle puncture was formerly much used in infants, a special short-needle with a short bevel and a guard being introduced through the posterior angle of the anterior fontanelle into the superior sagittal sinus. This method is not without risk, and now that other techniques have been perfected it should no longer be used.

Cord Blood.—This should not be forgotten as a simple and ready source of blood where the necessity for blood examination is likely to arise immediately after birth. It is easy to collect, and equally easy to discard should it not be wanted.

Heel Prick.—Lastly, in spite of the available methods of venepuncture in infants there are occasional cases where all efforts to get venous blood are unsuccessful. In these, and in cases where not more than 1 ml. of blood is required and sterility is not important, the possibilities of the heel prick

should not be forgotten. While it is capillary and not venous blood which is so obtained the great majority of clinical laboratory tests can be performed on it, especially as so many biochemical tests are now being adapted to micro-methods. The heel should be thoroughly warmed first and the skin sterilized; then, being held so that the first finger and thumb

FIG. 4.—Milking blood from the heel into a Lyon's tube.

of the left hand can milk the soft tissues (Fig. 4), the point of the heel is sharply jabbed with a needle with a cutting edge; the edge faces towards the toes, and as the needle is withdrawn it is dragged slightly distally to enlarge the opening in the skin. By this means a free flow of blood may be established, which is allowed to drop into a bijou bottle (where any necessary anticoagulant can be mixed with the blood by agitation) or collected into a Lyon's tube (a small glass tube carrying a hole near the top through which air can escape as the blood runs in) or Wright's capsule.

The methods outlined above are all well tried and have been found adequate for their purpose. Successful venepuncture calls for no comment, but failure to obtain blood can be the most exasperating and humiliating experience. Seeking to analyse the reasons for success or failure, there can be little doubt that the main essentials for success are a sharp needle, gentleness, and, above all, not being in too much of a hurry.

Next article on Clinical Pathology.—"Pleural, Pericardial, and Peritoneal Tap," by Dr. Thomas Parkinson.

Refresher Course Book.—The first collection of articles in the Refresher Course for General Practitioners (fully revised) are available as a book containing 55 chapters, price 25s. Copies can be obtained either direct from the Publishing Manager, B.M.A. House, Tavistock Square, London, W.C.1, or from booksellers.

THE DUKE OF EDINBURGH

HONORARY F.R.C.S.

The Duke of Edinburgh visited the Royal College of Surgeons on October 20 to receive the Honorary Fellowship of the College. He was welcomed by the President (Sir Cecil Wakeley) and the Vice-Presidents.

Sir Cecil Wakeley said that every Fellow, Member, and Diplomat of the College shared in their pleasure in conferring on the Duke the greatest honour within their power. The Royal College was closely associated with the Royal Family, and only two years ago the Queen, then the Princess Elizabeth, had been made an Honorary Fellow. He went on to mention the Duke's interest in many branches of science, and said that, over and above the deep affection with which they regarded the Queen and his Royal Highness, they looked upon them both as the personification of a new spirit in their midst—a spirit of confidence and new endeavour. The Senior Vice-President (Sir Reginald Watson-Jones) then presented the Duke for admission to the Honorary Fellowship and the President admitted him.

The Duke of Edinburgh said that he had been interested in a remark by Sir Cecil Wakeley to the effect that surgery was both an art and a science. He did not pretend to know anything about surgery, but, looking at it purely from the layman's point of view, he hoped that the practitioner artist and the scientific researcher would always continue to work in close harmony. He could not help thinking what a bad thing it would be if, with the increasing complexity in all branches of scientific work, the two were to grow apart. He spoke of his great pleasure in following in the footsteps of so many distinguished members of the Royal Family. With them he shared a common admiration for surgeons and surgery, and (they would forgive him for saying) a common ignorance of this noble craft. He added that he had a special satisfaction in becoming an honorary surgeon, because he felt that if ever he came under the surgeon's knife the surgeon would remember that he was operating on a colleague.

THE NEW ACADEMIC YEAR

WESTMINSTER MEDICAL SCHOOL

The inaugural address at Westminster Medical School was given on October 12 at Church House by Lord Radcliffe, Lord of Appeal. Mr. H. M. Clowes, chairman of the school council, presided, and those present included the Mayor of Westminster, the Directors-General of the three armed Services, and Lord Nathan, chairman of the board of governors of Westminster Hospital.

The Inquiring Mind

LORD RADCLIFFE said that he had been reading the reports of the recent First World Conference on Medical Education, and had been impressed by the frequency with which it was stated that the person training in medicine and surgery should not become so immersed in his subject as to forgo the claims of general education and culture. This was a precept not only for those intending a medical career but also for those who were entering other professions, such as engineering. A wider general education must support their specialized knowledge. The leaders of industry were asking for men not merely qualified in their special branches of study but with a wider general background. This was one of the problems of the age. Times had changed since the recognized curriculum could cover all general education. It was quite right for the young person to wish to be a specialist. His wits were alive, he realized the fascination of his particular subject, and dived head foremost into it with all the enthusiasm he could command. It was a good thing to become a man with a craft; it gave him self-confidence and a footing in life which he would otherwise fail to find. The days of universal knowledge