least a number of years. Recurrence after a strict course of medical treatment should be regarded as an indication for surgery, whether or not a strict post-ulcer regime has been followed.

Summary

The case records of 498 patients with chronic gastric ulcer who were treated in the gastro-enterological clinic at St. James's Hospital, London, during the years 1940-6 inclusive are analysed. Adequate follow-up data concerning nearly all were obtained in 1952.

The patients fall into two series, the one treated medically and the other surgically. In the former there are 262 and in the latter 254 patients, 18 appearing in both series.

The criterion for inclusion was the presence of an active chronic gastric ulcer, which in the medical series must have been seen gastroscopically. Patients with additional duodenal ulcers or scars have been excluded.

The ratio of men to women was 3.5:1.

The symptoms may start at any age, most commonly after 30, the average age being 44. The average age at the time of operation in those surgically treated was in the fifties.

The sites of gastric ulcers differ greatly between men and women. In women the site of predilection is the high posterior aspect of the lesser curve, and ulcers in the pyloric antrum are exceptional. In men the distribution along the lesser curve is more even, with a maximum incidence in the mid and lower body, while 10% are situated distal to the angulus.

The results of a fractional test meal, using gruel as the stimulant, are given. The age of the patient influences the level of acidity to only a slight extent, and it is concluded that the presence of a degenerate mucosa rather than the age of the patient accounts for the findings.

A quarter of the patients in each series have subsequently died. There is no predominant cause of death. The problem of the malignant degeneration of chronic gastric ulcers is considered. In none of those who later developed gastric carcinoma is there enough evidence to label them "ulcer-cancers," and it is concluded that if an association between gastric ulcer and gastric carcinoma exists it is due to a preference of both for the same type of degenerate mucosa.

The evidence concerning the incidence of tuberculosis following gastric resection is presented. No conclusion concerning this is reached.

In 28% of patients an incident of manifest bleeding occurred during the course of their ulcer histories. In 16% of these there was more than one episode.

In the medical series it is concluded that although treatment will nearly always heal the ulcer it will keep it healed in only a quarter of the milder or earlier cases. A quarter have symptomatic recurrences more often than once a year, and a further third have had subsequent resections performed for recurrence. Some of the factors which may influence the recurrence rates are analysed.

Three-quarters of those who have had no further trouble have given up dieting, while nearly two-thirds of those with frequent trouble have kept to a diet the whole time.

In the surgical series a very satisfactory result has been obtained in 80%, a moderately successful one in 10%, and a poor result in the remaining 10%.

A high proportion of those with poor results have some other chronic incapacitating disease, most frequently chronic bronchitis.

The choice of treatment for those presenting at hospital with a chronic gastric ulcer is discussed. It is concluded that surgery at present offers the best results, but before it is advocated certain factors, including local operative mortality figures, the severity and length of history of symptoms, and the presence of intercurrent disease, should be taken into account.

Mr. Colin Craig has very kindly allowed us to make use of his late results for patients operated on in 1946. We wish to thank the many people who co-operated in the follow-up of patients, and are grateful to the staff of St. James's Hospital for their interest and advice. We are particularly indebted to Miss Kirwin, who, in addition to secretarial duties, suggested ways and means which led to the tracing of many "lost" patients. The cost of the investigation was met by a grant from the Weir Research Fund.

REFERENCES

- Althausen, T. L. (1949). *Ann. intern. Med.*, **30**, 544.
Brinton, W. (1864). *Lectures on the Diseases of the Stomach*, 2nd ed. Churchill, London.
Capper, W. M., and Butler, T. J. (1951). *British Medical Journal*, **2**, 265.
Ivy, A. C., Grossman, M. I., and Bachrach, W. H. (1951). *Peptic Ulcer*, p. 468. Churchill, London.
Newcomb, W. D. (1932). *Brit. J. Surg.*, **20**, 279.
Palmer, W. L., and Humphreys, E. M. (1944). *Gastroenterology*, **3**, 257.
Pulvertaft, C. N. (1952). *Lancet*, **1**, 225.
Swynnerton, B. F., and Truelove, S. C. (1951). *British Medical Journal*, **2**, 1243.
Tanner, N. C. (1946). *Bristol med.-chir. J.*, **63**, 16.
(1951). *Edinb. med. J.*, **58**, 261.
Visick, A. H. (1948). *Ann. roy. Coll. Surg. Engl.*, **3**, 266.

CANCER AND THE COMMUNITY*

BY

PERCY STOCKS, C.M.G., M.D., F.R.C.P.

Cheshire and North Wales Branch, British Empire Cancer Campaign

In considering the scope of this discussion I have assumed that the Section of Preventive Medicine is not directly concerned with treatment, except in respect of diseases whose spread in the community can be reduced by early treatment of patients infective to others. This absolves me from dealing with the rather controversial questions of how far the community might be influenced to seek medical advice at an earlier stage of cancer and what effect that could have upon ultimate survival. Those aspects of the problem are important to the community and interesting to statisticians, whether in Toronto or London, but, while they may affect the fatality of cancer, they cannot, so far as we know at present, affect the incidence or prevention of the disease.

Attitudes to the Cancer Problem

Some will say that since we can do so little as yet to prevent cancer it is unrealistic to discuss it at all in this Section. Such an attitude seems generally to stem from a fixed idea that cancer is a complication of growing old, or a mode of departing from this life, which is inevitable for a proportion of people in the same sense that arteriosclerosis is inevitable, and I propose to deal with some of the misconceptions mixed up with that idea. According to my observations it is the medical people rather than the lay community who tend to the cheerless view that the best we can do is to try to

*Read in opening a discussion in the Section of Preventive Medicine at the Annual Meeting of the British Medical Association, Cardiff, 1953.

improve the effectiveness of treatment, and that it is futile to try to discover causes of cancer which would be capable of elimination. However, if doctors are at all happy about advances in the treatment of this disease the community certainly is not, and I sense a growing feeling of doubt whether the ordeal of treatment is worth while in many cases in order to achieve a short prolongation of life, coupled with some surprise and impatience because so little serious attempt seems to be made, apart from work on animals, to discover causes with a view to abating them. I find, for example, that, whilst only a very small percentage of people decline to furnish details about their environmental histories, many doctors are hesitant, for one reason or another, in furthering this type of research.

It is sometimes said, generally by young doctors, that cancer is of comparatively slight economic importance to the community, since it is predominantly a disease of advanced life. Comparison with tuberculosis a decade or two ago gave some justification for the view, but it is not true to-day, and it is time that error was exposed. In an address to the British Tuberculosis Association in 1948 (*Tubercle*, 1949, 30, 50) I calculated the years of future "working" life between the ages of 15 and 65 lost to the community in England and Wales through premature deaths caused by respiratory tuberculosis, cancer, and accidents in 1945. It was assumed that if the people who died of one of those causes had not done so in that year, they would have been subject from that time onwards to the ordinary risks of mortality at successive age periods as shown by the 1945 life table rates for the whole population of each sex. The future working years lost as a result of the cancer deaths was 355,000, compared with 435,000 lost through deaths by respiratory tuberculosis and 268,000 lost by fatal accidents. In view of the great fall in tuberculosis mortality since 1945 I have now repeated the whole calculation for the year 1950, using the life table for that year, and the result is shown in Table I. No account is taken of the loss through invalidity prior to death.

TABLE I.—*Estimated Loss of Future Working Life Due to Cancer, Tuberculosis, and Accidents in 1950*

Cause of Death in 1950	International Numbers	No. of Thousands of Person-years of Future Life Lost between 15 and 65		
		Males	Females	Total
Cancer	140-205	212	220	432
Tuberculosis, respiratory	001-008	143	125	268
Accidents	E800-962	211	61	272

The cancer total has increased from 355 to 432 thousands in the interval because the expectation of life based on 1950 death rates from all causes was much better at each age than in 1945, and because the new definition of cancer takes in leukaemia and Hodgkin's disease; this does not indicate a worsening of cancer mortality for both sexes taken together. For accidents the figure has hardly changed, owing to a balance between the reduction in deaths and the increase in normal life expectation. Accidents and cancer are now equally important for males according to this criterion, but for females cancer produces three and a half times the loss due to accidents. For respiratory tuberculosis there has been a great fall from 435 to 268 thousands, bringing the total much below that for cancer. Consequently, cancer has now become the cause of death which is of most serious economic importance when judged from

this standpoint. It is interesting to note that cancer of the lung and bronchus in 1950 contributed 65 to the 212 total for male cancer deaths, and this figure is of course increasing rapidly every year. Cancer of the stomach contributed 33 thousand years for males, so nearly half the total is due to these two forms of cancer in the case of men.

Possibilities of Reducing Cancer Incidence

Another prevalent idea among doctors, though not in the community as a whole, is that it is of little profit to talk about abatement of cancer until pathologists have established its aetiology through experiment, despite the fact that this thesis has been disproved in the history of other diseases. I was once asked by a group of demographers why we could be any more confident in predicting the future trend of the general death rates at particular ages than we are in predicting the trend of the birth rate. My reply was that whilst the latter is influenced by many conflicting desires and considerations which pull in opposing directions and are affected by the social circumstances of the time, conscious efforts to influence mortality in a free democracy are pulling all the time in the same direction, downwards, and, apart from wars, we can count on that continuing.

Furthermore, experience has shown that when the public conscience has become thoroughly aroused about deaths from a particular cause, that death rate has tended to come down whether or not the pathology was properly understood. Two decades ago no one thought that the national infant mortality rate would ever be reduced below 30 per 1,000; but world-wide interest in the problem accompanied by all-round efforts without much aid from new discoveries resulted in rates of 20 being attained by Sweden and 25 by Holland in 1950, and brought down the England and Wales rate to about 28 in 1952. About 1934 the public conscience became thoroughly aroused about maternal mortality, and the rate began to fall at once and continued to do so for years before the sulphonamides and penicillin came along and nearly finished the job. The trouble with cancer is that possibilities of abating its incidence have as yet hardly been regarded seriously except in respect of a few varieties of so-called industrial cancer whose causes are well established.

The search for connexions between human cancer and the countless chemical substances, natural and artificial, which we ingest, inhale, and absorb in the course of our normal lives has begun, however, and we really owe its inception to the observation of Percivall Pott, a surgeon at Bart's in the eighteenth century, that skin cancers of chimneysweeps arose from irritation by soot; there followed, more than a century later, proof that certain mineral oils were responsible for the epitheliomas occurring in cotton-spinners.

Chemical Promoters and Irritants

The number of different agents which have been proved to be carcinogenic to animals since Yamagiwa and Ichikawa first produced true malignant tumours by applying coal tar to the ears of rabbits in 1915 now runs into many hundreds. In 1922 Passey produced malignant growths by painting the skin of mice with an ethereal extract of soot, and in 1924-5 Kennaway was able to produce carcinogenic tars by methods which led to the conclusion that the carcinogenic agent was a complex hydrocarbon of the aromatic series. In 1927 Mayneord discovered that the carcinogenic tars and oils had a characteristic fluorescence spectrum, and Heiger then found that a chemical substance, 1:2-benzanthracene, gave such a spectrum. As a result of further intensive research Kennaway and Heiger were able to demonstrate for the first time that cancer could be produced by a pure compound 1:2:5:6-dibenzanthracene, and in 1933 Cook and Hewett synthesized 3:4-benzpyrene, a carcinogen present in pitch. More recently Berenblum and others have produced experimental evidence that after application of a

carcinogen certain non-specific irritants could determine whether or not tumours appeared and how long they took to appear.

Furthermore, it has been found that the addition of certain fats to the diet of experimental animals increases the incidence of tumours resulting from application of some carcinogens to the skin, whilst, on the other hand, hydrogenated coconut oil reduces the incidence. This work on non-specific co-carcinogens and anti-carcinogens might possibly open the way to cancer prevention on a large scale without waiting until the whole mechanism is understood. Consider the case of bladder cancer, which we know is produced with rather high frequency in men who work with certain chemicals of precisely known composition, while thousands of people who have never handled those chemicals industrially contract the same condition. Surely it is a reasonable inference that those other bladder cancers are initiated by contact in ordinary life with chemicals closely related to the known industrial carcinogens, and whether they do develop may depend on the action of some other chronic irritant. This form of cancer becomes, therefore, a problem for preventive medicine to investigate and is not merely a problem peculiar to one industry.

Statistical Research on Cancer

A third manifestation of the sceptical attitude to prevention in respect of cancer is the excuse that statistics cannot prove causation, and, since they can be made to prove almost anything, why take any notice of them? When statistics show a death rate to be falling the evidence is usually accepted without question and credit is taken for it by all who can lay claim to a part in that happy event. But if statistics show a rate to be rising for a particular disease it is at once objected that medical certification or changing fashion in nomenclature made the figures fallacious. The persistence of the old joke about statistics proving anything is convenient, because when statisticians arrive at unpalatable conclusions it can be said that so long as the evidence is solely statistical nothing need be done except wait for "real," that is to say experimental, evidence. If many thousands of people are dying annually from a disease and statistical evidence indicates that the probability is very high that the main cause lies in an article X, it is hardly sensible to sit back and wait, maybe 20 years, for the experimental evidence. Indeed, unequivocal evidence may never be obtained in a case of a disease which takes a very long time to develop in man after application of an irritant or of an initiator. In such a case the responsibility of advising the community is a serious one, and it is idle to suppose it can be evaded by pretending that statistical evidence alone is of no account.

We are faced with such a situation to-day in regard to lung cancer and tobacco-smoking, and, whilst it is fortunate that tobacco is not a necessity of life, the position is complicated by considerations of finance. In my opinion the proper course is to intensify research on the question from every aspect, and meanwhile inform the community of the statistical findings (as indeed has already been done by the B.B.C.), and say what the risk amounts to if what is suspected is correct, so that each individual can decide for himself or herself whether to take the risk or not. Although statistical studies alone may establish the existence of a real connexion between a factor X and a disease Y without revealing the precise reason for that connexion, the probability that in the absence of X there could be a great reduction in Y may be indicated as so high that some may consider that the use of X involves an amount of risk which it is unwise to take.

Whatever may be the final outcome of the tobacco question, cancer of the lung has now come into the field of preventive medicine. Non-smokers can get the disease and the statistics of deaths suggest that smoke from domestic chimneys may be a factor, so the problem to be solved is whether it arises through inhaling or ingesting carcinogenic or co-carcinogenic substances derived from coal or mineral

oil or used in the cultivation or manufacture of tobacco. Perhaps an initiator is present in one of these and a non-specific irritant in another, and statistical studies now in progress may help to answer this question.

Cancers of the oesophagus and larynx are more frequent amongst men in occupations associated with excessive drinking of alcohol, but there are other factors involved, and the sexes are differently affected. Cancers of the cervix uteri on the one hand and of the breast and ovary on the other are associated in opposite ways with marital history. For these and other kinds of cancer, statistics have indicated that environmental factors have some influence on incidence.

Incidence of Cancer of the Stomach

In the case of stomach cancer incidence is closely related to social and urban environment, and it shows other peculiar local variations for which no explanation has yet been found. Death rates in 1921-30 attributed to this cause showed considerable excess in London and aggregates of large towns at early ages, but the excess was not so pronounced after 65. Standardized comparative figures for years 1947-9, based upon the mortality expected by applying age-specific rates for all rural districts of England and Wales to the populations, are shown in Table II.

TABLE II.—Comparative Mortality Figures for Cancer of Stomach in Certain Large Towns in 1947-9 (With Rates for All Rural Districts of England and Wales as Base)

	Males			Females		
	15-	45-	65+	15-	45-	65+
Liverpool ..	185	156	144	175	128	131
Manchester ..	218	158	132	124	151	123
London county ..	155	114	101	135	109	106
Outer ring of Greater London	133	97	97	112	91	100
Birmingham ..	130	125	120	155	125	116
All rural districts	100	100	100	100	100	100

There has been no diminution of the urban excess previously noted. It is very pronounced in Liverpool and Manchester, and is also considerable in Birmingham, at every age amongst both men and women. In London the excess is evident only at the younger ages, and rates are higher at each age in the centre than in the outer ring. This suggests that some carcinogenic or co-carcinogenic substance, which might be ingested or inhaled in the first instance, is more often encountered in large towns than in rural areas; but we also know that certain rural areas present a large and at present inexplicable excess of mortality compared with other rural districts. I refer particularly to the counties of Wales which touch the sea coast, and will confine attention to the five in North Wales which are now being studied by the Cheshire and North Wales Branch of the British Empire Cancer Campaign with the help of the medical officers of health, the Liverpool and Welsh Regional Hospital Boards, and the University College of North Wales. Comparative indices for cancer of the stomach and breast in North Wales during the 29 years from 1921 to 1949, based as before on the mortality expected from rates in all rural districts at decennial age groups, are shown in Table III. The North Wales counties included are Anglesey, Caernarvon, Denbigh, Flint, and Merioneth; the West Midland group comprises Gloucester, Stafford, Warwick, and Worcester; and the South-east group consists of Kent, Surrey, Sussex East, and Sussex West. In this table, and also in Table II, duodenum is included with stomach.

North Wales showed a rural mortality attributed to stomach cancer about twice the national average at ages under 65, and about 1.8 times the average at ages over 65. Dividing into two periods 1921-39 and 1940-9, the excess is seen to have become only slightly less in the second than in the first period. Cheshire and Lancashire rural districts produced average rates, whilst the West Midland rates were about 11% below the national level, and the South-east

TABLE III.—*Comparative Mortality Figures for Cancer of Stomach and Breast in 1921-49 in Aggregates of Rural Districts of North Wales and Parts of England (With Rates for All Rural Districts as Base)*

Rural Districts of	Stomach				Breast	
	Males		Females		Females	
	15- 65+	15- 65+	15- 65+	15- 65+	15- 65+	15- 65+
North Wales { 1921-39 ..	184	173	224	203	95	91
{ 1940-49 ..	177	165	210	173	97	95
{ 1921-49 ..	182	169	220	192	96	93
Cheshire and Lancashire, 1921-49	101	97	99	104	95	101
West Midlands, 1921-49 ..	89	89	86	89	105	106
South-east, 1921-49 ..	81	83	73	76	105	100
All England and Wales ..	100	100	100	100	100	100

county rates were around 20% below. By way of contrast, breast cancer mortality in the 29 years showed no important differences between these areas, that in North Wales being 5% below average, and that in the West Midlands 5% above.

Justification for Research into Environmental Factors

Suggestions have been made that doctors in Wales tend to certify cancer of the stomach as the cause of death whenever haematemesis or obscure gastric symptoms make it seem feasible, and often with insufficient justification. No evidence that such a practice is in fact more prevalent in Welsh than in English rural areas has been produced. The relative excess is smaller at ages over 65 than in mid-life; and no compensating deficiency can be found in the group of deaths attributed to those conditions likely to be mistaken for gastric cancer, such as peptic ulcer, cancer of the liver, pancreas, and colon, other diseases of the stomach, and pernicious anaemia. Inquiries from a first sample of 100 certifiers of deaths said to be due to stomach cancer in North Wales show that there was necropsy, biopsy, operation, or x-ray evidence for 50%, compared with 65% of 100 deaths so certified in rural districts of Cheshire and Lancashire, and about 60% in the large towns other than Liverpool. If half the cases in which the diagnosis is unsupported by any of these tests are rejected, the North Wales rates would be reduced by a quarter and the Cheshire-Lancashire rates by a sixth, correcting the ratio between them to 1.8 at ages under 65 and 1.6 at ages over 65.

These preliminary statistical tests justify the Cheshire and North Wales Branch of the British Empire Cancer Campaign in their belief that there exists a real phenomenon to be explained, and in their determination to investigate race and residence, work and workplace, soil, air, and water, food, drink, and every conceivable vehicle of carcinogenic substances, or co-carcinogenic irritants. With this line of approach the community of the research area seem to be very much in sympathy. It is something they can understand and welcome because it holds out some hope of prevention, which from the patients' standpoint would be so much better than cure by surgery or radiation, no matter how skilful or high-powered that may be.

Contribution of Medical Officers of Health to Environmental Studies

At the International Conference in London in 1928, organized by the British Empire Cancer Campaign, Sir George Buchanan gave an address on "Collective Public Effort in Dealing with Cancer," which after 25 years is still very pertinent to the discussion to-day. Speaking of municipal work, he said: "The principle to be followed . . . is to secure the closest possible common action between the municipal health department and the authorities of the general and poor-law hospitals, the university if there is one, the voluntary agencies which support social work, and especially the staffs of hospitals and the medical practitioners as well as the local work of the B.E.C.C. where it is in operation. It is no part of the programme that municipal direction should

be made prominent; it is enough for the health department if it can help. Where the medical officer of health is wanted to take the initiative he will do so; otherwise he is content to be one of the cogs on a useful working wheel. Personally, I may express the view that cancer research has much to gain through his instrumentality. . . ."

I would like to take this opportunity of expressing my profound appreciation of the way in which the medical officers of health of the region in which I am working are, together with their assistants, helping the British Empire Cancer Campaign in just that spirit.

Sir John Robertson, at that time medical officer of health of Birmingham, uttered at the same conference some words on which I will close. "Inquiry by the health authority into the facts surrounding each death from cancer has as much value from an educational point of view as it has from an aetiological." How true this is we are finding out in the study I have referred to; education of surgeons, physicians, radiologists, and health officers about the essential gravity of cancer as a community problem is proceeding as a result of the inquiries in progress, and later it may even penetrate to the university. Whether the scientific results we hope for materialize or not, by such education the community will benefit eventually.

CANCER AND THE COMMUNITY*

BY

RONALD W. RAVEN, O.B.E., F.R.C.S.

Surgeon to Westminster (Gordon) Hospital; Surgeon to the Royal Cancer Hospital

The control of cancer is a world problem and one of the big challenges of the present day. Its magnitude requires an all-out effort to reach a solution. With our present knowledge of the disease a great deal can be done to help, but we must not be satisfied until it is under complete control. The object of this paper is to indicate the effects of cancer on the community and the ways in which the latter can co-operate in the fight for its mastery.

Effects of Cancer on the Community

These are of such severity that the disease can be described as one of the greatest scourges of mankind.

World Mortality.—The death rate from cancer varies in different countries, and the figures shown in the Table were published by the World Health Organization in 1951.

Deaths from Cancer by Country per 100,000 Population

Country	Males	Females
United Kingdom (1950) ..	205.8	184.0
France (1950) ..	175.8	169.5
Northern Ireland (1951) ..	161.3	145.1
Netherlands (1951) ..	151.4	147.2
U.S.A. (1949) ..	141.1	136.4
Canada (1950) ..	129.7	123.1
Israel (1951) ..	85.5	105.5
Japan (1950) ..	80.1	74.9

National Mortality.—Reference to the Registrar-General's Statistical Review of England and Wales for 1950 showed that 87,274 persons died from neoplasms, as compared with 72,613 in 1940. During this decade the death rate has fallen considerably from acute infec-

*Read in opening a discussion in the Section of Preventive Medicine at the Annual Meeting of the British Medical Association, Cardiff, 1953.