the author of a film scenario may show you the great surgeon peeling off his rubber gloves after removing the appendix, or the psychiatrist administering to his patient an electric shock, but somehow he does not find romance in the medical officer of health bringing a typhoid epidemic to an end, or in the vital statistician studying the number of deaths from carcinoma of the lung.

It is of fundamental importance to arouse or increase the interest of governments, doctors, and the public in all countries in vital statistics, for without this better technical methods by themselves can achieve little. It is undeniable that at present neither vital statistics nor even public health and preventive medicine excite the interest which their importance deserves and needs, and if we are to change the public attitude to them we must first try to discover why this is so.

Attitude Towards Preventive Medicine

Let us consider first the attitude of the medical profession, since in a sense the doctors are the key to the situation, partly because their direct help is essential, and partly because they are the natural propagandists for preventive medicine, with unique opportunities of obtaining the ear both of peoples and of governments. It might be thought that doctors who spend their lives dealing with the results of disease could be equally active in the sphere of preventive medicine, yet experience shows that this is not necessarily the case. It has to be remembered, first, that doctors who devote themselves chiefly to treatment usually have little time for much else; and, secondly, that the man or woman who is by temperament attracted to the practice of medicine with its concrete problems and personal relationships, though daily made aware of the importance of preventive medicine, is usually less interested in and fitted for what seem the more abstract problems of the aetiology and prevention of disease and the administrative measures for which the latter often calls. Conversely, the medical statistician and the medical administrator have chosen their sphere of work in preference to consultant or general practice.

Co-operation of Statistician and Physician

Hughlings Jackson quoted Herbert Spencer as saying that integration must keep pace with differentiation as division of labour necessitates co-operation, and nowhere is this truer than in preventive medicine. The statistician and the physician are not doing different jobs; they are dealing with different but inseparable aspects of the same job. Let me illustrate this by reference to poliomyelitis. It is a commonplace which is too easily forgotten that clinical observation is the foundation of all knowledge, for the nosological isolation of a disease depends in the first instance almost always upon a combination of clinical and pathological observation; and bacteriology and immunology are built upon the same twin foundations. But when we come to the practical fieldwork of epidemiology clinical recognition of paralytic and, still more, of abortive cases is basic. All this is perhaps obvious, but I stress it because I believe it points the way to increasing the interest of the clinician in vital statistics, which is to convince him that his contribution, just as in preventive medicine generally, must be made, not to what is apt to seem to him an obscure and abstract discipline, but in his own sphere, in which he is the specialist, and where the part he has to play is indispensable because it is irreplaceable.

That is what I meant by saying that the clinician and the statistician deal with different but inseparable aspects of the same job. The statistician depends on the clinician for the accuracy of his data, whether these be diagnoses of disease during life or causes of death, and upon the accuracy of these data his whole work depends. But the statistician is not using statistics for their own sake, but in order that the conclusions which he draws from them may be applied in practice, and in the practice not merely, perhaps not chiefly, of the administrator, but also of the general practitioner and consultant. The relationship between the

statistician and the clinician is thus a symbiosis, and nothing would better arouse the interest of the clinician than information which would convince him of the value to himself of the material which he supplies.

A Great Achievement

Vital statistics dull! Not myself a mathematician, perhaps I can afford to romanticize mathematics a little. It was Pythagoras who set number at the heart of the universe and saw in mathematical relations the foundation of everything that exists. I have often thought that the present age might fitly be described as Neo-pythagorean. Every branch of science has become increasingly dependent upon mathematics, and future historians of thought may well regard it as the supreme achievement of the human mind that it was able to create mathematical symbols of such complexity and subtlety as to provide what seems to be an inexhaustible mode of representing the universe, for on this symbolic representation depends in no small degree our power of action. It is surely of particular interest to us to-day that, if I understand them aright, mathematical physicists have found that when they seek to represent the behaviour of matter in its minutest subdivisions only statistical methods are applicable. And it is to the mathematics of statistics applied to human populations that we look for the source of our power to improve health and prevent diseases which cannot be controlled in any other way. This congress therefore has a twofold purpose—that we may learn from each other, and that we may collectively impress upon the widest possible public the importance for the welfare of the whole world of the task upon which experts in vital statistics are engaged.

SAMPLING IN MEDICAL RESEARCH

BY

COLIN WHITE, M.B., M.Sc.

Lecturer in Physiology, University of Birmingham

A hint that the importance of sampling has not been sufficiently recognized in medical work may be found in the common use of the word "specimen" at times when it would be more helpful to refer to a "sample." The former term tends to imply that one can rely on the material under examination, in the way a botanist usually does when classifying a plant; and that, if mistakes are made, they arise not because the material is atypical but rather because the knowledge of the examiner is imperfect. On the other hand, to use the word "sample" is to remind oneself of the widely recognized fact that samples may be good or bad, fair or biased, representative or misleading; and that all sampling entails, therefore, a sampling error, which is just as great a hazard for the professor or consultant as for the junior assistant.

To emphasize this hazard it may be helpful to examine a few cases in which the results of some experiment or survey have been affected by biased sampling. In the first three examples the bias was important enough to invalidate the conclusions.

Examples of Biased Sampling

Keys *et al.* (1947) have given a lively description of a common error regarding the effect of starvation on the weights of various organs and tissues. It is claimed in many textbooks that when an adult animal is starved the decrease in the weight of the heart is relatively much smaller than the decrease in the weight of the voluntary muscles. This idea arose from a commentator's account of some work by

C. Voit in which two cats, one starved and one well-fed, were used in an experiment. The details given in the original account of the work included the weights of certain organs and tissues in the two animals; and it is easy to calculate from these that, if the well-fed animal is regarded as a control, starvation in one particular cat reduced the weight of the heart by only 2.6% and the weight of the voluntary muscles by as much as 30.5%. Voit did not discuss this point, because it was merely incidental to his problem, but the figures were later picked out by others and illustrated by clear diagrams. Next a mechanism was described to explain how the heart could be maintained on the decomposition products of other tissues. Finally, it was emphasized how easily the results could be reconciled with the philosophical view that this was one more example of the wisdom of the body in maintaining the most important organs. It is difficult to find a medical student to-day who is unaware of these matters, because the story is so interesting and reasonable that it is easy to tell and easy to remember. Unfortunately, it is not true, for at least one of the two cats was atypical, so far as the weights of the organs were concerned. The sample was misleading.

The next example is from the field of public health. At the beginning of this century it was widely believed in Britain that the population was deteriorating in physique. Public opinion had been stirred by the claim that of every five men offering for enlistment in the Army only two were found effective soldiers after the lapse of two years. A committee was set up by the Privy Council to consider this problem. It stated (Report, 1904) that there was little evidence of progressive physical deterioration, but that the men offering for enlistment at that time were not representative of the people as a whole; they were, in fact, a biased sample, with average physique poorer than that of the population from which they were drawn.

The third example is a large-scale experiment on nutrition—the Lanarkshire milk experiment—which was carried out in 1930 and has been summarized and criticized by Student (1931). Over a period of four months 5,000 school-children received $\frac{1}{2}$ pint (425 ml.) of raw milk a day and a further 5,000 received the same amount of pasteurized milk. Another 10,000 children acted as controls. The whole 20,000 had their height and weight measured at the beginning and end of the investigation. It was an experiment planned on a grand scale and carried through with skill and industry; but, unfortunately, the methods of sampling were not rigorous enough. In the first place, the teachers were given some freedom of choice in allotting the children to the control or experimental group; and they exercised it by including too high a proportion of ill-nourished children among those who took milk. A second defect was that each school receiving milk was given either raw or pasteurized, so that the two kinds of milk were not tried out on different pupils from the same school; and, since the experiment was restricted to 67 schools drawn from a rather heterogeneous area, it is quite possible that the group of children drinking raw milk was not comparable to that receiving pasteurized.

In criticizing this experiment we have to remember that in 1930 detailed knowledge of the technique of sampling human populations was confined to relatively few experts. There were no books available on the subject of sampling, and the periodical literature was scanty and rather inaccessible. Moreover, faith in the reliability of large samples was so general among medical workers as to seem almost an unofficial extension of the Hippocratic Oath. But, acknowledging these difficulties, we still have to regard the experiment as a huge waste of effort, and as a warning to future workers of the danger of biased sampling.

In the following example there was no deliberate attempt to select a sample at all; but, in the course of work at a laboratory in Baltimore, Rh-typing was carried out on many people, and it might have been reasonable to suppose that, as the data accumulated, a more and more reliable sample of the population was being obtained. Glass (1950) noticed, however, that of the first 5,000 white subjects examined

15.6% were Rh-negative, but by the time 28,000 subjects had been examined the Rh-negative group had increased to 17.7%. This difference is large, and it is clear that those typed were not a consistently fair sample of the population from whom they were drawn. There was no reason why this biased sampling should have been expected, but bias is so often unexpected that it is wise to take routine precautions against it, if possible, whenever a sample is drawn. Many observers would think it unnecessary to carry out the sort of analysis undertaken by Glass, and would be surprised to find similar bias in their own data.

Random Sampling

The simplest precaution against bias in sampling is to take the sample at random, and, in the absence of background knowledge about the population, we cannot do better than this. It is true that the error involved in random sampling may be rather large; but the important advantages are that the sample itself provides a valid estimate of this error, and that the error can be reduced, if necessary, by increasing the number of cases. Random sampling is a strict process of selection, which is roughly equivalent to drawing lots, and which ensures that each member of the population has the same probability of being selected. Unfortunately, it is difficult to carry out; in fact, the random sample is a theoretical concept which never finds its exact counterpart in the world around us; for even the tossing of coins and the rolling of dice are only close approximations to random sampling.

The methods used for drawing samples in medical research usually fall far below these standards; there is often a factor of expediency involved—for example, when the items required are obtained from a relatively small, and conveniently accessible, part of the whole population. In such a case the risk is clear, but at other times the problems may be subtle as well as serious, for, as Feller (1950) has written, “In sampling human populations, the statistician encounters considerable and often unpredictable difficulties, and bitter experience has shown that it is difficult to obtain even a crude image of randomness.” Nevertheless, in such an important matter it is worth while defining the objectives and describing some common dangers.

As a first step the investigator should review the sampling process before the collection of data is begun, and should satisfy himself that there is no obvious source of error. This review must be penetrating: it is not enough to have a clear conscience about the sampling, because we may have this even when the subconscious is anything but clear; moreover, it is not enough to argue that the sampling is haphazard, because the definition of what is haphazard varies with our understanding of the problem, and we may err through ignorance; or, again, it is far too optimistic to hope that we will obtain a random sample merely because we can identify certain sources of bias, of which some are likely to work in one direction and some in the other. The ideal precaution to take is to ensure that the observer has no freedom of choice in selecting the sample. This is a job in which human beings can hardly be trusted: once the observer is allowed an element of personal choice the selection of the sample is likely to be influenced by the qualities of the objects sampled, and the element of randomness is jeopardized. One procedure which is often feasible in a well-planned experiment or survey is to number the population to be sampled and then use a table of random numbers to determine which members will be selected. The practical details of this method, and various alternative methods, are described by Yates (1949).

Sometimes an investigator, through no fault of his own, may have been unable to obtain a random sample, but he may be familiar with the material under study, and have a clear idea of the type of sampling process that has been used. In this case it may be reasonable to define a population, if necessary a hypothetical one, such that the sample can be regarded as having been drawn at random from this population—for example, certain results obtained in a

survey may not apply to children in general, but may be true of a population which one could define roughly as healthy children; the concept of such a class is perhaps rather vague but may seem adequate for a particular problem. Another example is provided by the report of Kinsey *et al.* (1948) on the sex life of the male. It is clear that the results obtained by these workers are not valid for the male population of the United States as a whole, because there is excessive weighting of certain groups, such as college students, residents in and around Indiana, and people with a professional interest in Kinsey's novel methods of interviewing. Nevertheless the subjects examined may perhaps be considered a random sample from some population that would still be of great interest. In this imaginary construction of populations, however, we should bear in mind the acid comment that certain samples are representative only of themselves.

Probability Sampling

Simple random sampling is the best known of a group of unbiased methods that are described collectively as probability sampling. The essential characteristic of these is that each item in the population has a known probability of being selected for the sample, and, as a result of this, valid errors can be calculated for the estimates made from the data. Descriptions of various types of probability sampling have been given by Yates (1949) and by Deming (1950). One example is stratified sampling. If we attempted to investigate the haemoglobin level of adults in a particular city by simply taking a sample at random, we could not define the value very closely unless we examined a large number of cases. The method would not be misleading, but it would be tedious if accurate information were needed. We could obtain a better estimate from fewer examinations if we stratified the population into groups according to criteria, such as age and sex, which are known to affect haemoglobin values, and then drew a random sample from each of the various strata. The method of stratified random sampling enables us to use background information efficiently. Since some information of this kind is almost always available to those who carry out large-scale surveys, it is rare to find such people using simple random sampling. The latter method is commoner in laboratory work, but is not always used; in fact, the large literature on what is called, perhaps somewhat misleadingly, the design of experiments is simply an account of various methods of sampling.

The advantages of employing some form of probability sampling are considerable. It is true that if we draw non-random samples we may select a group of items which represent the population very well, especially in respect of some quantity such as mean value on which we have deliberately concentrated; but there is a disturbing uncertainty in our conclusions, for we have no objective way of assessing the error made: the usual statistical methods do not apply when the samples are non-random. Of course, any sample at all yields some information about the universe from which it is drawn, but only in the case of probability sampling can the information be assessed in terms of the theory of statistics. This limited value of non-random samples has important effects on scientific procedure—for example, it is wrong to believe that you have only to increase the size of a sample enough in order to be able to draw powerful conclusions. Quality, in general, is more important than quantity; if the sampling is biased an increase in the number of individuals may merely commit you more and more firmly to a false hypothesis. It also follows that when authors are in disagreement the one with the larger number of cases is not necessarily correct.

Other Types of Sampling

A good case can often be made out for using methods of sampling which are known to be biased. Several instances of this occur in the process of measurement, which, in the present context, may be regarded as a method of sampling. If the bias is constant and can be assessed, then it may

simply be subtracted from the end-result, like a biochemist's correction for the blank. If the interest lies in the difference between two measurements, the same amount of bias may enter into each without affecting the results at all. Again, it may be possible to calibrate a biased method against an accurate method, and thereafter use the former, on the ground that it is cheaper, quicker, more convenient, or more likely to give reproducible results—for example, the measurement of the carbon dioxide percentage in "alveolar air" by using end-expiratory samples has the last three of these advantages. Finally, in a given problem the bias of a customary technique may be unimportant. It is fortunate when this is so, for the virtual elimination of bias in measurement—that is, the determination of absolute values—involves the most painstaking care, and often requires facilities that are available in only a few special laboratories.

Another type of biased sampling, in which some expert takes the responsibility of picking the items, has been described by Deming (1950) as judgment sampling. It would be risky to endorse such a method freely, but it is nevertheless common and important in medical research, and is especially useful in the pioneering phase of a problem. The more objective methods can be used when some promising ideas have been developed.

Perhaps the clearest statement of the role of judgment sampling in experimental medicine has been made by Claude Bernard (1865). In a discussion of the method of reporting the results of a group of experiments in which there had been some variations in conditions and techniques, he warned against any attempt to combine the details of several experiments into a kind of average description, and recommended the use of expert judgment: "Quand on a affaire à des expériences complexes et variables, il faut en étudier les diverses circonstances et ensuite donner l'expérience la plus parfaite comme type." A similar idea is expressed in a more colloquial way by the physiologist who inspects the kymograph record and remarks that he has obtained a "publication" tracing.

Judgment samples vary in value according to the ability of the expert; at times this has the unfortunate effect of concentrating attention and debate on the extent of this ability. There is, however, an alternative to argument directed at the men concerned, and that is to regard any conclusions from judgment sampling as tentative until the findings have been checked, preferably by an independent technique.

Quota sampling is a type of judgment sampling which is widely used for commercial and social surveys, but has not, so far, become important in medical work. In this method quotas are allotted to certain important classes—for example, the sample might be planned to contain so many people from the different age groups, so many of each sex, and so many from each social class. Once the quotas have been allotted the interviewers are free to obtain the necessary subjects in whatever way they can. The method may be disastrous, but it does work well at times when the planning is thorough. It has the advantage of cheapness.

Bias in Hospital Statistics

Suppose an investigator wishes to find out whether vascularization of the cornea seems to be associated with deficiency of riboflavin. In order to obtain enough cases for a satisfactory conclusion he might carry out his research on patients in a hospital. A control group of patients without deficiency of riboflavin might be provided from the same hospital, and perhaps steps would be taken to match the two groups in respect of age, occupation, and any other factor that might be known to influence the incidence of vascularization of the cornea. When the survey had been concluded the investigator would be able to draw up a 2×2 table showing how many subjects fell into each of the four possible groups, characterized by one or other abnormality, neither, or both. If the results did not seem clear-cut, perhaps the data would be treated by a standard statistical test such as the chi-squared test.

In an interesting paper Berkson (1946) has shown an important fallacy in this type of argument: his objection to the method can be stated in terms of biased sampling. Consider a population in which there is no association between the sign and the disease under study—that is, the percentage of people with vascularization of the cornea is the same in the deficient group as in the rest of the community. We may now imagine that the patients in certain hospitals are drawn from this community. If the probability that a person entered hospital differed according to whether he was deficient in riboflavin or not and also according to whether he had vascularization of the cornea or not, then the hospital population would not reflect the true incidence of these factors, and would give a false picture of the problem under investigation. The results, however carefully developed, would be an artifact. The true relationship between the two factors would sometimes be stronger and sometimes weaker than that indicated by the sample of hospital patients, for the bias might tend in one direction or the other according to the way the patients were drawn from the four groups in the community.

This is a far-reaching criticism extending, as it does, to many current projects in medical research; but it must be modified in one important respect. Suppose that the presence of duodenal ulcer influences the probability that a person would be admitted to hospital, but the nature of his blood group does not. Then, on these assumptions, the patients in hospital could be used to obtain a valid estimate of any hypothetical association between blood group and the incidence of duodenal ulcer. If the association were zero, for example, then such a state would be reflected in a group of hospital patients just as well as in a group taken from the community. The saving factor here is that, since the blood group is without effect on the probability that a person would enter hospital, the mere concentration in hospital of large numbers of patients with duodenal ulcer would not distort the true relation between the blood group and the disease. Such distortion will not occur unless both of the factors being examined affect the probability of becoming a hospital patient.

The problems raised by the use of hospital statistics may be illustrated clearly by a reference to thromboangiitis obliterans. It used to be accepted that this disease occurs chiefly in Jews; but, more recently, there have been numerous reports that contradict such a view. It is now recalled that Buerger and many of the other early investigators of thromboangiitis obliterans worked at hospitals where the patients were predominantly Jewish; so that the supposed racial predisposition to the disease was an error due to biased hospital statistics.

Bias in Post-mortem Statistics

The bias to which Berkson has drawn attention in the case of hospital statistics is even more likely to occur when statistics are collected from post-mortem records. The morbid anatomist probably has little to learn from current statistical methods of testing hypotheses, because, in general, his material is necessarily obtained by methods of sampling which invalidate the tests of significance that a statistician employs. An early paper by Stocks (1924) shows that he sensed this point, though he had no method of neutralizing the bias which it might cause, for, as Berkson has pointed out in regard to similar problems of hospital statistics, "There does not appear to be any ready way of correcting the spurious correlation . . . by any device that does not involve the acquisition of data which would themselves answer the primary question."

Post-mortem results may be equally subject to bias when they involve measurements. Marshall (1951) has carried out t-tests to determine whether the heart weight observed after death differs according as the subjects have normal or aberrant renal arteries. If we wish to apply these results to the population in general, however, we have to remember that no care or skill on the part of the pathologist can ensure that he obtains a random sample of material from

the general population; his sample is determined by factors over which he has little control. Whenever an investigator is forced to draw his sample, not from the population in which he is interested but from some pre-selected portion of it, he is in danger of obtaining a biased group. In effect there are then two stages of sampling; the investigator controls the second stage, and perhaps carries out his work with scrupulous care; but this may be of no avail if there has been bias in the first stage.

To illustrate this point, suppose that someone takes a random sample of schoolchildren in a certain area and finds out in each case how many there are in the family to which the child belongs. The sampling of the schoolchildren may fulfil all the requirements of random selection, but the data obtained will be biased if they are used to give an estimate of the average number of children per family. It is clear that various families have different chances of being included in the sample; an extreme case is that of the childless family which has zero chance of being selected.

Fallacy of the Interesting Cases

Biased sampling may arise when we concentrate our attention exclusively on the changes that develop in some interesting section of the total group. Consider the case of children who have either unusually tall or unusually short parents. These children, when fully grown, have heights which, on the average, are not as extreme as those of their parents; but it would be fallacious to conclude from this interesting but biased sample that the height of individuals is converging towards some modal value. Adults of the more extreme sizes are derived in part from parents belonging to the middle ranges, and when these parents are fairly represented in the sample it is found that the distribution of the height of adults remains fairly constant from one generation to the next.

The mistake of following the progress of members of the tails of a distribution, and forgetting about the rest, should not be made when the data are available for the complete series. But it sometimes happens in medical research that only the atypical subjects have been given a second examination. This may occur because they excite greater attention than their fellows and appear to repay closer study; they provide welcome relief from the dullness of examining a series composed mainly of normal people.

Some data provided by Toverud (1950) may be used in illustration of this problem. He was interested in the nutrition of pregnant women, and, as part of his investigation, he determined the incidence of anaemia among them. Of 1,077 women examined, 120 had haemoglobin values of less than 12 g. per 100 ml. These 120 women were treated by "regulating the diet and adding iron" and, on subsequent examination, it was found that 82 of the 109 available had haemoglobin values of more than 14.4 g. per 100 ml. This might be taken as evidence that about 11% of the pregnant women required treatment for anaemia and that the treatment used was successful in most of the cases. But the design of the investigation was such that no information could be obtained about those fluctuations in the haemoglobin level which were independent of treatment. Some of the normals, if re-examined, might have been found to have developed temporary anaemia; and some of the anaemic women might have recovered without treatment. These possibilities could have been investigated by making more measurements on the normal group and perhaps by withholding treatment for a time from the supposedly anaemic group—that is, by concentrating less on the interesting cases.

Mass radiography, and other similar preliminary tests, can be regarded as methods of separating off an interesting section of the population for more detailed study. In the initial stages, at least, it is essential that a sample of those who are classified as negative in the screening test should be given the same detailed study as the positive cases receive. Only in this way can it be found whether some subjects have been incorrectly diagnosed as negative. If you direct

attention to the positive group alone, you can never be certain that the screening process is sensitive enough for your needs—that is, you can never find out whether the test fails to detect early cases.

"Ex Post Facto" Experiments

A great deal of information relevant to problems in medicine has been collected in the form of routine records by agencies such as public health departments. Sometimes these data have been gathered without any special research project in mind, but when they are reconsidered they seem to be pertinent to some problem. These data then constitute what sociologists refer to as an *ex post facto* experiment.

Information obtained in this way is often illuminating, but there are certain limitations to its use, and these are chiefly concerned with the problem of biased sampling. Is it possible to define a universe of which the cases under consideration can be regarded as a sample, and has the sampling been random? If these questions cannot be answered satisfactorily then the standard methods cannot safely be used to test hypotheses concerning the data, and any conclusions drawn may lack objectivity.

REFERENCES

- Berkson, J. (1946). *Biometr. Bull.*, 2, 47.
 Bernard, C. (1865). *Introduction à l'étude de la médecine expérimentale*. Paris.
 Deming, W. E. (1950). *Some Theory of Sampling*. New York.
 Feller, W. (1950). *An Introduction to Probability Theory and its Applications*, vol. 1. London.
 Glass, H. B. (1950). *Cold Spr. Harb. Symp. quant. Biol.*, 15, 33.
 Keys, A., Henschel, A., and Taylor, H. L. (1947). *Amer. J. Physiol.*, 150, 153.
 Kinsey, A. C., Pomeroy, W. B., and Martin, C. E. (1948). *Sexual Behaviour in the Human Male*. Philadelphia.
 Marshall, A. G. (1951). *Lancet*, 2, 701.
 Report of the Privy Council upon Physical Deterioration (1904). *Lancet*, 2, 390.
 Stocks, P. (1924). *Biometrika*, 16, 364.
 Student (1931). *Ibid.*, 23, 398.
 Toverud, G. (1950). *Milbank mem. Fd Quart.*, 28, 7.
 Yates, F. (1949). *Sampling Methods for Censuses and Surveys*. London.

ADRENAL PSEUDOHERMAPHRODITISM

BY

L. R. BROSTER, O.B.E., M.Ch., F.R.C.S.
 F.A.S.A.(Hon.)

JOCELYN PATTERSON, M.Sc., Ph.D., F.R.S.Ed.

AND

B. CAMBER, M.B., Ch.B.

(From Charing Cross Hospital)

Adrenal pseudohermaphroditism, or pre-pubertal virilism, is a syndrome characterized by the masculinization of the female. It is associated with simple enlargement and pigmentation of the adrenal cortex, and the excretion of large amounts of 17-ketosteroid in the urine. The condition is clearly differentiated from true hermaphroditism and the intersexes in two respects. Whereas in both the latter there is a persistence of Wolffian derivatives and no rise in 17-ketosteroid excretion, the reverse is true of adrenal pseudohermaphroditism.

In relation to the aetiology of the disease it is of interest to note that Professor Barr, of London University, Ontario, has examined portions of the skin taken from some of our patients and found in every case the nuclei were of female morphology (personal communications). The available evidence does not advance us beyond the vague generalization that the condition is due to secondary hormonal influences acting upon a genetically determined female embryo during the early stages of foetal development. The critical period of any hormonal action would seem to be between the 11th and 21st weeks—that is, between the time of disappearance of the Wolffian derivatives and

the completion of Müllerian development. Whether the stimulus derives from the mother or arises from some factor inherent in the foetal endocrine system are questions which have yet to be determined. Whatever theory is advanced, it must take into account the strong familial element associated with the condition, the evidence for which is cited later.

Our work on this problem of the masculinization of the female began in 1926; since then we have collected 37 cases belonging to the group of congenital pseudohermaphroditism, and the present communication is based on clinical and biochemical data compiled on this series. Previous communications have dealt with a number of the early cases (Broster and Vines, 1933; Broster *et al.*, 1938), but within the past two years many of these, together with more recent patients, have been subjected to a closer and more critical study made possible through the collaboration of Dr. J. Swale, Professor J. Patterson, and Dr. B. Camber.

General Features

Symptoms.—The general features depend on the age of the patient and the severity of the lesion. In infancy the earliest sign is an enlarged clitoris with a persistent urogenital sinus. Later there may be precocious puberty as indicated by premature growth of pubic followed by axillary hair and raised ketosteroid output. As the patients grow up the somatic development is in advance of the chronological age and the short stature is characteristic. If, however, the patient is not seen until the age of normal puberty the diagnosis is made clear when the secondary sex characters of the female do not appear and facial and somatic hair of the male type grows and the voice becomes deep. More recently it has become known that patients suffering from congenital hyperplasia may die in early infancy or childhood from adrenal failure, and in six instances a history was obtained of episodes suggestive of Addisonian crises.

Hirsuties.—Except in infants, and one adolescent, hirsuties of male distribution has been a prominent feature in all our cases. In two of the older women temporal recession and apical baldness were present.

Breasts.—The great majority of our patients have shown no breast development at puberty. In only eight were they rudimentary, and some of these had received oestrogen therapy.

Sex Development.—On examination the clitoris is found to be enlarged and curved downwards. From the area of the fraenum two small folds of skin gradually diverge to encircle a small orifice which is the persistent urogenital sinus. No urethral orifice is visible externally. The cervix and uterus are small and generally not palpable. In only one patient was there a small urethrovaginal fistula. At laparotomy the ovaries have been in their normal position and have not shown any marked macroscopic changes. Some appear normal, while others are fibrotic and cystic. Histologically they show oogenesis and atretic follicles as well as cysts. Three patients about the age of 6 showed numerous ova, atretic follicles, and numbers of cysts typical of isosexual precocity. One of these (M. Ba.), who had normal follicles with some abnormally mature ova, showed a sexuality much in advance of her age. As is reported later, this sexual precocity was related to signs of undue maturity recorded by the electroencephalogram. A few reports recorded luteum cysts, and in one case there was no abnormality apart from suspended activity. Primary amenorrhoea is almost invariably present, although two patients had a slight show after puberty.

Clinical

History.—Significant features in the family histories of 50% (18) of the patients in this series were familial endocrinopathy, maternal disorder during pregnancy, mixed racial origins, or the presence in the individual of associated congenital abnormalities. With adrenal disease in general we have been struck by the frequency with which it occurs