Published in final edited form as:

Soc Networks. 2007 May ; 29(2): 279–299.

Partner Naming and Forgetting: Recall of Network Members⁺

David C. Bell, Ph.D. ^{*}

Department of Sociology, Indiana University-Purdue University Indianapolis

Benedetta Belli-McQueen, M.S., M.A., and

Center for Success and Independence, Houston, TX

Ali Haider, B.B.A.

Karachi, Pakistan

Network studies, whether structural studies of social networks or relationship studies of ego networks, depend on the accuracy of informants' self-report data (Marsden 1990; Tourangeau et al 2000). In turn, the accuracy of informants' self-reports will determine the accuracy of inferences drawn from such data (Brewer & Webster 1999; Stone et al 1999; Wasserman & Faust 1994). Sometimes all potential network members are known to the investigator (e.g., a list of employees for a company network). But in many cases there is no list and the investigator has to rely on informant free recall of alters.¹ In this paper, we will examine the accuracy of free recall using a unique dataset in which informants were interviewed about their networks up to 13 times at 3-month intervals. We will address two research questions. The first has to do with describing and explaining the forgetting of alters. The second question has to do with the relative accuracy of global estimates and naming as measures of network size.

Global Estimates vs. Naming

The simplest and most direct measure of ego network size is a global estimate, a single number produced by the informant as an estimate of the number of alters in a network (what Brewer et al 1998 refer to as a "direct estimate"). Psychologists of survey response as well as network researchers have identified several ways an informant may produce a global estimate of network size. First, the estimate may be produced by implicit enumeration (Blair & Burton 1987; Burt 1987; McCarty et al 2001; Sudman & Schwarz 1996; Tourangeau et al 2000). In this process, the informant mentally enumerates the alters and reports the count. Reports summarized by Tourangeau and colleagues (2000) suggest that, when informants provide a global estimate of the frequency of past behaviors, they use implicit enumeration about one fourth of the time. Implicit enumeration appears to be more common in estimates of numbers of alters. When informants are asked to estimate the number of lifetime sex partners, they use implicit enumeration 70% of the time (including 28% who have kept a running tally in memory) (Brown & Sinclair 1997, cited by Tourangeau et al. [2000]). In one network study, two-thirds

⁺Work on this paper was supported in part by grant DA08989 from the National Institute on Drug Abuse, principal investigators Isaac D. Montoya and David C. Bell. The authors would like to thank Devon Brewer, Matt Salganik, and anonymous reviewers for comments on a previous draft. Opinions expressed here are solely those of the authors. This research was conducted at Affiliated Systems Corporation, a private research organization in Houston, Texas.

* Corresponding author: David C. Bell Department of Sociology, Indiana University-Purdue University Indianapolis (IUPUI), 425 University Blvd. Indianapolis, IN 46202-5140, Telephone (317) 278-1336, email: dcbell@iupui.edu

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

¹Since our concern in this paper is with informants' self reports, when we say "network" without qualifier, we are referring to an informant's ego network. We will use "alter" to refer in general to a person to whom an informant has a link of some kind. When we specify a particular kind of relationship, we refer to the alter as a "partner" or "friend": sex partner, drug use partner, close friend.

of informants reported using implicit enumeration to arrive at their global estimates, especially for small numbers of alters (Brewer et al 1998).

In addition to implicit enumeration as a method by which informants directly estimate network size, Tourangeau and colleagues (2000) describe several other estimation methods, of which rates and impressions are relevant to network size estimation. Global estimation may be based on informants' awareness of an average rate at which events occur (Blair & Ganesh 1991; Menon 1993), which is then converted mentally into an estimate of number of alters. Another general method of global estimation is for an informant to rely on a general impression (Tourangeau et al 2000), which is essentially a matter of guessing. Impressions or guessing are often detectable as "heaping," in which estimates tend to be clustered around common numbers such as multiples of five or ten (Brewer et al 1998; Conrad et al 1998; Huttenlocher et al 1990). Global estimates of the number of sex partners are likely to use round numbers when there are more than 8 partners (Tourangeau & Smith 1998).

As an alternative to asking for a global estimate of ego network size, a researcher may ask a study informant for a list of partners, an explicit naming of the network. Compared to global estimates, the naming method provides greater depth of information because the investigator can ask about each alter, but this depth comes at the expense of valuable interview time.

Both naming and global estimates are self-reports, and thus both are liable to potential biases. Both often involve the same recall process. As we have seen, most people who make global estimates of number of alters say that they carry out an implicit enumeration. Naming makes this enumeration explicit. Naming can lead to informant exhaustion as informants come to believe they have spent "enough time" retrieving names (Krosnick 1991). However, since naming is more explicit and in network research is often under the gentle coercion of an interviewer, naming is inherently produced by enumeration (see a similar argument on interviewer demand in Schrppler 2004).

Recall and Forgetting

Whether in naming or in global estimates, there are a number of potential biases in self reports. An informant may guess at the size of their network without trying very hard to remember specific alters; they may forget one or more alters; they may omit alters because they do not know their names or identities; or they may intentionally or unintentionally conceal (or invent) some alters.

Alter Recall

When informants give a global estimate or when they are asked to name members of their ego network during interviews, the quality of the data collected depends on informants' ability to recall and report their alters. However, humans can be fragile measuring instruments. Studies comparing free recall to recognition of names from a list find that recall can be as low as 10% (Hammer 1984) and as high as 90% (Brewer & Webster 1999; Sudman 1988). Using partner concordance as a way to identify successful recall, studies have found varying levels of agreement between informant and alter (52–85%) for high school buddies (Alexander & Campbell 1964; Kirke 1996), adult best friends (Laumann 1969), and intimates (Hammer 1984; Shulman 1976). Research on partner concordance involving behaviorally based network alters have found agreement rates of 93% for sex partners, 82–83% for non-injection drug use partners, and 76–90% for drug injection partners (Adams & Moody 2005; Bell et al 2000). Studies using objective and independent measures of interaction have suggested that, for some kinds of relationships, respondents can recall fewer than half of all alters (Bernard & Killworth 1977; Bernard et al 1981; Bernard et al 1982; Milardo 1989). Some of this research is summarized in Brewer (2000) and Marsden (2005).

Forgotten Alters

Some partners may be unintentionally omitted when an informant attempts to enumerate alters (Brewer et al 1999). Omission can occur because of forgetting, a result of memory decay over time (Anderson 1990; Sudman & Bradburn 1982; Wixted & Ebbesen 1991). For example, memory decay may be involved when self-reported frequencies of partner interactions tend to fluctuate more the greater the interval between event and recall (Catania et al 1993; Kauth et al 1991). We consider forgetting in greater detail below.

Anonymous Alters

In addition to forgetting, where an informant has previously known the requested information but has forgotten it, are those cases where the informant does not know the information. Ego network interview name generators usually ask for a name, nickname or initials. If the researcher is interested in friendships or advice networks, anonymity may not be a problem. Friendships are developed over time, so the case of a friend whose name is unknown is ordinarily rare. Similarly, people usually receive advice from others they know and respect. However, some theoretically and personally meaningful relationships may form with persons whom an informant cannot identify by name. For example, a one-time drug partner met in a park, or a one-night stand met at a bar may be alters whose names are not reliably stored memory (Sudman & Bradburn 1982) and may never even have been communicated. Informants are likely to omit such relationships from a global estimate or a list of names unless specifically asked about alters whose names are not recalled.

Sources of Forgetting

Leaving aside the problem of anonymous alters whose names are not known, there are a number of potential factors that can tribute to forgetting.

Network Size

Forgetting can depend on network size (Brewer et al 1999; Brewer & Webster 1999). This principle suggests that when one has only one or two alters of a given type, it is relatively easy cognitively to keep track of them. However, as the number of alters increases, it appears to become increasingly difficult to recall additional alters. Behavioral estimates based on impressions are more likely as the number of events increases (Blair & Burton 1987; Conrad et al 1998).

Behavioral Specificity

In addition to depending on network size, recall of alters also clearly depends on the type of behavior that constitutes the relationship (Sudman & Bradburn 1982). Network name generators may ask about behavior-based relationships or about affective relationships (Campbell & Lee 1991; Milardo 1992; van der Poel 1993). In their study using a two-year time frame, Brewer and colleagues (1999) found higher levels of forgetting for drug injection partners than for sex partners. From the point of view of recall, behavioral links have the advantage that the distinction between link and non-link is relatively clear: either a behavior occurred or it did not. Kogovsek and Ferligoj (2005) report that responses to questions about behaviors of alters (frequency of contact) are more consistent and more accurate than emotional questions (closeness) (see also Clair et al 2003; Harvey et al 2004). Fischer (1982), for example, found a great deal of variation in how informants interpreted the term “friend.” Thus, an imprecise concept like friendship (Brewer & Webster 1999) or acquaintanceship (Brewer 1993; Sudman 1985) may have higher levels of “forgetting” because of the indeterminacy of the threshold (where does the difference between “friend” and “not-friend” lie on the dimension of emotional closeness) or the indeterminacy of the alter’s relation to the threshold at any

interview (is the emotional closeness to this alter above or below the cut-off). Furthermore, informants may have different thresholds for a given term (Feld & Carter 2002; Kashy & Kenny 1990). Against this pattern, Kirke (1996) found a high level of partner concordance over the terms “friend” and “pal” in her sample of adolescents in Ireland.

Salience

Certain name generators may refer to types of relationships that are highly salient and involving, and thus easier to remember than others (Sudman & Bradburn 1982). Along these lines, Marsden and Campbell (1984) suggest that emotional closeness is the best conceptualization of strong ties. Stronger friendships relationships are more likely to be recalled than weaker ones (Brewer 2000; Campbell & Lee 1991; White & Watkins 2000). Less salient and less absorbing interactions have been found to involve greater amounts of forgetting, such as telephone conversations (Milardo 1989) and other day-to-day communications (Bernard & Killworth 1977; Bernard et al 1981). Stable relationships are more easily recalled (White & Watkins 2000), as are higher frequency relationships (Burt & Ronchi 1994). Because of salience, rare events, high value events, and continuing events are most often recalled (Sudman & Bradburn 1982). Thus, occasional partners and partners not very close to the respondent may be more likely to be forgotten. In their study using a two-year time frame, Brewer and colleagues (1999) found higher levels of forgetting for drug injection partners than for sex partners (see also Bell et al 2000).

Time Frame

Network self-reports appear to be compounded of specific memory and enduring cognitive map (Freeman & Romney 1987). Specific memories decay over time, as discussed above. Thus, recall of specific events has been found to be worse for events distant in time than for those close in time (Bernard et al 1984; Graham et al 2003). In compensation for memory decay, people often form cognitive structures in memory, and these structures may be enduring and powerful (Killworth & Bernard 1976). My memory of whether we have met in the past week will be built up in part of my specific memory of individual interactions with you and my cognitive map which says that we “always” meet for lunch on Wednesdays. Recalled relationships may thus represent the long-term pattern of social relationships as recorded in the cognitive map (Freeman et al 1987). Implicit enumeration of behavioral counts is less likely as the time frame increases (Burton & Blair 1991). Blair and Burton (1987) report that episodic enumeration rarely occurs over time frames as long as 6 months. Bachman and O’Malley (1981) found that high school students reported higher rates of drug use for short time periods and tended to underestimate drug use over longer time periods. This pattern of time dependence in global estimates has been referred to as “telescoping.” Kogure and colleagues (2001) found that forward telescoping (overestimating recent events by including events from outside the time frame) occurred in a time frame of under two years, while backward telescoping (underestimating by considering only part of the time frame) occurred for events more than three years old (see also Bradburn 1983; Prohaska et al 1998). This pattern has been suggested to occur because persons have a sense of a “customary” or typical network (Brewer 2000; Freeman & Romney 1987; Hammer 1984; Sudman & Bradburn 1982). For example, to the extent that alters in one’s network may have episodic interactions, a global estimate with a short-term time frame may include some of the “customary” network members, even though there were no interactions within the time frame. Some empirical evidence has suggested that measures that tap into typical networks may have more predictive validity than short term measures (Chang & Krosnick 2003). When persons participate in joint drug injection or sexual behaviors, they are more likely to agree on their six-month behaviors than on their one-month behaviors (Bell et al 2000).

Implementation Issues in Collecting Network Data

The information that informants provide depends on their knowledge and their motivation to be accurate. Since informants are engaged in a process that in most cases they have not chosen (they have been recruited by strangers to provide information to other strangers for reasons known only to strangers), their commitment to absolute accuracy must be considered to be problematic. Informants are usually willing to give some effort to the process of providing information, but they cannot be expected to be exhaustive in their accessing of mental information, especially with interview fatigue from long interviews (Knoke & Kuklinski 1982).

In addition to the limits of informant effort and ability to be accurate, informants may sometimes be motivated by social desirability to provide inaccurate (biased) information. Research on computer-assisted interviewing has suggested generally that sensitive behaviors may prompt reticence from informants (Des Jarlais et al 1999; Kissinger et al 1999; Kurth et al 2004), although some studies have found mixed (Perils et al 2004) or negative results (Couper et al 2003). Sexual information may be especially problematic. It has been suggested that people feel threatened when asked to report on their sexual activities and thus hesitate to reveal their sexual behaviors to others (Bradburn 1983; Catania 1999; Catania et al 1986; Herold & Way 1988). On the other hand, other investigators have found that respondents can be quite forthcoming about their sexual behaviors (Kinsey et al 1948; Kinsey et al 1953; Laumann et al 2004; Laumann et al 1994). When rapport with interviewers is high, respondents have been found to reveal 93% of their sex partners (Bell et al 2000). Just as social desirability has been discussed as a possible influence on validity in sexual behavior self-reports, it has also been discussed in relation to drug and alcohol use. A study of drinking during pregnancy indicated that women reported higher during-pregnancy drinking rates 5 years after delivery than they had reported during pregnancy (Ernhart et al 1988). For drug use, underreporting is more severe depending on the stigmatization of the drug. Marijuana use appears to be reported more validly than cocaine use in the general population (Harrison 1995) and in arrestee populations (Fendrich & Xu 1994; Mieczkowski et al 1991).

Method

Data for this study were collected as part of the longitudinal [study], an investigation of relationships between a community (non-clinical) sample of drug-using and nonusing persons and their partners, and the effects of these relationships on HIV transmission risk. The sample for the [study] was drawn from [description omitted]. The sample had been recruited through random walk and peer-driven recruitment methods [citations omitted]. One of the consequences of this study design, in which informants were interviewed over a period of up to 7 years, is that study informants tended to be stable in residence and in relationships. Informants' relationships were not limited to the kind of short-term relationships that often predominate in transient populations such as college students. Thus, non-family relationships had a mean duration of some 9.9 years and a median of 7.0 years. Sex relationships had a median length of 1.8 years; drug use relationships, 1.6 years.

At baseline each study informant was informed as to the actual purpose of the study, types of information requested, the amount of time required, and their right to refuse participation. Interviews were conducted in English in a private room in a field research center. All informants were over the age of 18, verified by a driver's license or similar document. All informants gave written informed consent and were reimbursed \$25 for their time for each interview. Study procedures were approved by the Affiliated Systems IRB.

The complete sample of 202 informants is described in Table 1. Slightly more than half the sample were male. More than half were African American, about a fourth were Hispanic, and one-sixth were Anglo. The majority of informants were over 40. Almost half the sample had not been employed in the six months before the interview; almost a fourth of the sample were identified behaviorally as gay, lesbian, or bisexual. An informant who ever described sex with a partner of the same gender at any wave of the study was coded as gay/lesbian/bisexual. In addition, informants who described no sex partners at any wave were coded as not sexually active. Most of the eleven informants who were not sexually active reported that they were not in the mood for sex; three informants used abstinence to protect against disease. Almost 84% of the sample were drug users, defined as users of cocaine, heroin, or methamphetamine. Nearly half had used marijuana, more than half had used crack cocaine, and about a fourth had used powder cocaine in the previous 30 days.

The [study] collected longitudinal data from informants at three-month intervals for over three years. Attrition, incarceration, involvement in drug treatment, and other contingencies affected the number of subjects available for interview at any given wave. In all, 193 of the 202 informants participated in at least one follow-up interview (so that forgetting could be detected). Over half of the sample were interviewed at least 11 times out of 13 waves.

Measures

Individual-level data—At each wave, the data included a core instrument that collected individual-level demographic information and information on sexual activity and injection/non-injection drug use (crack cocaine, powder cocaine, heroin, methamphetamine, and other drugs). More detailed questions were asked about drug injection and sex because of the study's focus on HIV risk. Informants were asked, "In the last 30 days, how many men/women did you have any kind of sex with?" (male and female partners were asked about separately). "Sex" was defined as anything that either partner thought of as sex. Informants were asked how many of these men/women were persons whose name the informant did not know or could not remember. In retrospect, we wish we had asked about global numbers of drug use partners and about those whose names were not known, but we failed to do so in this study.

Relationship-level data—The instrument employed several name generators, where informants were asked to name all their sex and drug use partners in the previous 30 days, as well as close friends ("the people you consider to be close friends"). Drug use partners were alters with whom the informant used drugs (specifically crack/cocaine, heroin, or methamphetamine) in the same place at the same time. Alters could be drug partners with or without sharing drugs or sharing equipment. Drug injection partners and non-injection drug partners were named. For each name generator, the informant was prompted for any more alters in the category until the informant could name no more. An alter could be named in multiple categories (e.g., both close friend and drug partner).

Data collection was conducted by a trained interviewer using computer-assisted personal interview (CAPI) methods. Computer-assisted interviewing allowed us not only to solicit alter names, but also to compare the list of names spontaneously recalled to a list of previously named alters, so that each successive wave of data collection provided an opportunity to detect whether a study informant forgot a previously named partner. After all alters were spontaneously named, informants were presented with the name of each alter named at any previous wave but not named at the current wave. In this way we were able to determine why any alter had been omitted from the list of spontaneously generated names. Each such alter was coded as "forgotten" (if alter was currently a partner) or as "not a current partner" (if alter fell into the categories of "haven't seen in 30 days, not in contact now," "haven't seen in 30 days, still in contact," or "have seen, but not a partner now"). Similar procedures for identifying

forgetting have been used in other longitudinal studies (Morgan et al 1997; Wright & Pescosolido 2002).

There is a potential problem in free recall of alters across multiple waves of measurement, in that informants may use different names for the same alter. In our procedures, these name conflicts were caught when the informant was presented with the previously used name and identified this person as the same as one of the spontaneously named alters. Our software then assigned the previous alter's ID number and combined the two names for use in subsequent waves. Another issue occurred when an informant reported that they could not remember the previously named alter: the alter was coded "not a partner." Occasionally an informant would report that an alter had died. We recorded this, but we learned from experience not to remove an alter from the names list until they had been reported as dead three times.

Of course, there were occasions where a partner was not named because the informant did not know the alter's name. One of the goals of the analyses we report here is to explore this issue.

A series of questions was asked about each alter. Connection was operationalized by a set of 6 items asking how much the participant trusted, respected, cared about, did things for, was listened to, and told important things to the partner. Responses were recorded on a scale from 0 to 10. These responses were combined (as a mean) into a single measure of the level of emotional connection felt by the participant toward the partner (alpha reliability = .92). Connection is a measure of emotional intensity within a relationship. In addition, for sex relationships, we asked whether sex was "for sex" or "a close emotional connection."

Interaction frequency was measured by the number of days in the past 30 days that the informant and partner had seen or talked to each other. Interaction frequency is a measure of behavioral intensity of a relationship. Relationship duration was measured for close friends as the number of months they had known the partner. Similarly, for sex partners, informants reported the length of the sex relationship, and for drug use partners the length of the drug-using relationship. Relationship type. For each relationship, informants reported whether the partner was a spouse, other family member (child, parent, other family), boyfriend or girlfriend, best friend, or acquaintance (friend, acquaintance, drug buddy or other).

Over 13 waves of the study, almost 14,000 observations were collected on various social network alters who were active partners at the time of the interview; in addition, there were over 19,000 observations where alters were reported as not having been partners in the previous 30 days. Some of these alters were never named after the first mention; some were named in every subsequent wave after the first mention; and others were named intermittently. Of the 191 informants who reported any sex, 189 described sex relationships with a total of 830 partners, of whom 305 were sex partners in more than one wave (four informants gave global estimates of one sex partner for at least one wave, but did not have a sex partner whose name they knew); 159 informants described drug use relationships with 1,413 partners, of whom 461 were drug use partners in more than one wave; and 200 informants described close friendships with 3,522 alters, of which 2,307 were named as close friends in more than one wave. An examination of alters shows that 63% of sex partners were described as sex partners only once, even though the informant was reminded of them at each subsequent wave. Similarly, 67% of drug use partners and 34% of close friends were described only once. Only 16 alters were described as sex partners in all 13 waves even though 63 informants were interviewed all 13 waves.

Results

The analyses were designed around two research questions. The first question had to do with recall and forgetting. We examined forgetting as a function of network size, behavioral

specificity, and salience. To do this, we examined recall and forgetting for three relationship domains: 30-day sex partners, 30-day drug use partners, and close friends. The second question had to do with the association between global estimates and named estimates of network size. We were able to examine this question only in the sex partner domain.

Recall and Forgetting

We consider for this analysis only those alters who had been named in a previous wave of the study (that is, we did not consider the wave in which the alter was first named, because we could not detect forgetting until a subsequent wave). Results of these analyses are given in Tables 2 and 3. In these tables, results are weighted. That is, if a given alter was mentioned by a given informant n times (after the first mention), those mentions were weighted by $1/n$ to avoid over-representing stable informants who participate in many interviews and stable relationships described in many interviews. This weighting method effectively gives higher weight to forgetting occurring in short term relationships with only one or two recall/forgetting occurrences.

Recall and forgetting are related to number of partners in Table 2. Results are presented separately according to the number of alters named at a given wave and the type of relationship. There were 166 informants who described 348 sex partners (after the initial mention) at 1,243 occasions over the course of the study (because of weighting, each relationship has equal weight no matter how many times mentioned). During waves where the informant described only one sex partner, the sex partner was forgotten (remembered only after an interviewer reminder) on 1.1% of those occasions (2 out of 803). On the 213 occasions where participants described two sex partners, only five partners were forgotten (4.5%). Forgetting increased as the number of partners enumerated increased. This measure of the percent of described partners who were forgotten is referred to by Wright and Pescosolido (2002) as a “forgetfulness bias coefficient.”

When a person had a single drug use partner, there was a 7.3% chance that the participant would forget them when they were naming drug use partners. If there were four or more drug use partners, there was a 27.7% chance that a partner would be forgotten. When participants were asked to name close friends, there was a small chance of forgetting a single partner (5.9%), but a larger chance of forgetting one out of two or more partners (21.3–29.5%). Thus we see similar patterns of increasing forgetting with increasing numbers of partners. Most statistical tests are deferred until the multivariate analysis below. Forgetting was greater for drug use partners (18.5% overall) than for sex partners (5.9%), and even greater for those who were named as close friends (25.8%). Across the three domains, we found that the rate of forgetting was lowest for spouses, and found increasing forgetting for boy/girlfriends, best friends, other family, and highest for acquaintances. This pattern was observed even for those named with the “close friend” name generator (37% forgetting of “close friends” described later in the interview as friends or acquaintances, and 23% forgetting of non-spouse family members. Using our question about whether sex was “for sex” or “a close emotional connection,” we found 9.2% forgetting (corrected for clustering) when sex was for sex and 3.3% when sex involved an emotional connection ($\chi^2 = 19.6$, $df=1$).

To gain an increased understanding of forgetting, we predicted the log odds of forgetting in a logistic regression analysis adjusted for clustering of waves within relationships (Table 3). Because of the collinearity between connection and relationship duration on the one hand and relationship type on the other, we have conducted two analyses. We omitted connection and relationship duration from Model I and omitted relationship type from Model II. The reference population for this analysis consisted of drug use relationships with spouses. Compared to these reference relationships, results showed that forgetting was significantly less likely in sex relationships (28% as likely) and significantly more likely in close relationships (2.97 times as likely). In Model I, spouses and boyfriends/girlfriends were least likely to be forgotten and

did not differ from one another. Family members, best friends and acquaintances were significantly more likely to be forgotten than spouses and boy/girlfriends, and were significantly different from one another. In Model II, emotional connection and duration of relationship significantly reduced forgetting. In both models, behavioral intensity of interaction reduced forgetting.

We examined several methodological and control variables. Forgetting declined over time (so that it was not a matter of informants learning to avoid naming alters over time to shorten the interview; in fact, this result may suggest that informants initially avoided naming some alters but learned over time that this strategy did not work). Forgetting did not differ significantly for stable informants who participated in more interviews. Women, older, and employed participants were more likely to forget partners than men, younger, or less fully employed participants.

Naming vs. Global Estimates

The second question we addressed was the correspondence between global and naming estimates of network size. Unfortunately, we had the complete data to conduct this detailed analysis only for sex partners.

We compared the global estimate of the total number of sex partners to three estimates based on naming. The “raw named estimate” was the number of spontaneously recalled sex partners. The “corrected named estimate” was the sum of the number of spontaneously recalled and anonymous partners. The “full named estimate” was the sum of the number of recalled, anonymous, and forgotten partners. Table 4 shows how these four estimates of sex network size differed by gender and by drug use. Observations for each informant were weighted by the inverse of the number of waves the informant was interviewed. Men’s and women’s global estimates did not differ significantly, but women named significantly more sex partners than did men. This difference lost significance when anonymous partners were added. Raw named estimates were not reliable proxies for global estimates for men or women ($r = .61$ and $r = .48$).

Table 4 also examines the effect of drug use on estimated number of sex partners. The first and fourth categories are informants who were either always nonusers (of cocaine, heroin, or methamphetamine during the previous 30 days) or always users. The middle two categories involve 78 informants who sometimes reported no drug use and sometimes reported drug use. The “sometimes drug use” column refers to their report of sex partners during the waves when they were currently drug users. All four measures of sex network size differed significantly by drug use. Sex networks were smaller when informants had not been using drugs and larger when they had been using drugs. The raw, spontaneous named estimate is a reliable proxy for the global estimate of sex partners for nonusers ($r = .93$ and $r = .98$), but not for drug users ($r = .40$ and $r = .50$). The large size of sexual networks in these data for drug users are probably related to the predominant use of cocaine, a stimulant, in Houston.

Table 5 shows a further comparison of the estimates of network size. We see that the raw named estimate (spontaneously recalled) differed from the global estimate 12.5% of the time, and when it differed it gave an estimate higher than the global estimate about one fifth of the time and lower four-fifths of the time. The corrected named estimate (recalled + anonymous) differed 8.2% of the time with little imbalance between underestimates and overestimates of the global estimate. We found that the raw estimate omitted on average 0.53 sex partners relative to the global estimate, while the corrected estimate omitted 0.04 partner.

We performed a series of logistic regression analyses (not shown) where the dependent variable was whether there was a discrepancy between the named estimate and the global estimate

(ignoring direction of the discrepancy). In these analyses, waves were clustered within informants. These analyses, using the raw, corrected, and full named estimates each gave the same result: a significant positive effect of the number of sex partners on whether there was a discrepancy (raw, OR = 1.55; corrected, OR = 1.56; full, OR = 1.73; in each analysis, $p < .001$). In addition, gay/lesbian/bisexual informants displayed a greater discrepancy between named and global measures (raw, OR = 2.21; corrected, OR = 2.42; full, OR = 1.95; in each analysis, $p < .03$), as did drug users, but only for the raw estimate (OR = 2.31, $p < .01$). These results are similar to those reported by Brewer and colleagues (Brewer et al 1999; Brewer et al 2005).

Discussion

When trying to measure the size of ego networks, researchers face potential biases in informant self reports, depending on method, population, and domain. The researcher's methodological choices can reduce some of the bias.

Network size

We have seen that the rate of forgetting increases as network size increases. This result is consistent with the effect of network size reported in other studies of network recall (Brewer et al 1999; Brewer & Webster 1999) and of the recall of events (Blair & Burton 1987; Conrad et al 1998). We found that the discrepancy between global and named estimates was both small and relatively balanced for the corrected (spontaneously recalled + anonymous) and full (recalled + anonymous + forgetting) named estimates, while the most frequent and most imbalanced discrepancy was between the raw named estimate and the global estimate. Based on these discrepancies, we infer that informants who give a global estimate of network size do not generally construct a complete internal list of alters except when there are only a few alters. Nevertheless, even though global estimates for larger networks are guesses, they tend to be quite consistent with full detailed estimates that consider named as well as anonymous alters. If concern is to learn about relatively small networks, then measurement of such networks by either enumeration or global estimation should suffer little loss of information as long as there are no anonymous alters. However, attempts to identify larger networks are likely to suffer from higher levels of forgetting. Whether forgetting in large networks is substantively important will depend on one's research goals.

Behavioral specificity

Behavioral specificity tends to reduce forgetting. Both sex partners and drug use partners showed lower amounts of forgetting than alters identified as close friends by a name generator. One reason for this difference in level of forgetting is likely to be the greater behavioral specificity of sex and drug use relationships compared to close relationships. Other investigators have found similar recall benefits for behaviors over emotions (Clair et al 2003; Harvey et al 2004; Kogovsek & Ferligoj 2005). Furthermore, we have established that it is the behavior of drug use that affects forgetting, not some characteristic of persons who use drugs. Differences in forgetting of sex partners between drug users and nonusers were consistent for both global and named estimates of sex partners. Because we had 78 persons who used drugs on some waves and not on others, we were able to determine that the differences in sex partners was because of current drug use and not because of a general factor that differentiated those who ever use drugs.

From the point of view of recall, behavioral relationships have the advantage that the distinction between relationship and non-relationship is relatively clear. Relationships, on the other hand, that are characterized by feelings may be more difficult to distinguish, because closeness depends more on how one feels about a partner than what one has done with the partner.

At the same time that specificity improves recall, certain types of behavioral specificity may be problematic because of social desirability (Catania 1999; Catania et al 1986; Herold & Way 1988). Considerations of social desirability might have been expected to contribute to a greater discrepancy between global and enumerated estimates for gay/lesbian/bisexual informants than for straight informants. Although a trend in this direction was found in these data, the effect was not significant. Nevertheless, special care may need to be taken with this population because of stigma associated with gay sex.

Salience

The third element in recall and forgetting is salience. Our results show that sex partners, highly salient alters, are reported more reliably (with less forgetting) than drug use partners, as others have also found (Brewer 2000; Brewer & Garrett 2001; Brewer et al 1999; Brewer et al 1998). It seems reasonable that this is because sex relationships are more emotionally involving in general than drug use relationships. In support of this notion, we found more forgetting when sex was just “for sex,” compared to when sex was part of a close emotional relationship. Our two statistical models used different variables to account for type of relationship. The results from using descriptive categories of spouse-boy/girlfriend-best friend-acquaintance-family were essentially the same (although with larger level of variance explained) as when we used more parsimonious theoretical concepts of emotional connection, indicating that relationship type was largely a proxy for emotional closeness, as similarly suggested by Marsden and Campbell (1984).

An apparent contradiction to the importance of salience arose in these data. Close friends, a name generator associated with emotional closeness, showed the highest levels of forgetting. In our sample of adults from high drug use neighborhoods, 21% of all alters (and 53% of non-family alters) whom informants named as a “close friend” later in the same interview were classified as “friend,” “acquaintance,” or “drug buddy” (which we classified together as acquaintances). We also found a range of forgetting among those identified as close friends from 3–4% forgetting for spouses and boy/girlfriends compared to 38% forgetting of acquaintances. If all “close friends” involved emotional closeness, there are clearly differences in the emotional quality of these relationships. We suspect that the high level of “forgetting” of family members was not true forgetting but reflected an inference by informants that, in our interviews focusing on HIV risk, we did not “really” want to know about family.

We suspect, although we have no direct data, that the comparability of the global and named estimates in our data depend on both the behavioral specificity of the sex domain as well as its emotional salience. We suspect that as the cognitive and emotional complexity of the measurement domain increase, global estimates will become weaker and weaker approximations of named data (corrected for anonymous alters).

Time frame

The fourth element in the accuracy of informant reports of network size is time frame. Our results show, at least for sex relationships in the past 30 days, that global and named estimates of network size are comparable when anonymous partners are accounted for. The effect of forgetting appears to be particularly serious for global estimates within long time frames, when we compare our results with other studies. Brewer and colleagues (1998), looked at data from studies in two cities, in which informants described partners over 6 months (Colorado Springs), and one year and two years (Seattle). Although Brewer did not subject the differences to statistical analysis, he reported that for a 6-month time frame the named estimate was lower than the global estimate, while for one-year and two-year time frames, the named estimate was higher than the global estimate. In our results with a 30-day time frame, we found that raw named estimates, when inaccurate, were almost four times more likely to be lower than global

estimates. Thus, while informants may tend to miss partners (“guess low,” backward telescoping) in their global estimates when the time frame is long (and recall them as they provide names), it appears that when the time frame is short, informants tend to “guess high” (forward telescoping). Since research has shown that most informants attempt to conduct an implicit, silent enumeration to produce a global estimate (Brewer et al 1998), it appears that this implicit global enumeration is especially sensitive to forgetting of relationships distant in time.

Limitations—There are a number of potential limitations to this study. In the first place, this study takes place in a single city, so we do not know if forgetting is different in Houston from other locations. Second, we have only asked about certain kinds of relationships, so we can’t be sure that we would not have gotten different results from a different set of relationships. Furthermore, we only have global estimates for one of the relationship domains we studied. However, the relationship domains we studied vary in predictable ways along the theoretically meaningful dimensions of network size, behavioral specificity, and salience. Of course, since we do not know how many alters an informant has independently of the informant’s report, we cannot make definitive statements about absolute accuracy. We have, however, been able to make some estimates of the comparative validity of global and naming approaches.

One reason for the apparently low level of forgetting and moderate level of guessing may be the nature of our sample. Much of the literature on the threatening nature of sex partner questions appears to be concerned with largely middle and working class samples with informants with what may be thought of as “conventional” moral standards. At the same time, surveys on sexual behavior by Kinsey (1948;1953) and Laumann (2004;1994) have shown that such data can be collected. Our sample, in contrast, is drawn from low income high drug use neighborhoods and consists of over four-fifths “hard” drug users (cocaine, heroin, methamphetamine). Our drug users and the nonusers who live nearby do not seem to share the hypothesized reluctance to discuss sexual matters described for other populations. Our interviewers are not students, but are professional interviewers comfortable in these neighborhoods with this population. Our study informants not only seem to have little difficulty telling us who their sex partners are, but also describe their sexual behaviors in considerable detail and with considerable reliability (Bell et al 2000;Dowling-Guyer et al 1994;Needle et al 1995).

Another limitation of our method is that we could not identify forgotten partners in short-term relationships. A person had to be a partner at two waves three months apart in order for them to be identified as forgotten. A “one-night stand” or a one-month relationship, if forgotten, could never be detected unless the person became a partner named at a later interview. In this sample, a majority of both sex and drug use partners were reported only once. Not quite all of the one-mention alters were necessarily in short-term relationships, because some of them were described by the small number of study participants who were interviewed only one or two times.

Another potential limitation could result from our research design. One issue that can come up in a longitudinal study, especially a study with a sample from a comparatively recalcitrant population such as drug users, is that study informants have learned that every alter named will require a certain amount of time and effort to answer questions about that alter. As a result, one acknowledges that study informants can perceive an advantage to limiting the number of alters they name in order to shorten their interview. Thus, at least some of the alters who were “forgotten” may have been persons that the informant did not want to spend time talking about in the current interview. However, our procedure for catching such intentionally “forgotten” alters found little evidence for such suppression. Results showed that, contrary to this argument, informant forgetting *declined* rather than increasing over time. Furthermore, the low level of

discrepancy between global estimates (with little overhead related to size) and enumerated estimates also argues against study self-censorship. Obviously, if an informant deliberately and consistently withheld a given alter, the alter would not be detected as forgotten, and this would introduce a bias to the list of named alters. However, our longitudinal data suggest that this may not be a major problem, at least in the population we studied. Informants would have had to remember, from wave to wave over a period of over three years, which alters to “forget.” While such a level of consistency is possible, it is not very plausible among heavy drug users.

Recommendations—The results of the analyses reported here underline the importance of both theory and experience for measurement decisions. There are methodological decisions that one might make in order to improve recall and limit forgetting, but in some cases these decisions might thereby compromise one’s research aims. If one’s theory only requires the size of a network and does not require details about the relationships of which it is composed, then the global estimate seems to be as accurate as a corrected named estimate (including anonymous alters), at least for salient behavioral networks like sex networks. The ease of global estimates in terms of respondent burden also recommends the global method when only overall network size is needed.

At the same time, the amount of information that can be collected from naming methods recommends the naming method if this information is important to the research even though the effort from investigator, interviewer, and especially the informant can be much greater. If much information is collected on each alter, informants may become inattentive or resentful as the interview continues. A compromise between global and naming methods may be possible. Instead of asking in detail about each alter, the investigator could define theoretically meaningful subsets of alters and ask about each subset separately. For example, the investigator could ask for global estimates of the number of “main” sex partners (who might be defined as emotionally committed, intended long-term partners), regular sex partners (frequent but not committed), and casual sex partners (infrequent) and then ask questions about these types of partners rather than about each individually. For this method to work, the theoretically defined subsets will have to be empirically homogeneous so that information provided about each subset will be meaningful.

If the investigator wishes to know both details about relationships and the overall network size, it is clear from our results that using a names list alone will be inadequate for those kinds of networks where names may be unknown or forgotten. In these cases, we recommend asking explicitly about the number of alters whose names are not known. One could ask, as we did, the number of alters “whose names you do not know or can’t remember,” or number of alters the informant “could not recall or did not want to mention” (Brewer et al 1999).

In cases where alter naming is to be used, we recommend that global estimates be taken first and used as a general target for naming. A good strategy is to get a global estimate and then to continue to encourage recall up to that point, and then to continue urging recall to some extent after that point (Brewer et al 2005). This strategy may be particularly important for behaviorally unspecific relationships like friendship. We recommend that in all panel studies of social networks informants be presented with previously described partners to minimize forgetting and also to learn the nature of network turnover. Since the amount of actual forgetting was low for behavioral partners, the expense of longitudinal data collection simply to detect forgetting would seem to be unjustified in most cases for such networks. This procedure, of course, requires that investigators solicit identifiable names, requiring substantial efforts to maintain confidentiality of this information on the part of researchers.

One’s theoretical goals may determine the extent to which one can methodologically improve precision. Short time frames, behavioral specificity, high salience, and small network size may

be useful goals from the point of view of minimizing forgetting, but the feasibility of meeting these goals may depend on experience and theoretical focus. Small networks are feasible if one is studying sexual networks, unless one wishes to study prostitutes or some stimulant drug users (e.g., crack cocaine or methamphetamine) where large sexual networks are common. Behavioral specificity will improve recall, and if one wishes to study sexually transmitted diseases, the transmission of salient and identifiable pieces of information, or enacted social support, one can probably count on relatively good recall. However, if one's theoretical interest is in stress, environmental diseases, potential power, or potential social support, then behavioral data may be unavailable or of low validity.

The time frame of network measures can be important. We found little discrepancy between global and named estimates (controlling for anonymous partners) for a 30-day time frame. Other researchers have found much greater discrepancies over larger time frames. However, time frame may be dictated by the investigator's research question. In our research, we attempted to maximize both theoretical and measurement precision by using a 3-month repeated longitudinal design with a 30-day measurement window. We felt that the 30-day window would lower respondent burden for persons with changing networks and that we would recoup the loss of information with repeated longitudinal measurements. Of course, our method cost a few million dollars over the length of the study. On a more limited budget, we would have made different decisions.

The integral association of theory and measurement is clear in domains like friendship. If one is interested in a construct like friendship, one might find that it is difficult to specify a comprehensive set of relevant behaviors (McCallister & Fischer 1983). One may choose to use an imprecise name generator like "Name your friends" instead. On the other hand, much of the imprecision in the concept of friendship lies in notions of friendship that differ among informants (Fischer 1982). We recommend against a name generator like "close friend" (at least in adult samples like ours) unless additional information is collected on type of relationship in order to interpret the meaning of the name generator to the informant. The uninterpreted name generator seems to have such a broad range of meanings in our adult sample as to be potentially theoretically meaningless. Additional research is needed to determine whether our experience with the "close friend" name generator is a general phenomenon, whether it is characteristic of low income or drug-using populations, or whether it is a local phenomenon in Houston. Even when explicitly defined, a concept like friendship may not have an empirical metric. However, if one can formulate the concept of friendship in a theoretically precise way that can be communicated to the informant (e.g., whether friendship refers to short-term mood, medium-term feelings, or long-term persistence), then the imprecision of the informant's threshold level for the concept can be reduced.

This discussion has considered some of the ways that theory and measurement work together in successful research. In many cases, it will be true that theoretical goals will trump measurement goals. It will often require creative invention of new measures and new measurement techniques to narrow the theory-measurement gap. Furthermore, imprecision can never be eliminated from measurement, so it is important to develop new techniques to measure the level of precision. For example, our longitudinal design and computer-assisted access to prior interviews allowed us to measure the level of forgetting in three domains of relationships. These estimates can thus be used in statistical analyses to adjust for forgetting using methods that are beginning to be developed (Handcock & Jones 2004). Thus, investigators can have in their research arsenal techniques for reducing measurement imprecision such as forgetting and adjusting for the unavoidable imprecision that remains.

References

- Adams, J.; Moody, J. "To tell the truth": Measuring concordance in multiply reported network data. Presented at Sunbelt Social Networks Conference; Redondo Beach, CA. 2005.
- Alexander CN, Campbell EQ. Peer influences on adolescent educational aspirations and attainments. *American sociological review* 1964;29:568–575.
- Anderson, JR. *The adaptive character of thought*. Hillsdale, NJ: Erlbaum; 1990.
- Bachman JG, O'Malley PM. When four months equal a year: Inconsistencies in student reports of drug use. *Public opinion quarterly* 1981;45:536–548.
- Bell DC, Atkinson JS, Mosier V. The role of gatekeepers in limiting HIV transmission. *International journal of sociology and social policy* 2002;22:47–76.
- Bell DC, Montoya ID, Atkinson JS. Partner concordance in reports of joint risk behaviors. *JAIDS: Journal of acquired immune deficiency syndromes* 2000;25:173–181.
- Bernard HR, Killworth PD. Informant accuracy in social network data. II. Human communication research. 1977;4:3–18.
- Bernard HR, Killworth PD, Kronenfeld D, Sailer L. The problem of informant accuracy: The validity of retrospective data. *Annual review of anthropology* 1984;1:495–517.
- Bernard HR, Killworth PD, Sailer L. Summary of research on informant accuracy in network data, and on the reverse small world problem. *Connections* 1981;4:11–25.
- Bernard HR, Killworth PD, Sailer L. Informant accuracy in social-network data: V. An experimental attempt to predict actual communication from recall data. *Social science research* 1982;11:30–66.
- Blair E, Burton S. Cognitive processes used by survey respondents to answer behavioral frequency questions. *Journal of consumer research* 1987;14:280–288.
- Blair EA, Ganesh GK. Characteristics of interval-based estimates of autobiographical frequencies. *Applied cognitive psychology* 1991;5:237–250.
- Bradbourn, N. Response effects. In: Rossi, P.; Wright, J.; Anderson, A., editors. *Handbook of survey research*. Academic Press; New York: 1983. p. 289–328.
- Brewer DD. Patterns in the recall of persons in a student community. *Social networks* 1993;15:335–359.
- Brewer DD. Forgetting in the recall-based elicitation of personal and social networks. *Social networks* 2000;22:29–43.
- Brewer DD, Garrett SB. Evaluation of interviewing techniques to enhance recall of sexual and drug injection partners. *Sexually transmitted diseases* 2001;28:666–677. [PubMed: 11677390]
- Brewer DD, Garrett SB, Kulasingam S. Forgetting as a cause of incomplete reporting of sexual and drug injection partners. *Sexually transmitted diseases* 1999;26:166–176. [PubMed: 10100775]
- Brewer, DD.; Potterat, JJ.; Garrett, SB.; Muth, SQ.; Roberts, JM., Jr; Rothenberg, RB. Comparison of direct estimate and partner elicitation methods for measuring the number of sexual and injection partners. Presented at American Public Health Association; Washington, DC. 1998.
- Brewer DD, Potterat JJ, Muth SQ, Malone PZ, Montoya P, et al. Randomized trial of supplementary interviewing techniques to enhance recall of sexual partners in contact interviews. *Sexually transmitted diseases* 2005;32:189–193. [PubMed: 15729158]
- Brewer DD, Webster CM. Forgetting of friends and its effects on measuring friendship networks. *Social networks* 1999;21:361–373.
- Brown, NR.; Sinclair, RC. Estimating the number of lifetime sexual partners: Men and women do it differently. Presented at American Association for Public Opinion Research; Norfolk, VA. 1997.
- Burt RS. A note on the General Social Survey's ersatz network density item. *Social networks* 1987;9:75–85.
- Burt RS, Ronchi D. Measuring a large network quickly. *Social networks* 1994;16:91–135.
- Burton S, Blair E. Task conditions, response formulation processes, and response accuracy for behavioral frequency questions in surveys. *Public opinion quarterly* 1991;55:50–79.
- Campbell KE, Lee BA. Name generators in surveys of personal networks. *Social networks* 1991;13:203–221.
- Catania JA. A framework for conceptualizing reporting bias and its antecedents in interviews assessing human sexuality. *Journal of sex research* 1999;36:25–38.

- Catania JA, Gibson DR, Marin B, Coates TJ, Greenblatt RM. Response bias in assessing sexual behaviors relevant to HIV transmission. *Evaluation and program planning* 1993;13:52–72.
- Catania JA, McDermott LJ, Pollack LM. Questionnaire response bias and face-to-face interview sample bias in sexuality research. *Journal of sex research* 1986;22:52–72.
- Chang L, Krosnick JA. Measuring the frequency of regular behaviors: Comparing the “typical week” to the “past week”. *Sociological methodology* 2003;33:55–80.
- Clair S, Schensul JJ, Raju M, Stanek E, Pino R. Will you remember me in the morning? Test-retest reliability of social network analysis examining HIV-related risky behavior in urban adolescents and young adults. *Connections* 2003;25:88–97.
- Conrad FG, Brown NR, Cashman ER. Strategies for estimating behavioral frequency in survey interviews. *Memory* 1998;6:339–366. [PubMed: 9829096]
- Couper MP, Singer E, Tourangeau R. Understanding the effects of audio-CASI on self-reports of sensitive behavior. *Public opinion quarterly* 2003;67:385–395.
- Des Jarlais DC, Paone D, Milliken J, Turner CF, Miller H, et al. Audio-computer interviewing to measure risk behaviour for HIV among injecting drug users: A quasi-randomized trial. *Lancet* 1999;353:1657–1661. [PubMed: 10335785]
- Dowling-Guyer S, Johnson ME, Fisher DG, Needle R, Watters J, et al. Reliability of drug users’ self-reported HIV risk behaviors and validity of self-reported recent drug use. *Assessment* 1994;1:383–392.
- Ernhart CB, Morrow-Tlucak M, Sokol RJ, Martier S. Underreporting of alcohol use in pregnancy. *Alcoholism, clinical and experimental research* 1988;12:506–511.
- Feld SL, Carter WC. Detecting measurement bias in respondent reports of personal networks. *Social networks* 2002;24:365–383.
- Fendrich M, Xu Y. The validity of drug use reports from juvenile arrestees. *International journal of the addictions* 1994;29(8):971–985. [PubMed: 7960302]
- Fischer CS. What do we mean by ‘friend’? An inductive study. *Social networks* 1982;3:287–306.
- Freeman LC, Romney AK. Words, deeds and social structure: A preliminary study of the reliability of informants. *Human organization* 1987;46:330–334.
- Freeman LC, Romney AK, Freeman SC. Cognitive structure and informant accuracy. *American anthropologist* 1987;89:310–325.
- Graham CA, Catania JA, Brand R, Tuong T, Canchola JA. Recalling sexual behavior: A methodological analysis of memory recall bias via interview using the diary as the gold standard. *Journal of sex research* 2003;40:325–332. [PubMed: 14735406]
- Hammer M. Explorations into the meaning of social network interview data. *Social networks* 1984;6:341–371.
- Handcock MS, Jones JH. Likelihood-based inference for stochastic models of sexual network formation. *Theoretical Population Biology* 2004;65:413–422. [PubMed: 15136015]
- Harrison LD. The validity of self-reported data on drug use. *Journal of drug issues* 1995;25:91–111.
- Harvey SM, Bird ST, Henderson JT, Beckman LJ, Huszti HC. He said, she said: Concordance between sexual partners. *Sexually transmitted diseases* 2004;31:185–191. [PubMed: 15076933]
- Herold E, Way L. Sexual self-disclosure among university women. *Journal of sex research* 1988;24:1–14.
- Huttenlocher J, Hedges LV, Bradburn NM. Reports of elapsed time: Bounding and rounding processes in estimation. *Journal of experimental psychology: Learning, memory, and cognition* 1990;16:196–213.
- Kashy DA, Kenny DA. Do you know whom you were with a week ago Friday? A re-analysis of the Bernard, Killworth, and Sailer studies. *Social psychology quarterly* 1990;53:55–61.
- Kauth MR, St Lawrence JS, Kelly JA. Reliability of retrospective assessments of sexual HIV risk behavior: A comparison of biweekly, three-month, and twelve-month self-reports. *AIDS education and prevention* 1991;3:207–214. [PubMed: 1834142]
- Killworth PD, Bernard HR. Informant accuracy in social network data. *Human organization* 1976;35:269–286.
- Kinsey, AC.; Pomeroy, WB.; Martin, CE. *Sexual behavior in the human male*. Philadelphia: Saunders; 1948.

- Kinsey, AC.; Pomeroy, WB.; Martin, CE.; Bebbard, PH. Sexual behavior in the human female. Philadelphia: Saunders; 1953.
- Kirke DM. Collecting peer data and delineating peer networks in a complete network. *Social networks* 1996;18:333–346.
- Kissinger P, Rice J, Farley T, Trim S, Jewitt K, et al. Application of computer-assisted interviews to sexual behavior research. *American journal of epidemiology* 1999;149:950–954. [PubMed: 10342804]
- Knoke, D.; Kuklinski, JH. Network analysis. Newbury Park, CA: Sage; 1982.
- Kogovsek T, Ferligoj A. Effects on reliability and validity of egocentered network measurements. *Social networks* 2005;27:205–229.
- Kogure T, Hatta T, Kawakami A, Kawaguchi J, Makino T. Characteristics of proper names and temporal memory of social news events. *Memory* 2001;9:103–116. [PubMed: 11338936]
- Krosnick JA. Response strategies for coping with the cognitive demands of attitude measures in surveys. *Applied cognitive psychology* 1991;5:213–236.
- Kurth AE, Martin DP, Golden MR, Weiss NS, Heagerty PJ, et al. A comparison between audio computer-assisted self-interviews and clinician interviews for obtaining the sexual history. *Sexually transmitted diseases* 2004;31:719–726. [PubMed: 15608586]
- Laumann EO. Friends of urban men: An assessment of accuracy in reporting their socioeconomic attributes, mutual choice, and attitude agreement. *Sociometry* 1969;32:54–69.
- Laumann, EO.; Ellingson, S.; Mahay, J.; Paik, A.; Youm, Y. The sexual organization of the city. Chicago: University of Chicago Press; 2004.
- Laumann, EO.; Gagnon, JH.; Michael, RT.; Michaels, R. The social organization of sexuality: Sexual practices in the United States. Chicago: University of Chicago Press; 1994.
- Marsden PV. Network data and measurement. *Annual review of sociology* 1990;16:435–463.
- Marsden, PV. Recent developments in network measurement. In: Carrington, PJ.; Scott, J.; Wasserman, S., editors. *Models and methods in social network analysis*. Cambridge University Press; Cambridge: 2005. p. 8-30.
- Marsden PV, Campbell KE. Measuring tie strength. *Social forces* 1984;63:482–501.
- McCallister, L.; Fischer, CS. A procedure for surveying personal networks. In: Burt, RS.; Minor, MJ., editors. *Applied network analysis: A methodological introduction*. Sage; Beverly Hills, CA: 1983. p. 75-88.
- McCarty C, Killworth PD, Bernard HR, Johnsen EC, Shelley GA. Comparing two methods for estimating network size. *Human organization* 2001;60:28–39.
- Menon G. The effects of accessibility of information in memory on judgments of behavioral frequencies. *Journal of consumer research* 1993;20:431–440.
- Mieczkowski T, Barzelay D, Gropper B, Wish E. Concordance of three measures of cocaine use in an arrestee population: Hair, urine and self-report. *Journal of psychoactive drugs* 1991;23:241–249. [PubMed: 1774609]
- Milardo RM. Theoretical and methodological issues in the identification of social networks of spouses. *Journal of marriage and the family* 1989;51:165–174.
- Milardo RM. Comparative methods for delineating social networks. *Journal of social and personal relationships* 1992;9:447–461.
- Montoya ID, Patek JA, Covarrubias LD, Graves JA. Posttraumatic stress disorder among Hispanic and African-American drug users. *American journal of drug & alcohol abuse* 2003;29:729–741. [PubMed: 14713136]
- Morgan DL, Neal MB, Carder P. The stability of core and peripheral networks over time. *Social networks* 1997;19:9–25.
- Needle R, Fisher DG, Weatherby N, Chitwood D, Brown B, et al. The reliability of self-reported HIV risk behaviors of drug users. *Psychology of addictive behaviors* 1995;9:242–250.
- Perils TE, Des Jarlais DC, Friedman SR, Arasteh K, Turner CF. Audio-computerized self-interviewing versus face-to-face interviewing for research data collection at drug abuse treatment programs. *Addiction* 2004;99:885–896. [PubMed: 15200584]

- Prohaska V, Brown NR, Belli RF. Forward telescoping: The question matters. *Memory* 1998;6:455–465. [PubMed: 9829101]
- Schräpler JP. Response behavior in panel studies: A case study for income nonresponse by means of the German socio-economic panel (SOEP). *Sociological methods and research* 2004;33:118–156.
- Shulman N. Network analysis: A new addition to an old bag of tricks. *Acta sociologica* 1976;19:307–323.
- Stone VE, Catania JA, Binson D. Measuring change in sexual behavior: Concordance between survey measures. *The journal of sex research* 1999;36:102–108.
- Sudman S. Experiments in the measurement of the size of social networks. *Social networks* 1985;7:127–151.
- Sudman S. Experiments in measuring neighbor and relative social networks. *Social networks* 1988;10:93–108.
- Sudman, S.; Bradburn, NM. Asking questions: A practical guide to questionnaire design. Chicago: Aldine; 1982.
- Sudman, S.; Norman, M.; Bradman Schwarz, N. Thinking about answers: The application of cognitive processes to survey methodology. San Francisco: Jossey-Bass Publishers; 1996.
- Tourangeau, R.; Rips, LJ.; Rasinski, K. The psychology of survey response. New York: Cambridge University Press; 2000.
- Tourangeau R, Smith TW. Asking sensitive questions: The impact of data collection mode, question format, and question context. *Public opinion quarterly* 1998;60:275–304.
- van der Poel MGM. Delineating personal support networks. *Social networks* 1993;15:49–70.
- Wasserman, S.; Faust, K. Social network analysis: Methods and applications. Cambridge: Cambridge University Press; 1994.
- White K, Watkins SC. Accuracy, stability and reciprocity in informal conversational networks in rural Kenya. *Social networks* 2000;22:337–355.
- Wixted JT, Ebbesen EB. On the form of forgetting. *Psychological science* 1991;2:409–415.
- Wright ER, Pescosolido BA. Sorry, I forgot”: The role of recall error in longitudinal personal network studies. *Social networks and health* 2002;8:113–129.

Table 1

Sample Characteristics

	Frequency (%) (N = 202)
Gender	
Male	113 (55.9%)
Female	89 (44.1)
Race	
African American	111 (55.0)
Anglo	33 (16.3)
Hispanic	57 (28.2)
Asian	1 (0.5)
Age at baseline	
20 – 30	24 (11.9)
31 – 40	67 (33.2)
41 – 50	77 (38.1)
51 and over	34 (16.8)
Employment, Previous 6 Months (baseline)	
None	98 (48.5)
Odd jobs	24 (11.9)
Part-time	20 (9.9)
Full-time	60 (29.7)
Sexual Behavior (full panel)	
Only straight (heterosexual)	144 (71.3)
Ever gay/lesbian/bisexual	47 (23.3)
Not sexually active	11 (5.4)
Drug Use Last 30 Days (Multiple Responses)	
None	33 (16.3)
Alcohol	147 (72.8)
Marijuana	95 (47.0)
Crack cocaine	112 (55.4)
Powder cocaine	49 (24.3)
Heroin	30 (14.8)
Methamphetamine	16 (7.9)

Table 2

Forgetting by Domain

	Domain		
	Sex partner	Drug use Partner	Close friend
Number of partners			
One partner	1.1%	7.3%	5.9%
Two partners	4.5	9.6	21.3
Three partners	5.5	15.6	29.5
Four or more partners	20.1	27.7	26.1
Relationship type			
Spouse	0.3%	0.5%	3.4%
Other family	---	19.4	23.1
Boyfriend/girlfriend	2.7	3.7	4.4
Best friend	3.1	8.1	18.0
Acquaintance/friend	11.0	24.2	37.8
Meaning of sex			
Sex for sex	9.2%		
Sex for connection	3.3		
All partners	5.9	18.5	25.8
Number of informants	166	135	192
Number of partners	348	569	2,483
Number of observations	1,243	1,376	8,582

Note: Each relationship is weighted by $1/\underline{n}$, where $\underline{n}$ is the number of waves over which the relationship was observed.

Table 3

Predictors of Forgetting

	Model I		Model II	
	Odds Ratio	p	Odds Ratio	p
<u>Relationship domain</u> (relative to drug use)				
Sex relationship	0.28	.001	0.23	.001
Close relationship	2.97	.001	3.11	.001
<u>Relationship type</u> (relative to spouse)				
Family member	4.67	.001		
Boy/girlfriend	1.13	.788		
Best friend	3.04	.001		
Acquaintance	7.05	.001		
<u>Relationship quality</u>				
Connection			0.96	.003
Relationship duration			0.99	.050
Interaction frequency	0.97	.001	0.96	.001
Gay/lesbian/bisexual	1.14	.075	1.14	.075
<u>Control variables</u>				
Wave number	0.95	.001	0.95	.001
Number of waves	1.01	.475	0.98	.059
Female	1.23	.001	1.22	.002
Age	1.02	.001	1.02	.001
Employment status	1.26	.024	1.27	.021
Black	1.05	.640	1.06	.569
Hispanic	1.19	.092	1.21	.067
Pseudo R ²	.096		.081	
Wald chi-square	476.3 (df=15)		477.4 (df=13)	
p <	.001		.001	

Log odds describe the relative probability of forgetting.

N = 15,974 observations; Standard errors adjusted for 4346 clusters (alters)

Table 4

Estimated numbers of 30-day Sex Partners

	Men (n = 106)	Women (n = 85)	Always nonusers (n = 28)	Sometimes nonusers (n = 78)	Sometimes drug users (n = 78)	Always drug users (n = 85)
Number of sex partners						
Global estimate	1.44	1.97	0.90	0.84	1.66	2.26**
Raw named estimate	1.01	1.31*	0.84	0.80	1.10	1.40**
Corrected named estimate	1.43	1.90	0.85	0.82	1.67	2.18**
Full named estimate	1.44	1.91	0.85	0.83	1.70	2.20**
Correlation with global						
Raw named estimate	0.61	0.48	0.93	0.98	0.40	0.50
Corrected named estimate	0.98	0.99	0.94	0.99	0.98	0.99
Full named estimate	0.98	0.99	0.94	0.99	0.99	0.99

* p < .05
** p < .01

Table 5

Discrepancy Between Global and Named Estimates of 30-day Sex Partners

	Raw		Corrected		Full	
	Discrepant	Discrepancy	Discrepant	Discrepancy	Discrepant	Discrepancy
Named estimate higher	2.6%	-1.06	3.2%	-1.13	3.7%	-1.18
Named and global estimates equal	87.5	0	91.8	0	91.7	0
Global estimate higher	10.0	5.62	5.0	1.60	4.6	1.61
All observations	12.5	0.53	8.2	0.04	8.3	0.03

Note: Raw = raw named estimate; Corrected = corrected named estimate (raw + anonymous partners); Full = full named estimate (corrected + forgotten partners). Totals do not always add to 100% due to rounding