Published in final edited form as:

Tetrahedron. 2007 July 2; 63(27): 6317–6321.

Enantioselective Transaminations by Dendrimeric Enzyme Mimics

Ronald Breslow, Sujun Wei, and Craig Kenesky

Department of Chemistry, Columbia University, New York NY 10027

Abstract

PAMAM dendrimers have been constructed with a pyridoxamine core, and chiral capping amino groups. Transamination to form phenylalanine and alanine from their related ketoacids produced enantioselectivities induced by the formally remote chiral caps, supporting computer models that indicate folding of the dendrimer chains back into the core region.

1. Introduction

We have reported mimics of transamination,¹ both simple pyridoxamine derivatives²⁻⁸ and enzyme mimics with a pyridoxamine attached to cyclodextrins,⁹⁻¹³ to a synthetic macrocycle,¹⁴ to polyethylenimines of various sizes,¹⁵⁻¹⁷ and to dendrimers.¹⁸ The process led to the formation of amino acids from ketoacids, while converting the pyridoxamine to its pyridoxal form (Scheme 1). Turnover was achieved by a process of decarboxylative transamination to regenerate the pyridoxamine from the pyridoxal.^{19,20} With a modified polyethyleneimine, having a hydrophobic core, we were able to use the pyridoxamine as a reversibly bound cofactor rather than a covalently attached functional group, and achieved accelerations of transamination as large as 725,000-fold relative to pyridoxamine itself.¹⁹

Transamination produces an amino acid with a new chiral center, so synthesis with enantioselectivity is an important goal. We were able to achieve considerable enantioselectivity with a rigid pyridoxamine derivative carrying a chirally mounted amino group for proton transfers.² We also saw enantioselective transamination with a pyridoxamine covalently attached to a chiral triamine that was derived by reduction of a tripeptide (but not with the unreduced tripeptide).²¹ We then wished to create a polyamine catalyst that could produce enantioselective transamination by the simple action of the chirality of the polymer itself, as happens in natural enzymes.

We have described elsewhere the enantioselective transaminations by pyridoxamines attached to a linear polyamine derived by the polymerization of a chirally substituted oxazoline.²² Further work on this system is underway. However, dendrimers are three-dimensional macromolecules with defined structures—in contrast with polyimines—some as large as small proteins, so they have good potential as enzyme mimics.

In our previous study, we saw that a PAMAM dendrimer grown from a pyridoxamine core performed transaminations when basic amino groups were linked to the outside of the dendrimer, formally far from the pyridoxamine unit.¹⁸ Without those remote amino groups

Correspondence to: Ronald Breslow.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

little catalysis was seen. This reflects the fact that dendrimers can be flexible species in which the chains fold back into the otherwise empty space near the core, as our computer model of the pyridoxamine-dendrimer species indicated.²³ The tertiary amino groups that are part of the core of a PAMAM dendrimer are not effective catalysts, presumably because they cannot reach the core pyridoxamine. Thus it seemed likely that we would see chiral induction in transamination with a pyridoxaminedendrimer species carrying chirally-mounted amino groups on their periphery, as terminators of the dendrimer chains. This is what we have now observed.

2. Results and discussion

2.1. Synthesis of the dendrimers

The synthesis of our dendrimers started from protected pyridoxamine with an attached 2-aminoethanethiol spacer (**1**). The NH₂ group on the spacer was used as a starting point to synthesize PAMAM dendrimers. Following the typical two-step iterative sequences²⁴—attachment of two molecules of methyl acrylate to each NH₂ group, then aminolysis of each ester group with excess ethylenediamine—generations G-1 to G-4 were synthesized (scheme 2).

The amino groups on the periphery of each generation were modified with N,N-dimethyl-L-phenylalanine using water soluble 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)²⁵ as a coupling reagent. The extent of coverage of the periphery was indicated by comparing the NMR integrated areas of the phenyl protons to those of Boc groups in the pyridoxamine core, and further confirmed by mass spectra. The series of dendrimers were referred as G-n-Me₂ (n refers to generation). We saw that up to generation 3 the materials were reasonably pure, but at generation 4 the mass spectrum indicated a range of masses, reflecting incomplete attachment of all sixteen dimethylphenylalanine groups. Finally, the Boc and the MOM groups on the pyridoxamine were removed with 1N HCl. An example for the first generation is shown as structure **2**.

2.2 Transamination studies

The transamination reactions transforming α -keto acids to amino acids were performed as indicated in the legend of Table 1 (see also Experimental), and monitored by HPLC. The % ee values are summarized in Table 1.

To further understand the role of the chiral unit on the periphery, N-acetyl-L-phenylalanine was attached to G-2 (we refer it as G-2-Ac). After the removal of Boc and MOM groups in pyridoxamine, this molecule also transaminated phenylpyruvic acid to phenylalanine, but the product was racemic. Therefore, the chiral units on the periphery did not simply create a chiral environment; the basic dimethylamino group is also necessary. This observation is consistent with our previous findings that peripheral dimethylamino groups on a PAMAM dendrimer with a pyridoxamine core reach the core and catalyze transaminations, but N-acetyl terminated dendrimers do not significantly catalyze the transaminations.

We also examined dendrimers capped with phenylalanine and with N-methylphenylalanine instead of the N,N-dimethyl derivative. Interestingly, these performed transaminations but the products were racemic. Probably the conformations differ with these more hydrophilic caps, bringing the chiral centers away from the pyridoxamine. Reactions of the primary or secondary amino groups of these derivatives with the ketoacid substrates are also possible.

2.3 Molecular mechanics calculations

To gain a better understanding of the stereoselectivity of the transaminations taking place within the chiral environment of the dendrimers, we performed molecular mechanics calculations (MacroModel 9.1) of the pyridoxamine dendrimer complexes. Calculations were executed using the MMFFs (Merck Molecular Force Field) force field and a water GB/SA (Generalized Born Surface Area) solvent continuum to promote the hydrophobic packing of the dendrimer arms, as is expected in the real aqueous system. The resulting structure of the G3 pyridoxamine dendrimer is shown in Figure 1 in two different forms. In the tube version it is clear that the pyridoxamine unit is solvent exposed, and that some of the formally remote phenyl rings are in fact packed close to the pyridoxamine. This explains their ability to induce enantioselectivity in the products. In the space-filling structures with van der Waals radii added it is clear that the computed structures are solidly packed. The structure on the left has the same view as that of the tube version, while the one on the right has this rotated by 90° to show how solid the packing is. Thus the dendrimer is hardly “tree-like” unless the branches and trunk of the tree are as flexible as spaghetti. The computed structures of the other dendrimers are shown in Supplementary data.

The G1 and G2 complexes had significant conformational flexibility, with no well-defined three-dimensional structure. However, both pyridoxamine complexes possessed intramolecular hydrogen bonds between the pyridine nitrogen and the amide NH groups. Such

a hydrogen bond holds the pyridoxamine residue in the proximity of a chiral terminal group and promotes asymmetry in the transamination.

The G3 dendrimer differs from the smaller dendrimers in the existence of a well-defined structure stabilized by the hydrophobic packing of dendrimer appendages, and intramolecular hydrogen bonds between amide groups. Unlike the G1 and G2 structures, with dendrimer arms radiating outward from the pyridoxamine core, the outermost arms of the G3 dendrimer fold backwards into the interior, presumably due to the polar solvation medium. In this structure, the pyridoxamine residue exists on the surface of the dendrimer, reflecting the length of the thioethoxy linker and the backwards folding of the dendrimer arms. This arrangement of the pyridoxamine residue in proximity to the chiral terminal groups explains why the G3 compound shows the best enantioselectivity in these transaminations.

Knowing that the pyridoxamine residue rested on the surface of the G3 dendrimer, we reasoned that elaboration of the system to a fourth generation could encapsulate the catalytic core residue fully within the chiral hydrophobic environment. Upon simulation of the G4 dendrimer, we were surprised to find that the pyridoxamine residue did not reside within the interior of the complex. Instead, aggregation of the hydrophobic groups to form an interlocking network of dendrimer arms actually withdrew the chiral termini somewhat from the pyridoxamine, which, being the most hydrophilic region of the molecule, had a significant degree of aqueous solvation. This observation may account for the slight loss of enantioselectivity obtained with G4 relative to G3. Of course our mass spectrum indicates that the G4 system does not have its full complement of phenylalanine units, in contrast to G3, which may contribute to the experimental finding of slight loss in enantioselectivity.

3. Conclusions

Our results, in combination with our previous findings, are strong experimental evidence to support the dense core model of dendrimer structures in solution, where the flexible chains can fold back to fill otherwise empty space, as indicated in our computer models. We observe catalysis and chiral induction by groups that are formally far from the pyridoxamine core, in topological terms, but are clearly nearby in terms of their real geometry.

Furthermore, our dendrimeric enzyme mimics have achieved significant enantiomeric excesses in the transamination products, induced by this formally remote chirality. Coupled with our previous demonstration that decarboxylative transaminations can be used to promote turnovers in such models,^{19,20} the dendrimeric mimics are a real advance in the quest to achieve enzyme mimics with potency and utility.

4. Experimental

4.1 General

Solvents, inorganic salts and organic reagents were purchased from commercial resources and used without further purification unless otherwise noted. Merck pre-coated 0.25mm silica plates coating a 254 nm fluorescence indicator were used for analytical thin-layer chromatography. Flash chromatography was performed on 230-400 mesh silica (Silica Gel 60) from EM Science. Analytical HPLC was run on a Waters 600 liquid chromatography with a pumping system, a temperature controlled autosampler (Waters 717), and a diode-array UV-vis detector (Waters 2996). Waters Sunfire C18 reverse phase analytical columns (particle size 5 μ m, 4.6 $\times$ 150 mm) were used as the solid phase. NMR spectra were obtained on a Bruker DPX 300 MHz spectrometer. UV-vis spectra were taken on a Varian Cary IE UV-vis spectrometer. CIMS spectra were taken on Nermag R-10-10 instrument. FAB MS spectra were taken on a JEOL JMS-DX-303 HF instrument using either glycerol or p-nitrobenzyl alcohol

as matrices. Matrix assisted laser desorption ionization (MALDI) mass spectra were taken on a VOYAGER DE instrument with α -cyano-4-hydroxycinnamic acid as a matrix. Positive ion mass spectra were acquired in the linear mode.

4.2 Synthesis of the dendrimers

The dendrimers were grown from a protected pyridoxamine using methyl acrylate and ethylenediamine, as described in references 18 and 23. They were terminated with ethylenediamine, and the terminal amino groups were then acylated with phenylalanine derivatives and the protecting groups were then removed as described below.

G-1: ^1H NMR (300MHz, D_2O): 8.15, s, 1H; 5.09, s, 2H; 4.45, s, 2H; 3.89, s, 2H; 3.62, s, 3H; 3.29-3.19, m, 4H; 2.79-2.66, m, 8H; 2.66-2.52, m, 4H; 2.47, s, 3H; 2.49-2.29, m, 4H; 1.41, s, 9H. CI MS: 600.51(M+1).

G-2: ^1H NMR (300MHz, D_2O): 8.15, s, 1H; 5.08, s, 2H; 4.44, s, 2H; 3.89, s, 2H; 3.62, s, 3H; 3.31-3.17, m, 12H; 2.80-2.53, m, 28H; 2.50-2.26, m, 15H; 1.41, s, 9H. FAB+ MS: 1056.4(M+1).

G-3: ^1H NMR (300MHz, D_2O): 8.20, s, 1H; 5.12, s, 2H; 4.48, s, 2H; 3.93, s, 2H; 3.66, s, 3H; 3.32-3.21, m, 28H; 2.95-2.58, m, 60H; 2.54-2.35, m, 31H; 1.45, s, 9H. MALDI MS $+$: 1970.49.

G-4: ^1H NMR (300MHz, D_2O): 8.21, s, 1H; 5.13, s, 2H; 4.49, s, 2H; 3.94, s, 2H; 3.67, s, 3H; 3.32-3.16, m, 60H; 2.95-2.57, m, 124H; 2.54-2.37, m, 63H; 1.46, s, 9H. MALDI MS $+$: 3792.76.

G-1Me₂: All modified dendrimers were prepared analogously to the following procedure: G-1(0.1017g, 0.17mmol) and N,N-dimethyl-L-phenylalanine (0.0721g, 0.374mmol) were dissolved in 2ml CH_3OH /2ml CH_3CN . After 5 minutes, 4-(4,6-dimethoxy-1, 3, 5-triazin-2-yl)-4-methylmorpholinium chloride (0.1032g, 0.374mmol) was added into the solution. The mixture was stirred overnight. After removal of the solvent, the residue was dissolved in CH_2Cl_2 , which was washed successively with saturated aqueous Na_2CO_3 .

solution and brine. Removal of the solvent gave the product as a slightly yellow solid. Yield: 50%. ^1H NMR (300MHz, CD_3OD): 8.13, s, 1H; 7.30-7.15, m, 10H; 5.03, s, 2H; 4.46, s, 2H; 3.92, s, 2H; 3.58, s, 3H; 3.57-3.51, dd, 2H; 3.25-2.94, m, 12H; 2.90-2.73, m, 6H; 2.72-2.54, m, 14H; 2.46, s, 3H; 2.43-2.28, m, 4H; 1.41, s, 9H. CI MS: 982(M+33).

G-2Me₂: Yield: 42%. ^1H NMR (300MHz, CD_3OD): 8.13, s, 1H; 7.37-7.08, m, 20H; 5.03, s, 2H; 4.45, s, 2H; 3.89, s, 2H; 3.54, s, 3H; 3.25-2.86, m, 32H; 2.84-2.61, m, 16H; 2.60-2.46, m, 7H; 2.45-2.20, m, 36H; 1.42, s, 9H. FAB+ MS: 1758.06(M+1), 1780.09(M+Na).

G-3Me₂: Yield: 37%. ^1H NMR (300MHz, CD_3OD): 8.13, s, 1H; 7.37-7.08, m, 40H; 5.03, s, 2H; 4.45, s, 2H; 3.91, s, 2H; 3.59, s, 3H; 3.32-2.85, m, 70H; 2.84-2.61, m, 34H; 2.60-2.46, m, 11H; 2.45-2.18, m, 76H; 1.42, s, 9H. MALDI MS+: 3372.54(M+2).

G-4Me₂: Yield: 30%. ^1H NMR (300MHz, CD_3OD): 8.15, s, 1H; 7.32-7.10, m, 80H; 5.05, s, 2H; 4.47, s, 2H; 3.91, s, 2H; 3.59, s, 3H; 3.32-2.84, m, 144H; 2.83-2.61, m, 72H; 2.60-2.46, m, 19H; 2.45-2.16, m, 156H; 1.44, s, 9H. MALDI MS+: 5000-7000. Expected Mass: 6885.6

4.3 Deprotection of the dendrimers

All modified dendrimers were deprotected similarly to the following procedure: 50mg of G-1Me₂ was dissolved in 1ml of HCl(1N). The reaction was stirred at room temperature overnight. The solvent was removed by vacuum evaporation. The residue was dissolved in deionized water directly for the transamination study without further purification.

4.4 Transaminations by the dendrimers

100 μl of aqueous dendrimers solution (ca. $5.0 \times 10^{-3}\text{M}$), 100 μl of aqueous α -ketoacids solution ($5.0 \times 10^{-2}\text{M}$) and 25 μl of aqueous EDTA solution ($1.0 \times 10^{-2}\text{M}$) were added to a vial. The pH of the solution was adjusted to 7.5 with NaOH ($1.0 \times 10^{-2}\text{M}$) and HCl ($1.0 \times 10^{-2}\text{M}$). The reaction at room temperature usually took 6-24 hrs.

4.5 Assay of the amino acid products

The reaction mixture from transamination was extracted with CH_2Cl_2 , and then the water layer containing the amino acid was collected and the volume of the aqueous media was reduced to 80 μl . 10 μl of a derivatizing solution containing 0.2M o-phthalaldehyde and 0.2M N-acetyl-L-cysteine in methanol, and 10 μl of an aqueous buffer solution of 1.0 M K_2HPO_4 (pH 8.0) were added. The resulted solutions of the diastereomeric 1-thioisoindoles were analyzed by reverse-phase HPLC using aqueous buffer/THF/acetonitrile with UV-vis detection at 344 nm.

The retention time for the derivatized D-phenylalanine, L-phenylalanine, D-alanine and L-alanine were ~21.1, ~22.3, ~16.5 and ~18.8 min respectively.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work has been supported by the NIH and NSF.

References

1. Liu, L.; Breslow, R. Artificial Enzymes. Breslow, R., editor. Wiley-VCH; Weinheim, Germany: 2005. p. 37-62.

2. Zimmerman SC, Czarnik AW, Breslow R. *J Am Chem Soc* 1983;105:1694–1695.
3. Zimmerman SC, Breslow R. *J Am Chem Soc* 1984;106:1490–1491.
4. Breslow R, Czarnik AW, Lauer M, Leppkes R, Winkler J, Zimmerman S. *J Am Chem Soc* 1986;108:1969–1979.
5. Chmielewski J, Breslow R. *Heterocycles* 1987;25:533.
6. Breslow R, Chmielewski J, Foley D, Johnson B, Kumabe N, Varney M, Mehra R. *Tetrahedron* 1988;44:5515–5524.
7. Lei Liu L, Ronald Breslow R. *Tetrahedron Lett* 2001;42:2775–2777.
8. Liu L, Rozenman M, Breslow R. *Bioorg Med Chem* 2002;10:3973–3979. [PubMed: 12413849]
9. Breslow R, Hammond M, Lauer M. *J Am Chem Soc* 1980;102:421–422.
10. Breslow R, Czarnik AW. *J Am Chem Soc* 1983;105:1390–1391.
11. Czarnik AW, Breslow R. *Carbohydrate Res* 1984;128:133–139.
12. Breslow R, Canary JW, Varney M, Waddell ST, Yang D. *J Am Chem Soc* 1990;112:5212–5219.
13. Fasella E, Dong SD, Breslow R. *Bioorg Med Chem* 1999;7:709–714. [PubMed: 10400323]
14. Winkler J, Coutouli-Argyropoulou E, Leppkes R, Breslow R. *J Am Chem Soc* 1983;105:7198–7199.
15. Liu L, Rozenman M, Breslow R. *J Am Chem Soc* 2002;124:12660–12661. [PubMed: 12392403]
16. Liu L, Breslow R. *J Am Chem Soc* 2002;124:4978. [PubMed: 11982360]
17. Zhou W, Liu L, Breslow R. *Helv Chim Acta* 2003;86:3560–3567.
18. Liu L, Breslow R. *J Am Chem Soc* 2003;125:12110. [PubMed: 14518994]
19. Liu L, Zhou W, Chruma JJ, Breslow R. *J Am Chem Soc* 2004;126:8136–8137. [PubMed: 15225053]
20. Chruma JJ, Liu L, Zhou W, Breslow R. *Bioorg Med Chem* 2005;13:5873–5883. [PubMed: 15953728]
21. Zhou W, Yerkes N, Chruma J, Liu L, Breslow R. *Bioorg Med Chem Lett* 2005;15:1351. [PubMed: 15713385]
22. Breslow R, Bandyopadhyay S, Levine M, Zhou W. *ChemBioChem* 2006;7:1491–1496. [PubMed: 16977662]
23. Supporting Information of Reference 18
24. Tomalia DA, Baker H, Hall M, Kallos G, Rock J, Simth P. *Polym J* 1985;17:117.
25. Kunishima M, Kawachi C, Iwasaki F, Terao K, Tani S. *Tetrahedron Lett* 1999;40:5327. Kunishima M, Kawachi C, Hioki K, Terao K, Tani S. *Tetrahedron* 2001;57:1551. Kunishima M, Kitao A, Kawachi C, Watanabe Y, Iguchi S, Hioki K, Tani S. *Chem Pharm Bull* 2002;50(4):549. [PubMed: 11964009]

Figure 1.

The calculated structures of the G-3 dendrimer with an attached pyridoxamine in water solution. Structure a, the tube version, shows how close the pyridoxamine is to some of the dimethylphenylalanine units that are attached to the ends of the dendrimer chains, explaining why the transaminations are enantioselective. Structure b, the space filling version of structure a, shows that the dendrimer is tightly packed, with no interior space. Structure c is the same space filling version rotated 90°, to show again the tight packing.

Scheme 1.

Scheme 2.

Table 1

The % S ee values in the transamination reactions between the dendrimers and phenylpyruvic acid or pyruvic acid in aqueous media^{a-c}

dendrimer	Phenylpyruvic acid	Pyruvic acid
G-1-Me2	23.9 ± 0.7	21.3 ± 0.4
G-2-Me2	42.8 ± 0.2	39.4 ± 0.9
G-3-Me2	45.4 ± 1.2	42.2 ± 3.7
G-4-Me2	38.9 ± 0.4	42.4 ± 2.3

^a All % ee values are averaged from two or three trials.

^b Transamination reaction condition: dendrimers(ca. 5.0×10^{-3} M), α -keto acids (5.0×10^{-2} M), EDTA(1.0×10^{-2} M), pH = 7.5, t = 20-25°C.

^c The amino acid products were derivatized with o-phthalaldehyde and N-acetyl-L-cysteine before analysis by reverse-phase HPLC, as described elsewhere [ref ⁸, and Experimental].