for cases of lung and heart disease. It is also a useful technique in the management of the second stage of labour in primigravidae.

The risk of aspiration pneumonia is obviated, and for this reason the technique can help to reduce mortality and morbidity.

There were no foetal deaths that could be attributed to the anaesthetic used.

It is suggested that this is a simpler form of analgesia than caudal and spinal analgesia, and one which allows for full oxygenation of the mother and the foetus.

The maternal complications were comparatively slight and the length of stay in hospital was not increased, though 9 out of the 57 patients had oedema, bruising, or haematoma formation.

In three cases another form of anaesthesia was given in order to accomplish a technically more difficult delivery.

My thanks are due to Miss Janet Bottomley and Mr. Oswald Lloyd for their permission to perform the pudendal nerve blocks and to report the results.

REFERENCES

- British Medical Journal*, 1955, 2, 489, 676, 736, 791.
Dieckmann, W. J. (1945). *Amer. J. Obstet. Gynec.*, 50, 28.
Douglas, G. W., and Vosburgh, L. F. (1951). *Ibid.*, 62, 1253.
Dutton, W. A. W. (1955). *Lancet*, 2, 1368.
Eastman, N. J. (1941). *Proc. Amer. Congr. Obstet. Gynaec.*, 1, 191.
Gilliat, W. (1949). *Trans. 12th Brit. Congr. Obstet. Gynaec.*, p. 269. London.
Hall, C. C. (1940). *J. Amer. med. Ass.*, 114, 728.
Klink, E. W. (1953). *Obstet. and Gynec.*, 1, 137.
Law, R. G., and Ransom, S. G. (1954). *British Medical Journal*, 1, 562.
Lock, F. R., and Greiss, F. C. (1955). *Amer. J. Obstet. Gynec.*, 70, 861.
Mendelson, C. L. (1946). *Ibid.*, 52, 191.
Merrill, R. B., and Hingson, R. A. (1951). *Curr. Res. Anesth.*, 30, 121.
Parker, R. B. (1954). *British Medical Journal*, 2, 65.
Willcox, A. (1949). *Lancet*, 1, 438.

FLUID CONTENT OF THE NUCLEUS PULPOSUS AS A FACTOR IN THE DISK SYNDROME

FURTHER OBSERVATIONS

BY

H. D. BUSH, Ph.D., A.Inst.P.

W. G. HORTON, B.Sc., A.R.I.C.

D. L. SMARE, Ph.D., M.Sc.(Tech.), F.Inst.P.

Bradford Technical College

AND

A. NAYLOR, Ch.M., M.Sc., F.R.C.S.

Consultant in Orthopaedic Surgery, Bradford Hospitals

The possibility that the hydrophilic property of the nucleus pulposus of the intervertebral disk has some bearing upon the development and symptomatology of low back pain, sciatica, brachial neuritis, and disk prolapse was suggested by Charnley (1952) and by us (Naylor, 1951; Naylor and Smare, 1953). The swelling of the nucleus due to an increased fluid content is thought to cause increased internal pressure within the disk, thus stretching the annulus fibrosus and stimulating the nerve-endings contained therein. Whilst the adult intervertebral disk has a poor blood supply, a considerable fluid interchange can occur, as shown by the rapid absorption of 35% diodrast when used in disk nucleography (Cloward and Buzaid, 1952). Charnley further showed that the absorption of fluid by the isolated nucleus could result in considerable pressure being developed when the nucleus was contained in a plethysmograph.

These results were obtained with nuclear material which was (a) removed from the disk, the annulus being disregarded; and (b) was not maintained at constant volume.

A preliminary investigation was carried out by two of us (Naylor and Smare, 1953) to discover whether the nucleus contained within a lumbar disk could absorb enough fluid to develop a high intra-disk pressure that might account for the symptoms associated with disk lesions. Isolated cadaveric disks, from subjects of varying ages, were immersed in water or isotonic saline solution and the pressure developed within the disk was recorded by means of a mercury manometer. Pressures of between 8 and 300 cm. mercury (1.5–58.1 lb. per square inch) were obtained from young subjects, but no pressures were recorded in disks from aged subjects in whom advanced degeneration was found to be present. A more comprehensive survey has now been made on disks from subjects in the age range 19–87 years, using the apparatus described below.

Apparatus

A diagram of the apparatus is shown in Fig. 1. A standard Bourdon gauge type pressure indicator A, reading up to a maximum of 60 lb. per square inch (4.2 kg. per square cm.), was connected to a copper tube, B, of $\frac{1}{2}$ in. (1.3 cm.) internal diameter. Tube E, a glass capillary about 8 in. (20 cm.) long and 1 mm. internal bore, was attached by pressure-tight joints to B and to a 20-gauge serum needle F. A metallic bellows, C, closed at one end, was connected to tube B and could be compressed by means of screw D. The whole apparatus above point X was filled with water, while below this point there was an unbroken column of mercury as far as the tip of the serum needle. The mercury-water interface was arranged to be about halfway along the glass capillary, and consequently was always visible.

When used the needle was inserted through the annulus fibrosus of the disk into the nucleus pulposus. On absorption of fluid by the nucleus, expansion of the latter forced mercury up the serum needle. The bellows, C, was compressed until the mercury meniscus returned to its original position. The pressure recorded by the gauge was thus the pressure developed in the nucleus. An indicator (not shown in Fig. 1) was used to enable the mercury meniscus to be adjusted with precision.

Experimental Procedure

Fresh lumbar disks, usually in sets of five (L 1–2, L 2–3, L 3–4, L 4–5, L 5–S 1), were obtained from necropsy subjects with a plate of bone on each side (about $\frac{1}{4}$ in. (6 mm.) thick, this being essential for maintaining the nucleus under conditions corresponding as far as possible with those *in vivo*). Forces generated by the increase in volume of the nucleus will consequently result in a slight increase in the circumference of the annulus fibrosus and not in a bulging of the bone end-plates. The annulus of the disk under test was pierced by a hollow needle of the same gauge as that used in the apparatus, and the needle attached to the apparatus inserted along the track made. This minimized the risk of blockage of the needle by tissue. Each disk was then totally immersed in distilled water contained in a Petri dish.

FIG. 1.—Diagram of apparatus (see text).

By means of this procedure, pressure readings were taken at intervals of about three hours over a period of up to 48 hours. The Petri dishes were immersed in a temperature-controlled water bath, enabling all experiments to be carried out with disks maintained at a temperature of 22° C.

Results

Some 137 disks from 29 subjects (both male and female) aged from 19 to 87 years were examined on the apparatus. The largest number was in the age group 41–60 years; unfortunately, fewer specimens below 41 years were available. Fig. 2 shows the results obtained. In 30 disks no

FIG. 2.—Scattergram of pressure recorded in 107 cases.

pressure was recorded, final pressures of between 0.5 and 42 lb. per square inch (0.04 and 3 kg. per square cm.) being recorded by the other 107. The numbers of disks (expressed as a percentage) developing a given pressure for the age groups considered—namely, 19–40 years (29 disks), 41–60 years (59 disks), 61–90 years (49 disks)—were plotted on histograms (Fig. 3).

The following features were noted: (1) The wide variation in pressure recorded not only with age but also by disks from any one subject. (2) The relatively large percentage of disks recording pressures of 30–45 lb. per square inch (2.1–3.16 kg. per square cm.) in the age group 41–60 years. This feature may also be characteristic of the age group 19–40 years, but lack of data precludes any certainty. (3) The relatively low pressures developed in subjects over 60 years of age. (The above features can also be inferred from Fig. 2.)

The rate of pressure increase varied considerably for each individual disk. No consistent correlation between age and rate of pressure increase was apparent.

At the end of each set of experiments the disks were removed and sectioned with a view to attempting to correlate the pressure developed with the conditions prevailing in each nucleus. As expected, degeneration (as measured by increased hardness and discoloration from white to brown or grey) and an increased number of cavities, usually filled with mercury, appeared frequently in aged subjects. On inserting a serum needle into a disk the mercury has been observed to fall, indicating cavities of up to 0.1 ml. in volume in the proximity of the needle-tip. This is consistent with the demonstration of cavities by means of nucleography (Lindblom, 1950; Erlacher, 1952).

Discussion

The hydrophilic properties of isolated nuclear material may be explained by a process regulated by two opposing factors—namely, (a) osmotic pressure differences between the protein gel and surrounding medium, this being duly influenced by the mucopolysaccharide content; and (b) the cohesive forces of the tissue. Equilibrium is established when these forces are equal and opposite.

Sylvén *et al.* (1952) have shown that more hydrophilic groups of mucopolysaccharide become attached to the collagen molecule with advancing age of the nucleus, and this may result in decreased water-binding capacity. The precipitation of large orientated collagen fibres and possibly of elastin (Happey, Horton, and Naylor, 1956) will decrease the gel concentration within the nucleus. Both the above-

FIG. 3.—Percentage of disks developing a given pressure, arranged according to age groups.

mentioned factors should lead to a decrease in the swelling and water uptake of the nucleus with advancing age. This is in agreement with the observations of Göcke (1932).

In our experiments concerning the increase in pressure in the nucleus pulposus, the nucleus has been confined within the annulus fibrosus. This represents a close approach to conditions appertaining *in vivo*.

It is suggested that the pressure developed in the nucleus pulposus in this state may be affected by the following factors: (1) the amount of hydrophilic mucopolysaccharide material; (2) the concentration of protein gel in the nucleus; (3) the presence of cavities of differing size in the nuclei pulposi, which occur more often in disks from aged subjects; and (4) variations in the elasticity of the annulus fibrosus. Experimental evidence (Happey, Horton, and Naylor, 1956)

has been obtained which indicates that the elasticity of the annulus fibrosus may vary with its position in the disk, and, furthermore, the elasticity may be different for each individual disk.

The differences in mucopolysaccharide content occurring in the nuclei of disks from any given subject are probably insufficient to account for the large differences in pressure encountered within such disks. It is suggested that the controlling factor may be the variations in the osmotic pressure due to differences in gel concentration within each nucleus pulposus. Any decrease in concentration resulting from increased precipitation of collagen would automatically reduce the water uptake and hence decrease the pressure developed. The implications are the development of a high pressure in one disk together with the action of body weight and muscle action, which may be sufficient to cause herniation in this disk but not in another. In this connexion it should be noted that local variations in the precipitation of protein within a nucleus occur in, and after, the fourth decade. On the basis of the above argument, this may lead to differences in pressure within the nucleus when immersed in aqueous solutions.

The observations of Göcke (1932) on isolated nuclei are in agreement with our observations on nuclei surrounded by an annulus—namely, that water uptake and therefore pressure decrease with advancing age. This may be readily explained, since the frequency of occurrence of cavities and decreased number of free polar mucopolysaccharide groups augmented by the increased gel concentration will tend to lower the pressure developed in aged subjects. Disks from subjects in the age group 20–55 years develop a high intradisk pressure, after which the pressure progressively decreases with age (Figs. 2 and 3). This observation may afford some explanation of the well-known fact that not only is disk prolapse more common before the age of 50 years than after and rare after 60, but that attacks in a patient become less frequent with advancing age, since the decreasing pressure developed should make excessive stress on the annulus less likely.

A considerable number of disks in all age groups failed to record a pressure. This may result from the appearance of cavities due to removal of the forces of compression from the disk (these cavities should be distinguished from those due to chemical changes). There is no apparent correlation between the presence of these cavities and the development of small or zero pressures, for disks with large cavities do occasionally record considerable pressures.

Summary

Further extended measurements on the intradisk pressure developed on immersion in distilled water have been made, using large numbers of disks. Though disks from the same subject were found to generate widely differing pressure, there is a general trend towards a lower pressure with increasing age. The possible causes of such behaviour are discussed.

One of us (W. G. H.) is indebted to the Bradford Education Authority and the Bradford "A" Group Hospital Management Committee for a joint scholarship to carry out this work.

REFERENCES

- Charnley, J. (1952) *Lancet*, 1, 124.
 Cloward, R. B., and Buzaid, L. C. (1952). *Amer. J. Roentgenol.*, 68, 552.
 Erlacher, P. R. (1952). *J. Bone Jt Surg.*, 34B, 204.
 Göcke, C. (1932). *Arch. orthop. Unfall-Chir.*, 31, 42.
 Haphey, F., Horton, W. G., and Naylor, A. (1956). In preparation.
 Lindblom, K. (1950). *Acta radiol. (Stockh.)*, 34, 321.
 Naylor, A. (1951). *Ann. roy. Coll. Surg. Engl.*, 9, 158.
 — and Smare, D. L. (1953). *British Medical Journal*, 2, 975.
 Sylvén, B., Snellman, O., Paulson, S., and Hirsch, C. (1952). *Acta orthop. scand.*, 22, 175.

NEPHROSIS TREATED WITH INTRAVENOUS T.A.B.

BY

DENNIS MENDEL, M.D., M.R.C.P.

*Senior Lecturer in Physiology, University College, Ibadan;
Formerly R.M.O., Lymington Hospital, Hants*

AND

R. J. MCGILL, M.B., M.R.C.P.Ed.

*Physician, Royal South Hants Hospital, Southampton,
and Lymington Hospital, Hants*

In the search for effective treatment for nephrosis—Type II nephritis of Ellis (1942)—the observation that infection might be associated with remission (Aldrich, 1926; Ball and Evans, 1932) led to the use of measles by Janeway and Hutchings (1947) and of malaria by Gairdner (1952) and Byrne (1952), the last of whom related the success of this form of treatment with the irregular remissions produced by corticotrophin (A.C.T.H.) (Barnett *et al.*, 1950; Riley, 1950).

Intravenous T.A.B. vaccine, a simple and well-known means of producing pyrexia, was used in the case reported here, and it regularly brought about a diuresis. The better-controlled production of pyrexia so induced allowed study of the mechanism involved. Below we give our observations on the diuresis, albuminuria, plasma proteins, plasma cholesterol, and the 17-ketosteroid excretion.

Case Report

A woman aged 53 noticed ankle oedema in September, 1951, which rapidly extended. Treatment at another hospital in April, 1952, with mercurial diuretics (mersalyl), plasma transfusion, urea, and a high-protein/low-salt diet, had not been beneficial, and Southey's tubes had been used. Two weeks after her discharge oedema recurred, and this steadily progressed in spite of mersalyl and urea. She had been bedridden for some weeks before admission. There was no previous history of acute nephritis.

On admission in December, 1952, she had gross generalized oedema, with pleural effusions and ascites, and the knees could not be flexed to the slightest degree. The blood pressure was 140/70. The fundi were normal. The thyroid was firm, with slight generalized enlargement, but there was no clinical evidence of toxicity or myxoedema. No other abnormality was detected.

Her urine contained albumin (Esbach, 20 g. per litre), but there was no other significant abnormality. Blood urea was 30 mg. per 100 ml.; plasma albumin, 2 g. per 100 ml.; plasma globulin, 1.5 g. per 100 ml.; plasma cholesterol, 680 mg. per 100 ml.

A low-salt/high-protein diet and unrestricted fluids were given throughout the period of observation, and fluid balance and Esbach estimations were recorded daily. The patient was at rest in bed for the first 12 weeks.

After observation for several weeks, during which mersalyl failed to produce a diuresis, intravenous T.A.B. therapy was started. The 24-hour fluid intake varied between 0.9 and 1.2 litres, and the 24-hour urinary output between 0.3 and 0.7 litre during this control period.

T.A.B. vaccine, containing 2,000 million organisms per ml., was used. The dose required was diluted to 1 ml. with sterile normal saline. Doses of 200–300 million organisms produced moderate fever (maximum temperature 102° F.—38.9° C.) and a generalized reaction (vomiting, backache) lasting up to 24 hours. Smaller doses had no appreciable effect. Five out of six injections in this dose range produced a successful response.