has been obtained which indicates that the elasticity of the annulus fibrosus may vary with its position in the disk, and, furthermore, the elasticity may be different for each individual disk.

The differences in mucopolysaccharide content occurring in the nuclei of disks from any given subject are probably insufficient to account for the large differences in pressure encountered within such disks. It is suggested that the controlling factor may be the variations in the osmotic pressure due to differences in gel concentration within each nucleus pulposus. Any decrease in concentration resulting from increased precipitation of collagen would automatically reduce the water uptake and hence decrease the pressure developed. The implications are the development of a high pressure in one disk together with the action of body weight and muscle action, which may be sufficient to cause herniation in this disk but not in another. In this connexion it should be noted that local variations in the precipitation of protein within a nucleus occur in, and after, the fourth decade. On the basis of the above argument, this may lead to differences in pressure within the nucleus when immersed in aqueous solutions.

The observations of Göcke (1932) on isolated nuclei are in agreement with our observations on nuclei surrounded by an annulus—namely, that water uptake and therefore pressure decrease with advancing age. This may be readily explained, since the frequency of occurrence of cavities and decreased number of free polar mucopolysaccharide groups augmented by the increased gel concentration will tend to lower the pressure developed in aged subjects. Disks from subjects in the age group 20–55 years develop a high intradisk pressure, after which the pressure progressively decreases with age (Figs. 2 and 3). This observation may afford some explanation of the well-known fact that not only is disk prolapse more common before the age of 50 years than after and rare after 60, but that attacks in a patient become less frequent with advancing age, since the decreasing pressure developed should make excessive stress on the annulus less likely.

A considerable number of disks in all age groups failed to record a pressure. This may result from the appearance of cavities due to removal of the forces of compression from the disk (these cavities should be distinguished from those due to chemical changes). There is no apparent correlation between the presence of these cavities and the development of small or zero pressures, for disks with large cavities do occasionally record considerable pressures.

Summary

Further extended measurements on the intradisk pressure developed on immersion in distilled water have been made, using large numbers of disks. Though disks from the same subject were found to generate widely differing pressure, there is a general trend towards a lower pressure with increasing age. The possible causes of such behaviour are discussed.

One of us (W. G. H.) is indebted to the Bradford Education Authority and the Bradford "A" Group Hospital Management Committee for a joint scholarship to carry out this work.

REFERENCES

- Charnley, J. (1952) *Lancet*, 1, 124.
Cloward, R. B., and Buzaid, L. C. (1952). *Amer. J. Roentgenol.*, 68, 552.
Erlacher, P. R. (1952). *J. Bone Jt Surg.*, 34B, 204.
Göcke, C. (1932). *Arch. orthop. Unfall-Chir.*, 31, 42.
Happley, F., Horton, W. G., and Naylor, A. (1956). In preparation.
Lindblom, K. (1950). *Acta radiol. (Stockh.)*, 34, 321.
Naylor, A. (1951). *Ann. roy. Coll. Surg. Engl.*, 9, 158.
— and Smare, D. L. (1953). *British Medical Journal*, 2, 975.
Sylvén, B., Snellman, O., Paulson, S., and Hirsch, C. (1952). *Acta orthop. scand.*, 22, 175.

NEPHROSIS TREATED WITH INTRAVENOUS T.A.B.

BY

DENNIS MENDEL, M.D., M.R.C.P.

*Senior Lecturer in Physiology, University College, Ibadan;
Formerly R.M.O., Lymington Hospital, Hants*

AND

R. J. MCGILL, M.B., M.R.C.P.Ed.

*Physician, Royal South Hants Hospital, Southampton,
and Lymington Hospital, Hants*

In the search for effective treatment for nephrosis—Type II nephritis of Ellis (1942)—the observation that infection might be associated with remission (Aldrich, 1926; Ball and Evans, 1932) led to the use of measles by Janeway and Hutchings (1947) and of malaria by Gairdner (1952) and Byrne (1952), the last of whom related the success of this form of treatment with the irregular remissions produced by corticotrophin (A.C.T.H.) (Barnett *et al.*, 1950; Riley, 1950).

Intravenous T.A.B. vaccine, a simple and well-known means of producing pyrexia, was used in the case reported here, and it regularly brought about a diuresis. The better-controlled production of pyrexia so induced allowed study of the mechanism involved. Below we give our observations on the diuresis, albuminuria, plasma proteins, plasma cholesterol, and the 17-ketosteroid excretion.

Case Report

A woman aged 53 noticed ankle oedema in September, 1951, which rapidly extended. Treatment at another hospital in April, 1952, with mercurial diuretics (mersalyl), plasma transfusion, urea, and a high-protein/low-salt diet, had not been beneficial, and Southey's tubes had been used. Two weeks after her discharge oedema recurred, and this steadily progressed in spite of mersalyl and urea. She had been bedridden for some weeks before admission. There was no previous history of acute nephritis.

On admission in December, 1952, she had gross generalized oedema, with pleural effusions and ascites, and the knees could not be flexed to the slightest degree. The blood pressure was 140/70. The fundi were normal. The thyroid was firm, with slight generalized enlargement, but there was no clinical evidence of toxicity or myxoedema. No other abnormality was detected.

Her urine contained albumin (Esbach, 20 g. per litre), but there was no other significant abnormality. Blood urea was 30 mg. per 100 ml.; plasma albumin, 2 g. per 100 ml.; plasma globulin, 1.5 g. per 100 ml.; plasma cholesterol, 680 mg. per 100 ml.

A low-salt/high-protein diet and unrestricted fluids were given throughout the period of observation, and fluid balance and Esbach estimations were recorded daily. The patient was at rest in bed for the first 12 weeks.

After observation for several weeks, during which mersalyl failed to produce a diuresis, intravenous T.A.B. therapy was started. The 24-hour fluid intake varied between 0.9 and 1.2 litres, and the 24-hour urinary output between 0.3 and 0.7 litre during this control period.

T.A.B. vaccine, containing 2,000 million organisms per ml., was used. The dose required was diluted to 1 ml. with sterile normal saline. Doses of 200–300 million organisms produced moderate fever (maximum temperature 102° F.—38.9° C.) and a generalized reaction (vomiting, backache) lasting up to 24 hours. Smaller doses had no appreciable effect. Five out of six injections in this dose range produced a successful response.

Five individual periods of diuresis have been studied. Some of the features of the response during two of the periods (second and third) are shown in the Chart. The data for the whole group are described below.

Diuresis.—Relative oliguria occurred on the day of injection. Diuresis followed in 24–36 hours and became manifest by a sudden large nocturnal output. The duration of the response was from 7 to 10 days, but lasted 35 days on the solitary occasion when a second injection was given during a successful response. The termination of the

100 ml. at the onset, and 672 and 696 mg. per 100 ml. at the termination. During the diuresis lasting 35 days 920 and 938 mg. per 100 ml. were recorded on the 21st and 35th days respectively.

17-Ketosteroids.—The 24-hour urinary excretion of 17-ketosteroids was measured at random intervals during the periods of diuresis. Values of 1.3 and 2.9 mg. per 24 hours were obtained at the onset of diuresis, 2.2 mg. 48 hours after the onset of diuresis, and 1.9 and 3.1 mg. on the 14th and 35th days in the case of the long-continued diuresis.

Discussion

Intravenous T.A.B. vaccine had a definite diuretic action in this case of nephrosis. The previous unrelenting course of the disease and the fact that individual large diureses occurred within a constant interval after T.A.B. would eliminate the occurrence of spontaneous remission.

The mechanism of the diuresis has been shown to be independent of plasma protein concentration, although secondary conservation of albumin occurs and cannot easily be explained by the three known mechanisms of producing a diuresis in nephrosis—namely, increase in plasma osmotic pressure, osmotic diuresis, and increased renal loss of sodium.

The action of pyrogens upon renal blood flow (Smith, 1951) is to induce an initial short period of ischaemia, followed by hyperaemia lasting four hours or more and reaching values of 30–60% above normal. There is, however, no change in filtration rate, and these transitory changes cannot be held to be responsible for the continued diuresis.

It has been shown by Slessor (1951) that hormones of the cortisone type may be concerned with the inactivation of the antidiuretic hormone. In the light of this work it is here suggested that the diuresis occurring during fever therapy for nephrosis might be due to increased corticoid output inactivating antidiuretic hormone, thus causing a primary water diuresis. Although in the above case we have been unable to show that following T.A.B. therapy there is an increase in urinary corticoids associated with a lowering of blood cholesterol, the diuresis in response to corticotrophin (Riley, 1951) is also independent of plasma proteins, and is associated in most cases with a lowering of blood cholesterol.

Summary

A case of Type II nephritis treated with intravenous T.A.B. is recorded.

Intravenous T.A.B. regularly produced a diuresis and a decrease in albuminuria. The diuresis was independent of changes in plasma proteins or cholesterol. A possible mechanism is suggested.

We are indebted to Mr. G. H. Pitt for referring this case to us.

REFERENCES

- Aldrich, C. A. (1926). *Amer. J. Dis. Child.*, **32**, 163.
Ball, W. G., and Evans, G. (1932). *Diseases of the Kidney*. Churchill, London.
Barnett, H. L., McNamara, H., McCrory, W., Forman, C., Rapoport, M., Michie, A., and Barbero, G. (1950). *Amer. J. Dis. Child.*, **80**, 519.
Byrne, E. A. J. (1952). *Lancet*, **1**, 844.
Ellis, A. (1942). *Ibid.*, **1**, 1.
Gairdner, D. (1952). *Ibid.*, **1**, 842.
Janeway, C. A., and Hutchings, G. (1947). *Amer. J. Dis. Child.*, **73**, 242.
Riley, C. M. (1950). *Ibid.*, **80**, 520.
— (1951). *Pediatrics*, **7**, 457.
Slessor, A. (1951). *J. clin. Endocr.*, **11**, 700.
Smith, H. W. (1951). *The Kidney*, p. 446. Oxford Univ. Press, New York.

Chart showing part of day-to-day reactions to T.A.B.

response, recognized by a rise in urinary albumin to more than 10 g. per litre and by a fall in urinary output below 1.2 litres, was also a sudden event. The average daily urinary output for the five periods was 1.8, 1.5, 2.2, 1.8, and 1.3 litres. Fluid intake remained between 0.9 and 1.2 litres. The net fluid loss for the whole of the five periods was 29.9 litres. A weight loss of approximately 40 lb. (18 kg.) was recorded.

Albumin Loss.—At the onset of diuresis the total urinary loss of albumin increased (see Chart). This was followed after 48 hours by a decrease in total albumin loss. Compared with an initial average loss of 12 g./24 hours, the average 24-hour losses during the five periods of treatment were 1.5, 7.1, 8.0, 6.5, and 4.7.

Plasma Proteins.—Determinations of plasma albumin and globulin were made at the onset and termination of three of the periods of diuresis. At the onset of diuresis these values were albumin 1.4, 1.7, and 1.6 g. per 100 ml., and globulin 1.6, 1.8, and 2.6 g. per 100 ml.; and at the termination of diuresis, albumin 3.2, 2.4, and 1.4 g. per 100 ml., and globulin 1.3, 2.1, and 1.6 g. per 100 ml. On the solitary occasion when diuresis continued for 35 days the plasma proteins at the end of this period were albumin 1.6 g. and globulin 2.9 g. per 100 ml. It can be seen that the diuresis was independent of plasma protein concentration.

Plasma Cholesterol.—Estimations of plasma cholesterol were similarly made at the onset and termination of two periods of diuresis. These values were 474 and 642 mg. per

Miss Gertrude L. Annan has been appointed librarian of the New York Academy of Medicine in place of Miss Janet Doe, who is retiring after thirty years' service at the Academy. Miss Doe has made a number of valuable contributions to medical bibliography, including a notable *Bibliography of the Works of Ambroise Paré*, 1943, and the editing of the *Handbook of Medical Library Practice*, 1943; 2nd ed., 1956.