FIG. 4.—Case 2. Photomicrograph of section of liver showing small miliary metastases of the same reticulum-cell tumour as Figs. 3 and 4. ($\times 340$.)

Comment

Death was due to bronchopneumonia following pulmonary, hepatic, and lymph-node metastases of a malignant reticulum-cell tumour, probably arising in the mesenteric lymph nodes.

Discussion

While the cause of idiopathic steatorrhoea (as of sprue) is as yet unknown, it has been recognized for some time that a small proportion of all cases of steatorrhoea are caused by simple obstruction of mesenteric lacteals by a widespread mesenteric lymphadenitis. Ryle (1936) described how such obstruction may occur from tuberculous mesenteric nodes with the production of the sprue syndrome, and Hurst, Payling Wright, and Ryle (1942) describe the history and necropsy of such a case. Hamilton Fairley and Mackie (1937) described a sprue-like syndrome in adults associated with lymphoma, lymphadenoma, and lymphosarcoma of mesenteric nodes, and caused, as they suggest, by defective intestinal absorption from lymphatic obstruction. In two of their four cases anaemia was present, macrocytic in one and normocytic in the other.

For neoplastic disease of mesenteric nodes to produce the sprue syndrome without other evidence of disease it is necessary that the neoplastic condition be a low-grade one, of slow growth, occurring primarily in mesenteric nodes, remaining localized in that area for a considerable interval before becoming generalized and manifesting itself in peripheral lymph nodes or elsewhere. These conditions existed in the two cases here described. That the presenting symptom in Case 1 was that of macrocytic anaemia of pregnancy is not surprising, since pregnancy is a recognized precipitating factor of both sprue and idiopathic steatorrhoea (Morgensen, 1937). That the two cases here described were under observation for four years and two years respectively without developing palpable peripheral lymphatic nodes, hepato-splenomegaly, or other signs of generalized spread indicates the slow-growing, relatively benign nature of the neoplastic process; and careful study of biopsy and necropsy material from these cases has led us to place them in Robb-Smith's category of reticulum-celled medullary reticulosis (Letterer-Siwe syndrome). While this condition has most often been recorded in children (Wallgren, 1940; Varga *et al.*, 1948), where it may be confused with aleukaemic leukaemia, Robb-Smith has observed the condition in adults, in whom a very different clinical picture is produced; the condition runs a much more chronic course, is associated with lymphatic node enlargement, splenomegaly, severe anaemia, and pyrexia; and symptoms of steatorrhoea have been observed. This condition merges with the lesions of the lipoidoses and eosinophil granuloma of bone, in which distinction between a chronic inflammatory condition and true neoplasia is difficult, and, on the other hand, with the more obviously malignant "reticulum-cell sarcoma" of Willis (1948). Willis's Case XII in particular shows some striking similarities to the cases here described.

Summary

Two cases of Letterer-Siwe disease in adults are recorded, with early involvement of mesenteric lymph nodes, giving rise to the sprue syndrome with severe macrocytic anaemia responding only to folic acid. The clinical picture and the pathology of the disease are discussed.

My thanks are due to Professor Payling Wright for helpful criticism; to Dr. J. McMurray for the necropsy report of Case 2, and for providing necropsy material; and to Dr. B. B. Hosford for his co-operation.

REFERENCES

- Fairley, N. Hamilton, and Mackie, F. P. (1937). *British Medical Journal*, **1**, 375.
 Hurst, Sir A., Wright, G. Payling, and Ryle, J. A. (1942). *Guy's Hosp. Rep.*, **91**, 26.
 Morgensen, E. (1937). *Quart. J. Med.*, **6**, 119.
 Robb-Smith, A. H. T. (1938). *J. Path. Bact.*, **47**, 457.
 Ryle, J. A. (1936). *Natural History of Disease*, p. 151. London.
 Schafer, E. L. (1949). *Amer. J. Path.*, **25**, 49.
 Strang, C. (1950). *Brit. J. Surg.*, **38**, 489.
 Varga, C., Richter, M. N., and De Sanctis, A. G. (1948). *Amer. J. Dis. Child.*, **75**, 376.
 Wallgren, A. (1940). *Ibid.*, **60**, 471.
 Willis, R. A. (1948). *Pathology of Tumours*, 1st ed., p. 777. London.

EFFECT OF MITRAL VALVOTOMY ON RENAL FUNCTION

BY

JOHN HAMMOND, M.B., M.R.C.P.

Research Assistant

JOHN MACKINNON, M.D., M.R.C.P.

Formerly Research Fellow

AND

WILLIAM WHITAKER, M.D., B.Sc., M.R.C.P.

Formerly Senior Lecturer in Medicine

University Department of Medicine, Royal Hospital, Sheffield, and Regional Cardiovascular Unit, City General Hospital, Sheffield

After surgical intervention some patients with mitral stenosis complicated by congestive heart failure improve to such an extent that they no longer require treatment. Merrill (1946) and Mokotoff *et al.* (1948) suggested that the abnormal renal circulation, which has been demonstrated in patients with chronic rheumatic heart disease both with and without congestive heart failure (Merrill, 1946; Aas and Blegen, 1949; Davies and Kilpatrick, 1951; Werkö *et al.*, 1955), was directly responsible for impairment of salt and water excretion in these patients and for the development of heart failure. Therefore it seemed desirable to investigate renal function before and after mitral valvotomy and to attempt to relate any changes to those in the clinical state.

Material and Methods

Fifteen patients, of whom all but one were female, were studied before and after mitral valvotomy.

The clinical state of the patients was assessed both before and after operation, using the grades of disability defined by Baker *et al.* (1952); these may be briefly summarized as 0, no disability; 1, slight disability; 2, moderate disability; 3, severe disability; and 4, total incapacity. The first post-operative assessment of the clinical state was made six months to one year after operation.

Observations on the renal circulation were made about two hours after a meal with the subjects propped up

comfortably in bed at an angle of approximately 45 degrees. The renal plasma flow was measured by sodium *para*-amino-hippuric acid clearance and glomerular filtration rate by inulin clearance, using the technical method described by Dick and Davies (1949). Measurements were made during two accurately timed periods of urine collection of approximately 20 minutes. The filtration fraction, that fraction of the plasma passing through the kidney which is filtered by the glomeruli, was calculated by dividing the glomerular filtration rate by the renal plasma flow. The renal blood flow was calculated from the renal plasma flow and the haematocrit reading.

The diuretic response to ingested water was tested before and after operation in five patients (Cases 1, 2, 8, 10, and 11). Fluids and food were withheld from 6 p.m. on the preceding evening. Between 9 and 9.15 a.m. the patient drank a litre of water and emptied the bladder at 9.15 a.m. The bladder was emptied every 15 minutes for the next four hours and the volume of each 15-minute specimen was measured. From these volumes the maximal rate of urine flow in millilitres per minute and the volume excreted in the first two hours were calculated.

These investigations were carried out a few days before and from two to five weeks after valvotomy.

Results

The effect of mitral valvotomy on the renal circulation, water excretion, and the clinical state is shown in Tables I and II.

TABLE I.—*Effect of Mitral Valvotomy upon Clinical State and Renal Circulation*

Case No.	Age and Sex	Follow-up (Months)	Clinical State (Grades 0 to 4)	Haematocrit (%)	Renal Blood Flow (ml./min.)	Renal Plasma Flow (ml./min.)	Glomerular Filtration Rate (ml./min.)	Filtration Fraction
1	38 F	19	{ a 2 b 1	36 40	695 815	445 489	95 96	0.21 0.20
2	34 F	20	{ a 3 b 1	36 42	350 412	224 239	52 63	0.23 0.26
3	38 F	22	{ a 3 b 2	46 40	417 408	225 245	67 69	0.30 0.28
4	23 F	9	{ a 3 b 1	37 36	560 674	353 432	80 104	0.23 0.24
5	38 F	35	{ a 4 b 2	47 52	323 342	171 164	65 72	0.38 0.44
6	41 F	42	{ a 4 b 4	51.5 37	497 327	241 206	75 83	0.31 0.40
7	35 F	8	{ a 4 b 4	37 41	421 721	248 455	93 108	0.375 0.24
8	43 F	35	{ a 2 b 1	40 43	638 581	383 331	108 93	0.28 0.28
9	40 F	32	{ a 2 b 1	43 40	418 365	239 237	57 66	0.24 0.28
10	38 F	13	{ a 4 b 2	42 39.5	466 411	270 248	72 56	0.27 0.23
11	43 F	36	{ a 3 b 1	35 41	471 276	306 157	76 44	0.25 0.28
12	39 M	6	{ a 3 b 2	46 41	624 732	337 432	81 139	0.24 0.32
13	36 F	36	{ a 4 b 1	37.5 38	428 586	268 357	81 76	0.30 0.21
14	25 F	20	{ a 2 b 1	46.5 45.5	577 597	309 325	106 81	0.34 0.25
15	40 F	20	{ a 3 b 1	48 44	440 465	22.9 260	74 78	0.32 0.30

a = Before operation. b = After operation.

1. *Effect upon Clinical State.*—Before operation five patients (Cases 5, 6, 7, 10, and 13) were in grade 4, six (Cases 2, 3, 4, 11, 12, and 15) in grade 3, and four (Cases 1, 8, 9, and 14) in grade 2. All except three were reassessed one year after operation. One of these three (Case 4) had improved from grade 3 to grade 1 nine months after operation and another (Case 12) had improved from grade 3 to grade 2 six months after operation; both patients subsequently failed to attend for follow-up. The third patient (Case 7)

TABLE II.—*Results of Water Diuresis Tests Before and After Mitral Valvotomy*

Case	2-hourly Volume (ml.)	Maximal Diuresis	
		Rate (ml./min.)	Time (minutes)
1 { after before	296 767	4.9 10.9	67 52
2 { after before	390 222	6.0 4.2	112 52
8 { after before	417 658	7.3 10.0	67 67
10 { after before	247 500	12.5 6.0	97 82
11 { after before	384 341	6.7 5.5	97 97
Normal range (Taylor, 1955)	702 to 1,338	9.7 to 16.9	48 to 104

first appeared slightly improved, but died nine months after valvotomy with congestive heart failure. Six of those followed up for at least one year (Cases 1, 3, 8, 9, 12, and 14) improved by one grade, but four of them (Cases 1, 8, 9, and 14) were only moderately disabled (grade 2) prior to operation. A decrease in disability of two grades was observed in six patients (Cases 2, 4, 5, 10, 11, and 15) and a decrease in disability of three grades in one (Case 13). The remaining patient (Case 6) showed no improvement in the first year and remained in grade 4, but after this she gradually improved to grade 3. In the other cases followed up for more than one year (Table I) there was no change in the clinical condition, but one patient (Case 5) developed hypertension and subsequently died three years later from a cerebral haemorrhage.

2. *Effect Upon the Renal Circulation.*—Accepting a change of at least 20% as being significant, valvotomy was followed by an increase in renal blood flow, renal plasma flow, and glomerular filtration rate in Case 4, and in renal blood and plasma flows unaccompanied by a change in glomerular filtration rate in Cases 7 and 13 (Table I). In Case 12 the renal blood flow showed a slight increase, but because of a fall in the haematocrit reading after operation the renal plasma flow increased to a much greater extent, and this was accompanied by an increase in the glomerular filtration rate. In Case 2 the glomerular filtration rate increased and in Cases 10 and 14 it decreased, with little change in renal blood and plasma flows. In Case 11 valvotomy was followed by a decrease in renal blood flow, renal plasma flow, and glomerular filtration rate, and in Case 6 a slight fall in renal blood flow associated with a decrease in haematocrit caused a marked fall in renal plasma flow, the glomerular filtration rate showing little change. In the remaining cases (Nos. 1, 3, 5, 8, 9, and 15) the changes in renal blood flow, renal plasma flow, and glomerular filtration rate were much smaller (Table I). As a result of these changes in renal plasma flows and glomerular filtration rates the filtration fractions increased in three cases (Nos. 5, 6, and 12), decreased in three (Nos. 7, 13, and 14), and showed little or no change in the remainder (Table I).

3. *Effect Upon the Diuretic Response to Ingested Water.*—After operation the volume of urine excreted in the first two hours after the ingestion of 1 litre of water increased in three cases (Nos. 1, 8, and 10), decreased in one (No. 2), and showed little change in the other, Case 11. In Cases 1 and 8 the maximal rate of excretion increased, whereas in Cases 2, 10, and 11 it decreased. The time of maximal rate of urine flow was earlier after water ingestion in Cases 1, 2, and 10, and the same in Cases 8 and 11 (Table II).

Discussion

All patients in the present series showed abnormally low values for renal blood flow, renal plasma flow, and glomerular filtration rate compared with those found in normals by Davies and Kilpatrick (1951), using a similar clearance technique. These findings are similar to those reported by

Blegen and Aas (1950), Heller and Jacobson (1950), and Werkö *et al.* (1952), and in many of the cases confirm that changes in renal circulation in patients with valvular heart disease occur long before the onset of congestive heart failure. After operation the renal blood flow and renal plasma flow often increased, but did not return to the normal range in any of the patients, and only in Case 12 did the glomerular filtration rate reach normal levels.

The clinical state following valvotomy improved in all except two patients (Cases 6 and 7), but this improvement was not always associated with one in the renal circulation, nor was it consistently accompanied by a change in the diuretic response to water ingestion. In the group with congestive heart failure (patients in grade 4), one patient (Case 13) improved to such an extent—grade 4 to grade 1—that she was only slightly disabled after operation and her renal blood and plasma flows increased greatly. In Cases 5, 6, and 10 there was less improvement in the clinical state, and this was associated with much smaller changes in renal circulation (Table I). The fifth patient with congestive heart failure (Case 7) showed a dramatic increase in her renal blood and plasma flows after valvotomy, but despite this she improved only slightly at first and then relapsed into congestive heart failure. Although all 10 patients without heart failure improved clinically after operation, only two (Cases 4 and 12) showed an improvement in renal circulation, and in one (Case 11) the renal blood and plasma flows and glomerular filtration rate decreased. These results are in accordance with those of Werkö *et al.* (1955) and Judson *et al.* (1955), who reported variable changes in renal blood flow after valvotomy.

Before operation all five patients whose response to the ingestion of 1 litre of water was tested excreted less than 702 ml. of urine in two hours (Table II), which was the lowest value found in normal fasting subjects by Taylor (1955). In all but one patient the maximal rate of diuresis was also below normal, but the time of the maximal urine flow was not delayed beyond the normal range given by Taylor (1955), except in one patient (Table II). After operation there was an improvement in the diuretic response in three patients. The maximal rate of diuresis increased to within normal limits in two patients, and in one of these the two-hourly urine volume also reached normal limits. In all patients maximal diuresis occurred within the normal time range (Table II). There was no apparent correlation between the improvement in diuretic response and changes in the renal circulation, since the patients whose two-hourly urine volumes and maximal rates of diuresis increased showed no significant change in renal blood flow or glomerular filtration rate, and Case 2, whose glomerular filtration rate increased, had a diminished diuretic response after operation. These results are in keeping with those of Judson *et al.* (1955), who reported that post-operative increases in water excretion could not be consistently correlated with any specific change in renal function.

Although these results show that sometimes there is a marked improvement in the renal circulation following mitral valvotomy, the series is too small to allow these changes to be correlated with recovery from or persistence of congestive heart failure. It is obvious that further studies are required, especially in patients with congestive heart failure who recover completely after operation, before it is possible to decide whether improvement in the renal circulation as a result of the operation is directly related to an increase in the capacity to excrete salt and water and the disappearance of congestive heart failure.

Summary

The renal circulation was studied before and after mitral valvotomy in 15 patients, and in five of these the diuretic response to water ingestion was also tested.

Five patients had congestive heart failure prior to operation; six others were severely and four moderately disabled.

Before operation the renal circulation was impaired in all patients and the diuretic response was abnormal in the five patients in whom it was studied.

After operation one patient died with congestive heart failure, another was unchanged, and the remaining 13 showed improvement in their clinical state. The renal circulation improved in five patients, was further reduced in four, and showed little change in six. In three of the five patients studied the diuretic response improved, and in one it became more abnormal.

From this small series there was no obvious correlation between changes in the renal circulation, diuretic response, and the clinical state, although one patient with congestive heart failure who was greatly improved showed a considerable increase in renal blood and plasma flows after operation. The results indicated a need for further study, especially in patients with congestive heart failure prior to operation.

We wish to thank Dr. J. W. Brown, Dr. C. E. Davies, Professor C. H. Stuart-Harris, Mr. J. T. Chesterman, and Mr. A. W. Fawcett for permission to investigate patients under their care, and Mr. A. F. Nichols and Miss S. Exley for technical assistance.

REFERENCES

- Aas, K., and Blegen, E. (1949). *Scand. J. clin. Lab. Invest.*, **1**, 22.
 Baker, C., Brock, R. C., Campbell, M., and Wood, P. (1952). *British Medical Journal*, **1**, 1043.
 Blegen, E., and Aas, K. (1950). *Acta med. scand.*, **138**, 391.
 Davies, C. E., and Kilpatrick, J. A. (1951). *Clin. Sci.*, **10**, 53.
 Dick, A., and Davies, C. E. (1949). *J. clin. Path.*, **2**, 67.
 Heller, B. I., and Jacobson, W. E. (1950). *Amer. Heart J.*, **39**, 188.
 Judson, W. E., Hatcher, J. D., Hollander, W., and Halperin, M. H. (1955). *J. clin. Invest.*, **34**, 1297.
 Merrill, A. J. (1946). *Ibid.*, **25**, 389.
 Mokotoff, R., Ross, G., and Leiter, L. (1948). *Ibid.*, **27**, 1.
 Taylor, W. H. (1955). *Clin. Sci.*, **14**, 725.
 Werkö, L., Ek, J., Bucht, H., and Eliasch, H. (1952). *Scand. J. clin. Lab. Invest.*, **4**, 15.
 ———, Varnauskas, E., Bucht, H., Thomasson, B., and Eliasch, H. (1955). *Amer. Heart J.*, **49**, 823.

SEA-SNAKE BITES

BY

H. A. REID, M.B., M.R.C.P.Ed., D.T.M.&H.

Specialist Physician, General Hospital, Penang, Malaya

Sea-snakes are the most abundant and venomous of the world's reptiles. They inhabit the shores of the Indian and Pacific Oceans, ranging from the Persian Gulf to south Japan, the coasts of tropical Australia, and the southern Pacific islands. In this vast triangle all fishermen know the sea-snake and its hall-mark, the flat rudder-like tail. Water-snakes, also common along these shores, have round tapering tails (see Fig. 1). They are harmless—a fact familiar to the fishermen.

The frequency of particular genera and species of the sea-snake family (Hydrophiidae) varies in different parts of the world, but, generally speaking, the commonest species is *Enhydrina schistosa* (common sea-snake). The adult is 3–4 feet (0.9–1.2 metres) long, about 1–2 inches (2.5–5 cm.) thick, a uniform grey above and whitish below. It is easily recognized by the deep cleft in its chin (see Fig. 2). Young ones have dark-grey bars. Most other species are banded. Some species of *Hydrophis* grow to a length of 9 feet (2.7 metres), but are relatively slender, with small heads. The genus *Astrotia* are the most massive, being up to 4 inches (10 cm.) thick. *Pelamis platurus* has the widest distribution, and has been noted off the western coasts of tropical America and the