of neurological signs, apart from those of increased intracranial pressure, and the absence of convulsive attacks, may have been due to the protection of the cerebral cortex by the thick membrane. Although attempts to isolate organisms from three specimens of pus from the bilateral subdural collections were made, nothing was grown; his clinical condition was one of moderate toxicity, with a persistent low fever and high white-cell count. Vomiting and lethargy were the striking abnormal signs throughout the course of his illness. At operation the tough membrane seemed too firmly attached to the underlying cortex to justify its removal, but, by removal of the bone flap and by some expansion of the cortex, the subdural space became quickly obliterated. This child gave no evidence of having had meningitis, although he was said to have had an upper respiratory infection a short time before admission. The advisability of removing well-formed membranes around subdural haematomas in infancy would appear to be re-emphasized by this experience.

The condition of Case 4 was very poor from the time of admission to hospital early in the illness. However, in view of the convulsions and lack of improvement under treatment, subdural taps should have been performed before the elapse of a week. In addition, bilateral burr-holes should have been made despite, on fontanelle punctures, the appearance of subdural fluid on the left side only.

Summary

Five cases of infected subdural effusion seen in young children in America are reported. Four were secondary to acute meningitis, the other the sequel to a traumatic subdural haematoma.

In infants, recovery from meningitis, otherwise adequately treated, may not occur because of the unsuspected development of subdural effusions. Their early recognition and treatment will also diminish the incidence of serious post-meningitic neurological defects.

It is our impression that more intensive antibiotic therapy than is often employed should be instituted in children with meningitis and that careful observation of the neurological state be made as often as possible so that subdural effusions may be recognized early.

BIBLIOGRAPHY

- Arnold, G. G. (1951). *J. Pediat.*, 39, 191.
 — (1952). *Ibid.*, 40, 757.
 Bloor, B. M., Grant, R. S., and Tabris, J. A. (1950). *J. Amer. med. Ass.*, 142, 241.
 Epstein, J. A., and Goldzier, S. E. (1953). *A.M.A. Arch. Neurol. Psychiat.*, 69, 242.
 Guthkelch, A. N. (1953). *British Medical Journal*, 1, 233.
 Jones, H. E. (1952). *Lancet*, 1, 891.
 McKay, R. J., jun., Ingraham, F. D., and Matson, D. D. (1953). *J. Amer. med. Ass.*, 152, 387.
 — Morissette, R. A., Ingraham, F. D., and Matson, D. D. (1950). *New Engl. J. Med.*, 242, 20.
 Mutch, N. (1941). *Lancet*, 2, 751.
 Smith, M. H. D., Dormont, R. E., and Prather, G. W. (1951). *Pediatrics*, 7, 34.
 Schwartz, A. B. (1916). *Amer. J. Dis. Child.*, 11, 23.
 Spitz, E., Pollak, A., and Angrist, A. (1945). *Arch. Neurol. Psychiat.* (Chicago), 53, 144.

By the end of this year the mileage of water mains in the country will be half as great again as it was when the war ended. Every year mains of a total length sufficient to reach from London to New York are being laid. These facts were given recently by Mr. ENOCH POWELL, Parliamentary Secretary of the Ministry of Housing and Local Government. "The number of households in the country without a piped water supply," he added, "has been halved in the last five years, and schemes before the Minister at the moment will go far towards halving the remainder. Even in the present period of financial stress rural water schemes are being approved at the rate of 6½ million pounds' worth a year, and nearly all attract Government grants under the Act of 1944."

SUBDURAL HAEMATOMATA AND EFFUSIONS IN INFANCY*

BY

GEORGE ORMISTON, M.D., F.R.C.P.Ed.

Consulting Paediatrician, Southampton and Winchester Group Hospitals

In discussing subdural effusions in infancy I can add nothing except my own limited experience to what has been said so well and so fully by Ingraham and Matson (1944, 1954) in the United States and by Guthkelch (1953) in England. There are few reports on the subject by British paediatricians, a fact which leads to the assumption that subdural effusions do not occur in Britain so often as on the other side of the Atlantic. This may seem to reflect credit on the conduct of our obstetrics in so far as birth injuries can be held responsible for such effusions; but first let us be sure that we are not sometimes missing the diagnosis. It is because of this possibility that I take leave to remind you of the features by which these intracerebral collections of fluid can be identified.

Aetiology and Pathology

Subdural haematoma resulting from birth trauma is due to the rupture of small vessels passing from the pia through the arachnoid and subdural spaces on their way to enter the venous sinuses, notably the sagittal sinus. It is likely that the decisive factor producing this rupture is unusual stress on the foetal head, leading to abnormal head-moulding and overriding of the cranial bones. However, one has seen extensive moulding and overriding without this damage, so that other influences such as hypoprothrombinaemia or immaturity of the endothelial tissues may be at work as well.

A remarkable thing is the extent of these effusions as seen at operation. Most often bilateral, and usually larger on one side than the other, the clot may cover the frontal lobe and extend down the temporal lobe to the floor of the middle fossa, and backwards a shorter distance over the parietal lobe. This does not necessarily correspond with the area covered by the original haemorrhage, which may be quite small. Ingraham and Matson (1949) concluded from the available experimental evidence that the blood becomes surrounded by a fibrinous wall which organizes and acts as a semi-permeable membrane, so that surrounding fluid passes into the sac so formed. As the blood disintegrates the oncotic pressure of the fluid rises and more fluid is drawn in. The increase in bulk of the encysted fluid leads to the rupture of the small capillaries in the wall of the sac, still more blood gets into the sac, and the cycle is repeated. Gitlin (1955) does not agree, however, that high oncotic pressure is the decisive mechanism leading to swelling of the sac, but presents data from which he postulates that the increase in the accumulation of fluid is an effusion through irritated and damaged capillary walls, and that repeated haemorrhages into the sac do not, in fact, occur.

From the practical point of view symptoms and signs arise when the growth of the cyst causes a crucial increase in intracranial tension, which happens rather suddenly between the third and eighth months of life, though it may be sooner or later. It is not clear why most cases do not reveal themselves in the early neonatal period. Some infants undoubtedly suffer small "silent" effusions after apparently uneventful deliveries, while others exhibit mild and transient symptoms, doubtless often put down to birth injury from cerebral anoxia and haemorrhage. In still others the signs are overt and respond completely to immediate treatment, as the following history relates.

*A lecture delivered at Nicosia on March 23, 1956, to the Cyprus Branch of the British Medical Association.

Case 1.—A female infant was seen at the age of 4 days on May 3, 1949. The mother, a primipara, was aged 40. The first stage of labour lasted 30 hours. Easy forceps delivery was carried out in the second stage. The infant appeared normal at birth. Within 48 hours she developed cyanotic attack, stiffness of limbs, exaggerated reflexes, and left-sided convulsions. The anterior fontanelle was tense and bulging. A subdural tap performed at right angles to a depth of 0.5 cm. yielded 30 ml. of dark red blood. Only a little blood-stained C.S.F. was obtained from the left angle. Bilateral subdural taps on subsequent days yielded minimal amounts of blood from the right angle. Lumbar puncture gave a fluid heavily blood-stained for five days, then xanthochromic, then clear. Fits ceased after the first subdural tap. At the age of 2½ years the child was perfectly normal in every respect.

Rarely, the haematoma is located in the posterior fossa rather than over the vertex, and will be missed unless the tap is repeated in this situation.

Case 2.—A male infant seen at the age of 8 hours after a difficult vertex delivery was dusky in appearance, was breathing irregularly, and was limp and unconscious. The fontanelle was slightly full ("boggy") and not tense. A lumbar puncture yielded bloody C.S.F. under moderate pressure. Bilateral subdural taps were negative. The baby died a few hours later, and at necropsy a large haematoma was found posteriorly under the tentorium.

Diagnosis

Ingraham and Matson's (1949) analysis of their large series gives all the information concerning age, sex incidence, nature and frequency of signs and symptoms, etc., that could be desired, and need only be summarized here. They show that males are affected twice as often as females and that most infants come under hospital care before the sixth month of life, the symptoms in the majority having been of recent origin—days to a few weeks. The symptoms are convulsions, vomiting, irritability, infection, stupor, and a history of trauma in diminishing order of frequency and in variable association. Similarly the signs are fever, hyperactive reflexes, bulging fontanelle, anaemia, enlargement of the head, abnormal fundi, paralysis, and fracture of the skull.

The diagnosis is clinched by the performance of a subdural tap after the hair has been shaved off the fontanelle and neighbourhood. The needle used should be short, about ½ inch (0.8 to 1 cm.) long. It is easily made from a Wassermann-size needle cut off the appropriate distance from its base with a bevel less acute and a point less sharp than the original. The body and limbs are firmly wrapped up ("mummified") and the infant is laid supine on a firm table, the nurse facing the operator, who should be comfortably seated on a chair opposite the vertex with a tray for the necessary instruments at his right hand. The nurse can easily keep the infant quite still if she places her forearms on each side of the trunk and holds the head between her hands. After the skin has been sterilized the needle is inserted at the lateral angle of the fontanelle. The skin and then the aponeurosis are pierced at right angles to the surface. The fluid should appear immediately, and can be led away through a short length of rubber tubing attached to the needle base by an adapter, or an all-glass syringe can be used to measure the fluid. It is wise to restrict the initial amount withdrawn to 10 to 20 ml. from each side, for the procedure must be done bilaterally. The puncture holes should be sealed with collodion.

Nature of Subdural Fluid

The character of the fluid varies according to the duration of the haematoma. It may be dark red soon after birth, but in a few weeks becomes dark brown or chocolate-coloured, and, later, amber.

Case 3.—A female infant, a first child, was born at home by difficult forceps delivery. She sucked well till the sixth day, when she became lifeless, then made good progress till the 25th day, when she had twitching limited to the right face, arm, and leg. Similar fits occurred on two subsequent days, with right conjugate

deviation of eyes, tremor, and deviation of tongue to same side. The skull appeared uniformly enlarged, circumference 15½ in. (38.7 cm.). Both fontanelles were much enlarged; the occipital suture was widened to admit two finger-tips. There was right facial paresis, all limbs were slightly spastic, and arm and leg reflexes were hyperactive. C.S.F.: xanthochromic; pressure 220 mm. Hg, subsiding to 150 when crying stopped; Nonne-Apelt reaction positive; protein, 65 mg. per 100 ml.; cells less than 1 per c.mm. From the left lateral angle 19 ml. of chocolate-coloured fluid was withdrawn by subdural tap, and a drop or two only of C.S.F. from the right. Taps on the following days were negative, except for a few drops of clear fluid from the left side. One week later Mr. J. Pennybacker excised a large membrane from the left hemisphere. It contained "muddy clot as usual." The haemoglobin fell to 45% post-operatively and a transfusion of blood was given. A slight squint persisted for a year, when the child was well and mentally normal and active, though perhaps not quite as clever with the right hand as the left.

Some Difficulties in Diagnosis

I propose to use other examples from the 10 patients that I have seen at two provincial hospitals in England to illustrate some of the difficulties that may arise in diagnosis.

Vomiting is an important symptom. It occurred in nearly 48% of Ingraham and Matson's (1949) cases and in at least 50% of Guthkelch's. It may occur alone and the underlying cause may be missed.

Case 4.—A male infant was admitted under my care at 6 months of age with drowsiness, an enlarged head, and a tense fontanelle. When a few weeks old he had been treated at another hospital because of severe vomiting—"cause unknown"—and discharged when the vomiting ceased. The symptoms remained in abeyance for several months. At operation (Mr. Pennybacker) the membranes were excised on both sides.

Case 5.—A male infant was admitted at 4 months with the complaint of "projectile vomiting, duration one day." He was said to have vomited once a day since birth; but there was only an occasional vomit in hospital. His weight was 13 lb. 6 oz. (6 kg.), and in the absence of signs and further symptoms he was soon discharged. He was readmitted a month later because of a convulsion, having continued to vomit once a day since discharge, though he had gained a further pound (0.5 kg.) in weight. He was now irritable, with a big head, tense fontanelle, slight stiffness of the limbs, and hyperactive reflexes. Repeated subdural taps were followed by the removal of a membrane from each hemisphere.

Finally, it is important not to be deceived by a negative subdural tap, for tragic consequences may result, as Case 6 shows.

Case 6.—A female infant was admitted to the gastro-enteritis unit at 6 months. She was a second child. Delivery was normal, birth weight 7 lb. (3.2 kg.). The parents had recently recovered from gastro-enteritis. The baby had vomited and been irritable with slight tremor in all limbs the day before admission. On the day of admission she had right-sided convulsions, recurrent small effortless vomits, a temperature of 104° F. (40° C.), no neck stiffness, and a "flat" fontanelle. C.S.F.: clear, with normal pressure; 2 cells per c.mm.; and a normal chemistry. Next day she had right hemiparalysis, left facial weakness, and tremor in the left arm. A subdural tap was negative. The tap was repeated the same day and a little blood obtained, thought to be traumatic. Illness continued, with high fever, coma, stiffness, and twitching. A subdural tap 12 days later yielded nothing on left, pure unclotted blood from right. Disks were normal; some fullness of retinal veins. A neurologist suggested a diagnosis of allergic encephalitis and advised cortisone. Six weeks after admission the temperature range was 100 to 101° F. (37.8 to 38.3° C.). The C.S.F. was again normal except for 100 mg. of protein per 100 ml. Death occurred 2½ months after admission. At necropsy there was a very large collection of xanthochromatous fluid on each side of the midline. Some old blood clot adhered to the under surface of the dura. The brain was grossly flattened, and there was a communicating internal hydrocephalus. The subdural fluid protein content was 1,000 mg. per 100 ml.

The failure of the first subdural tap may have been due to blood clot obstructing the needle, which may not have been pushed far enough. The unclotted blood withdrawn at the second tapping may have come from a vessel or from the cyst as a result of the intermittent bleeding postulated by Ingraham and Matson.

Subdural Effusions Complicating Meningitis

Although I can give only two examples of effusion complicating meningitis, I feel that in past years, before certain antibiotics were obtainable in England and before treatment of the meningitides had been accurately worked out, I must have had others.

Case 7.—A female neonate was treated at a maternity hospital for 16 days for what was undoubtedly a septicaemia with meningitis, organism undiscovered. She was transferred to a children's hospital because of convulsions. The urine contained pus and *E. coli* and the fontanelle was bulging. C.S.F.: 32 cells per c.mm., mostly lymphocytes; protein, 130 mg. per 100 ml.; globulin, slight excess; chlorides, 635 mg. per 100 ml.; sugar, 35 mg. per 100 ml. A bilateral subdural tap yielded an amber-coloured fluid containing a protein concentration of 1,000 mg. per 100 ml. Subdural taps were repeated on the following days, and antibiotics were continued. The infant was discharged a month after admission, and at 7½ months of age was sitting up unsupported, appearing normal in every way.

Case 8.—A male infant was transferred to my care at 3 months of age because of vomiting. He had been under treatment with various antibiotics for "acute pneumonitis." The head was retracted, neck stiff, fontanelle full, and the disks were normal. C.S.F.: 38 cells per c.mm.; 160 mg. protein, 710 mg. chloride, and 37 mg. sugar per 100 ml. From 10 to 40 ml. of yellow fluid was withdrawn from each side by subdural tap on 16 successive days—a total of 563 ml. of subdural fluid. Mr. Pennybacker subsequently operated and removed a membrane from the left hemisphere.

Treatment

Specific treatment is primarily the concern of the paediatrician, who should withdraw the subdural fluid on each side daily until it becomes difficult to obtain. This is really a method of producing gradual decompression, and may take anything from a few days to a few weeks. An antibiotic such as procaine penicillin 300,000 units or oxytetracycline 20 mg. per lb. (40 mg. per kg.) body weight should be given daily as a prophylactic or to clear up infection in the respiratory passages or elsewhere. A considerable amount of protein is lost in the subdural fluid, and this should be made good by one or more transfusions of citrated blood. Some infants show an increasing anaemia, so that transfusions of blood serve a double purpose. The nutrition is as often as not below average, sometimes much below, and good feeding during the pre-operative period enhances the prospects of successful operation.

When the infant is transferred to the care of the surgeon the presence of solid clot or membrane is determined by inspection after making burr-holes in the temporal region in the first place, and elsewhere if necessary. If clot and membranes are found, bone flaps are turned down and the tough outer membrane is attached to the dura, and the thin membrane and clot over the pia are removed on one side and, at a later date, on the other.

As has been seen, subdural haematomata that are discovered within a few days of birth can be successfully treated by subdural taps without resort to operation.

Subdural effusions as a complication of meningitis undoubtedly occur more often than has been realized, particularly after infection by *H. influenzae*—not only one of the most frequent infections in early infancy but also one of the most difficult to treat—and also after pneumococcal and meningococcal infections. Smith (1953) states that subdural taps alone seem to constitute adequate local therapy in these effusions, which generally clear up within three weeks. On the other hand, McKay and his colleagues (1953) achieved success by subdural tap in only 15 out of 50 cases, for the remaining 35 needed craniotomy as well.

One may conclude that small sterile effusions often arise in the course of treated meningitis and clear up spontaneously. Effusions which become clinically recognizable may be sterile or infected, and clear up in response to subdural taps and continued antibiotic therapy. If definite membrane formation has taken place and the contents of the sac are infected, subdural taps alone are unlikely to effect cure, and

craniotomy will be necessary. Subdural taps should be initiated when adequate antibiotic treatment of the infection does not bring down the fever, sterilize the spinal fluid, and abolish symptoms in three or four days.

Discussion

The detection of subdural effusions may fall to the lot of all who have to do with the medical care of young infants—the obstetrician, the paediatrician, and the general practitioner—whether in hospital or in the home. As has been seen, certain signs and symptoms found in such effusions are common to a number of unrelated conditions. Vomiting, anaemia, and wasting, and even convulsions in early infancy are so much more related to digestive disturbances and infection that the real nature of the trouble may escape recognition unless careful examination of the nervous system is a repeated routine. A simple and neglected procedure is the measurement of the circumference of the heads of young infants from time to time as a means of estimating normal and abnormal growth.

A history of a normal or non-instrumental labour should not put the physician off guard, for the real quality of the labour is often unknown unless good hospital records are kept; and, in any case, birth injuries occur in deliveries that have been apparently normal. It must be remembered that post-natal injuries, or a knock on the head, can cause subdural effusions of blood, though the history of such injury is always more forthcoming, and evidence of fracture in the skull x-ray film is more often associated with it, than with injuries incurred at birth.

Conclusion

Subdural haematomata and inflammatory effusions are best prevented by good obstetrics, and by early diagnosis and correct antibiotic therapy. When they do occur, early recognition and appropriate treatment should reduce to the minimum the incidence of sequelae such as paralysis, spasticity, mental defect, and death. Ingraham and Matson (1954) are able to state authoritatively after 14 years' follow-up that 70 to 80% of the children who had been surgically treated appear to be developing mentally for their age, and that residual neurological signs and post-operative convulsions are uncommon. The retarded and deficient children have been for the most part those with large chronic haematomas with marked atrophy of the hemispheres well established when the membranes were removed.

REFERENCES

- Gitlin, D. (1955). *Pediatrics*, 16, 345.
Guthkelch, A. N. (1953). *British Medical Journal*, 1, 233.
Ingraham, F. D., and Matson, D. D. (1944). *J. Pediat.*, 24, 1.
— (1949). *Advanc. Pediat.*, 4, 231.
— (1954). *Neurosurgery of Infancy and Childhood*. Thomas, Springfield, Illinois.
McKay, R. J., Ingraham, F. D., and Matson, D. D. (1953). *J. Amer. med. Ass.*, 152, 387.
Smith, M. H. D. (1953). *Bull. Tulane Med. Fac.*, 12, 105.

In 1955 there were 759,571 donations of blood, compared with 700,202 in 1954. This is the largest number recorded. 116,410 people gave their blood for the first time to the National Blood Transfusion Service last year, and at the end of December the total strength of the donors' panel was 591,204, compared with 540,389 a year before. The National Blood Transfusion Service needs at least 650,000 regular donors to make sure that adequate supplies of blood are available to meet the needs of the hospitals, to replace the older donors when they reach the age of 65, and to provide adequate reserves of blood plasma. Anyone who is in normal health and between the ages of 18 and 65 is welcomed at the donor sessions, which are held frequently in convenient places throughout England and Wales.