which no warning had been given from a preliminary test dose injected into the pulmonary artery. It is concluded that the administration of a test dose of contrast medium is an unreliable method of detecting hypersensitivity, and may in fact induce a false sense of security.

The treatment of the condition presented a difficult problem. As might have been expected, the methods of treatment applicable to pulmonary oedema occurring in left ventricular failure were ineffective here. Intravenous hydrocortisone, on the other hand, appeared to exert a specific and prolonged anti-allergic action in reducing pulmonary oedema. The antihistamine drugs proved of little value. Nikethamide was useful in reviving consciousness and the cough reflex so that fluid could be expectorated.

Intravenous hydrocortisone induced immediate improvement on three separate occasions, and undoubtedly saved this patient's life. Intramuscular cortisone, which was given initially, was presumably ineffective owing to poor absorption in the presence of severe peripheral vasoconstriction. The use of intravenous hydrocortisone in overwhelming allergic reactions is a rational therapeutic measure, and the present case history confirms its usefulness. Our experience indicates that the drug should be immediately at hand in all centres employing the intravenous injection of contrast media.

Summary

A case of gross pulmonary oedema following selective angiocardiology in a patient with primary or chronic thrombo-embolic pulmonary hypertension is reported. Recovery was clearly attributable to prolonged intravenous infusion of hydrocortisone. A test dose of sodium acetrizate failed to provide warning of hypersensitivity.

REFERENCES

- Dotter, C. T., and Jackson, F. S. (1950). *Radiology*, **54**, 527.
McAfee, J. G. (1955). *Amer. J. med. Sci.*, **229**, 549.
Morgan, R. H. (1950). *Amer. J. Roentgenol.*, **64**, 189.

FREQUENCY IN THE FEMALE DUE TO URETHRO-TRIGONITIS CAUSED BY INTESTINAL INFECTION

BY

J. E. SEMPLE, F.R.C.S.

Surgeon to St. Paul's Hospital, London

In his Hunterian Lecture of 1933 Winsbury-White proved by means of injected animal specimens that infection of the genital organs in the female caused marked pathological changes in the urethra and trigone of the bladder. This fact had been readily accepted as occurring in the male as a result of the anatomical relation of the prostate to the bladder neck, but it was not recognized as occurring so readily in the female. By the routine use of a urethroscope, characteristic changes at the bladder neck were found to be invariably associated with disturbances of micturition.

This is the original work which established urethro-trigonitis as a clinical entity and the commonest condition found in female patients attending a urological clinic, but, as Winsbury-White (1947) states, chronic urethro-trigonal changes are not uncommon in women with healthy cervixes. And it is not uncommon even in patients who had had the genital organs removed some time prior to the onset of the urethro-trigonitis.

General Review of Urethro-trigonitis in the Female

In a review of female patients suffering from urethro-trigonitis and attending a urological clinic over the last seven

years it was found that the incidence of this condition was remarkably constant from year to year, varying only between 25 and 30% of the total female attendances of 453. The majority of cases occurred between the ages of 30 and 50, but it might occur in any of the age groups. The most common symptom was frequency of micturition, which was pronounced in 80% of the patients. Dysuria was present in 50%, as was urgency of micturition. In spite of the old teaching that residual urine is rare in the female, an intravenous pyelogram in all cases revealed that 66% of these cases had a definite volume of residual urine left after micturition. The laboratory reported that 16% of the catheter specimens taken as a routine contained pus cells and that organisms were present in only 12%, the predominant organism being *Bact. coli*. Winsbury-White states that urethro-trigonitis is a predisposing cause of *Bact. coli* infection of the urinary tract, and that it does not produce such an infection by itself but only by causing conditions which favour the onset of infection by reason of obstruction of the bladder neck. The diagnosis was made by routine examination with a urethroscope, and characteristic changes in the urethra and bladder neck were noted in every case. The cervix was examined as a routine, and an erosion or evidence of cervicitis was present in 66% of cases; of the remainder there was no evidence of genital infection.

In each case the internal sphincter was dilated as a result of the urethroscopic examination, and dilatation was repeated in accordance with Winsbury-White's method at not less than six-weekly intervals and to a dilatation not greater than F.28. In 25% treatment was speeded up by fulguration of the polypi and granulations and hillocks: 66% were cured and 28% much improved.

Urethro-trigonitis Not due to Infection of Genital Tract

There remain, therefore, 34% of cases in which no infection of the cervix or genital organs can be found to account for the pathological changes in the urethra and trigone.

In a number of patients an association between the large intestine and the neck of the bladder had been noticed: the first instance was that of a patient in whom an acute attack of diverticulitis of the sigmoid colon followed each dilatation of the urethra and in whom marked urethro-trigonal changes had been found on urethroscopy. On further inquiry the patient stated that when she had an attack of diverticulitis the symptoms of urethro-trigonitis were very much worse.

A frequent complaint in these patients is of discomfort and even pain in the left iliac fossa, which may be associated with some irregularity of the bowels. On abdominal palpation there is not infrequently definite tenderness over the sigmoid and pelvic colon, which may in some patients be palpable.

Out of 43 cases examined radiologically by means of a barium enema, 11 showed lesions of the large intestine which were closely related to symptoms of urethro-trigonitis verified by urethroscopy. Of the 11 cases, 8 showed numerous diverticula of the lower colon with radiographic appearances strongly suggesting inflammatory changes in the surrounding bowel. Two patients had been previously diagnosed as having diverticulitis with rigidity and thickening of the sigmoid colon. In one case a barium enema showed a carcinoma of the recto-sigmoid junction though there had been no initial symptoms suggesting a neoplasm of the bowel but only those of urethro-trigonitis.

Chronic constipation was not accepted as a likely cause of urethro-trigonitis, as it does not of itself produce inflammatory changes in the wall of the bowel; nor were lesions of the anal canal included as a cause, although on anatomical grounds there would seem to be no reason why they might not form a source of infection.

In the absence of animal experiments, which made Winsbury-White's work so conclusive, a study of the relationship of the lymphatics of the lower part of the large intestine with those surrounding the bladder neck, although

not showing such a close relationship as that of the cervix to the floor of the bladder, nevertheless shows a ready lymphatic connexion. This is well shown by Rouvière (1932), and there would seem to be no reason why infection should not spread from the lymphatics of one organ to those of the other, which would appear, on clinical grounds, to be the case.

The results of treatment of the bowel condition combined with that of the bladder neck go far to support this theory. Dilatation of the bladder neck and even fulguration of polypi and granulations have not produced very encouraging results in such cases, but when the lesion of the bowel is alleviated the urethro-trigonitis responds to treatment very much more readily. Thus the bladder neck was treated in exactly the same way as when the infection came from the cervix, care being taken not to irritate the urethra by too frequent dilatation—that is, not more than once every six weeks and usually at less frequent intervals.

Treatment of the bowel depends on the type of lesion and its extent. In the less extensive lesions, in which surgery is not called for, the institution of a low residual diet which does not irritate the mucosa of the bowel and limits fermentation, combined with regulation of bowel action by "petrolagar" with or without phenolphthalein, alone may give much improved results, but infection can be further limited by the administration of phthalylsulphathiazole in an initial dose of six 0.5-g. tablets followed by two tablets every six hours for fourteen days. This course can be repeated at fortnightly intervals until the infection has been greatly diminished. (In the more resistant cases an initial course of streptomycin and sulphaguanidine followed by phthalylsulphathiazole may be necessary.) As a result of this routine the symptoms of urethro-trigonitis were improved in each of the cases treated.

Summary

Symptoms of urethro-trigonitis were present in 25 to 30% of female patients attending an out-patient clinic, and this condition was verified by urethroscopy in which characteristic changes were found present at the bladder neck. These changes have been proved to arise from lesions of the cervix and genital tract, and were found in 66% of these cases. Of the remaining 34%, a third were found to have radiologically demonstrable lesions of the large intestine. Treatment of the intestinal infection, combined with dilatation and, where necessary, fulguration of the bladder neck, gives excellent results as compared with dilatation and fulguration alone.

REFERENCES

- Rouvière, H. (1932). *Anatomie des Lymphatiques de l'Homme*, pp. 340, 397, and 413. Masson, Paris.
Winsbury-White, H. P. (1933). *Brit. J. Urol.*, 5, 249.
— (1947). *Med. Wld (Lond.)*, 66, 170.

The Ministry of Education on September 13 asked local education authorities to review their arrangements for providing primary and secondary education for children in hospital and to see that they are as comprehensive as possible. Though much has been done, the Ministry claims that "there are still some educable children in hospital for whom no provision has been made." This may be provided either by teachers visiting or by setting up hospital schools. In 1955 there were 120 hospital schools in England and Wales, with 6,476 children on the roll; 1,425 other children were receiving individual tuition. Where there is no hospital school or regular teaching group, the Ministry of Health has asked hospitals to inform education authorities about the admission of any child likely to remain in hospital who is yet well enough to have some lessons. The circular calls for a good supply of library books, commenting: "Reading matter given by well-intentioned visitors may not always be suitable or adequate."

ACCIDENTAL CARBON MONOXIDE POISONING

DANGERS OF INADEQUATE VENTILATION

BY

F. S. FIDDES, M.D.

Department of Forensic Medicine, University of Edinburgh

Of late there have been frequent expressions of concern at the increasing number of deaths from accidental coal-gas poisoning. A high proportion of such accidents are associated with various predisposing or contributory conditions in the victim, which have been described collectively as "the 4 D's of accidental carbon monoxide poisoning"—the decrepit, the diseased, the drugged, and the drunk. Others are caused by faulty appliances or by ill-advised amateur adaptations, and occasionally responsibility can be traced to defects in gas-pipes or joints which are actually outside the premises occupied by the victims.

The danger of coke braziers and coke-burning stoves is well known, but deaths still occur from these sources, through either ignorance or disregard of warnings. It is insufficiently appreciated, however, that even coal-burning stoves and fires may be dangerous in certain circumstances, and that gas-fires and other appliances, though free of defects in themselves, may also be a source of danger if the conditions do not ensure free and complete combustion of the coal-gas. The following cases, resulting in the deaths of three healthy adult males, illustrate the circumstances in which fatal poisoning from such sources may arise, and point to the need for a greater awareness of the potential danger.

Case 1

A man aged 50, employed as watchman on a building site, was in the way of spending part of the night asleep in a workmen's hut, and one morning he was found lying dead on the bunk with which the hut was fitted. The post-mortem appearances, external and internal, pointed to death from carbon monoxide poisoning, and the blood contained 55% carboxyhaemoglobin. There was no other significant finding at necropsy, apart from recent pleural adhesions over the left lung and older adhesions over the right lung. The man had recently resumed work after several weeks in hospital suffering from pneumonia.

The hut was of the wheeled type, adapted for towing, and the door at the rear was divided horizontally into two parts, upper and lower. There were two small windows, each capable of being opened. Inside the hut there was a fixed stove, with a chimney leading out through the roof. The capacity of the interior of the hut was approximately 600 cubic feet (17 cubic metres). On the night in question, which was very cold with a gusty wind blowing, both halves of the door had been closed and only one of the windows was slightly open. A fire had been burning in the stove, and this had apparently been well stoked before the man lay down to sleep. There was also an oil-lamp burning.

It was quite certain that no coke had been used in the stove, the only fuel available being coal of normal household quality. The stove and chimney were in good condition, without obvious defects. In these circumstances it was thought desirable to confirm the possibility of a lethal concentration of carbon monoxide having accumulated in the hut, and with this in view a fire of the same coal was lit in the stove and allowed to burn for about one and a half hours. The stove had previously been cleared of ashes and the fire burned freely. One window was slightly open. Samples were then taken of the flue (chimney) gases, with