Conclusion

Although the importance of a synecological or holistic approach is stressed, it would be a mistake to take this too literally. Ecological processes are so complicated that there is a danger of getting involved in a mass of irrelevant ecological detail, and of forgetting that when studying a zoonosis the natural history of the causative organism is our chief concern. The infecting organism is in fact the pivot round which the ecological investigation revolves.

If the views expressed here have any validity, they probably apply not only to the zoonoses but to parasitic infections in general.

Summary

The zoonoses, which are defined as *infections of man naturally acquired from other vertebrates*, are treated as a problem in ecology. This entails studying the interrelation between man, animals, a causative organism, the environment, and sometimes arthropods. Such an approach is holistic or synecological—wholes being regarded as more important than parts. Holism is a dynamic not a static conception.

The evolution of the zoonoses is discussed, particularly in relation to plague, relapsing fever, and leishmaniasis. The most important reservoirs of zoonoses and other parasitic infections are usually resistant or relatively insusceptible animals rather than highly susceptible ones. Plague and Chagas's disease illustrate this. Resistant and unduly susceptible animals occupy different "niches" in nature. By "niche" is meant the place of an animal in its biotic environment. Zoonoses often have a "focal distribution" in nature. Thus wild rodents infected with plague may occur in "pockets," and the vector mites of scrub typhus congregate in "typhus islands."

An unstable environment often favours the transference of zoonoses to man and animals. Examples are the plague-infected plain of Rongai, in Kenya, the typhus-infected forest clearings in Malaya, and the yellow-fever infected forest edges in Uganda. Ecologically unstable areas are termed "ecotones." Zoonoses can also be transmitted in or near relatively stable sites such as huts, rodent burrows, caves, and termite hills. These are known as "habitat niches." Animals and arthropods in a community are linked by food, and the importance of what are known as "food chains" and "food cycles" is discussed in relation to the transference of zoonotic infections from one host to another.

Reference is also made to the fluctuations in numbers of various animals in nature, and how this affects the incidence of zoonotic disease. Certain highly susceptible rodents are periodically decimated by plague; this breaks the link with man, and human infections cease for the time being.

I wish to thank the Director of Medical Services, Kenya, for permission to publish this paper.

REFERENCES

- Audy, J. R. (1954). *British Medical Journal*, **1**, 960.
 — and Harrison, J. L. (1951). *Trans. roy. Soc. trop. Med. Hyg.*, **44**, 371.
 Baltazard, M., Mofidi, C., and Bahmanyar, M. (1947). *Ann. Inst. Pasteur*, **73**, 1066.
 Boiron, H., Koerber, R., and Carronier, B. (1948). *Bull. Soc. Path. exot.*, **41**, 81.
 Elton, C. (1925). *J. Hyg. (Camb.)*, **24**, 138.
 — (1927). *Animal Ecology*. London.
 Garnham, P. C. C. (1952). *Trans. roy. Soc. trop. Med. Hyg.*, **46**, 397.
 Heisch, R. B. (1949). *British Medical Journal*, **1**, 17.
 — (1950). *Ann. trop. Med. Parasit.*, **44**, 260.
 — and Garnham, P. C. C. (1948). *Parasitology*, **38**, 247.
 — Grainger, W. E., and D'Souza, J. St. A. M. (1953). *Trans. roy. Soc. trop. Med. Hyg.*, **47**, 503.
 Meyer, K. F. (1942). *Medicine (Baltimore)*, **21**, 143.
 Nicolle, C., and Anderson, C. (1927). *Arch. Inst. Pasteur, Tunis*, **16**, 123.
 Smuts, J. C. (1936). *Holism and Evolution*. London.
 Wenyon, C. M. (1926). *Protozoology*. London.

CHEMICAL AND CLINICAL PROBLEMS OF THE ADRENAL CORTEX*

BY

F. T. G. PRUNTY, M.D., F.R.C.P.

Professor of Chemical Pathology in the University of London; Physician, St. Thomas's Hospital

Some Factors Governing the Response to Corticotrophin

Growth Hormone

In 1951 Stack-Dunne and Young stated that they were able to obtain preparations from the anterior pituitary having two distinct kinds of activity. These separate preparations produced either a fall in the adrenal ascorbic-acid content or an increase in adrenal weight. They used growth hormone as a source of the adrenal weight factor, but did not consider that this factor was necessarily growth hormone itself. There seemed to be clinical data supporting the view at the same time, for at the same meeting to which Stack-Dunne and Young gave their paper we presented our early data on discrepancies between human response and the assay potency as determined by the Sayers method. However, as we have seen, this seems to have been explained on grounds of destruction or inactivation of the corticotrophin in the patients receiving it. The substance promoting adrenal weight and associated with, but not necessarily identical with, growth hormone also causes increased mitotic activity in the adrenal cortex and differs from the ascorbic-acid-depleting substance which is probably responsible for release and synthesis of corticosteroids (Cater and Stack-Dunne, 1955).

Forsham *et al.* (1955) summarized some experiments using a substance prepared by Rinfret from pig pituitary which was not itself steroidogenic but which, when given to a patient subsequently given corticotrophin, augmented the urine steroid response measured by the Porter-Silber method. A commercial growth-hormone preparation did not have a similar effect. We were fortunate enough to have at our disposal a small quantity of one of Young's growth-hormone preparations which he thought would be active with respect to the adrenal weight factor, although it had not been specifically tested for this activity. This substance was administered to a patient with Simmonds's disease whose adrenal had already been and continued to be stimulated with corticotrophin. We judged that in these circumstances the patient would also be deficient in growth hormone or the adrenal weight factor, and would be quite sensitive to it. There was a rapid increase in blood-sugar level, presumably due to the growth hormone itself, given intramuscularly 25 mg. twice daily for five days, but no increase in the ketogenic steroid or ketosteroid excretion. This negative result cannot be construed to be entirely contradictory to the hypothesis presented. It may well be that too little material was given for too short a time, or, as we shall see, that the absence of some other factor was limiting the response.

There is a clinical example before us worthy of attention with regard to this matter—namely, acromegaly. It is a commonplace that the adrenal cortices of patients with acromegaly are large and often contain subcapsular nodules. There is no unanimity of opinion about their

*Conclusion of the Humphry Davy Rolleston Lectures delivered at the Royal College of Physicians of London on May 15 and 17. The first part was published last week.

steroid production, but this might be expected to be low if the acidophil tumour caused sufficient reduction of anterior pituitary tissue. Adrenocortical insufficiency appears to be rare in acromegaly (Rummell, Baer, Hollister, and Kolb, 1951). In six patients with acromegaly the ketosteroid excretion was normal in three and low in three (Fraser *et al.*, 1941). In another group, three women and three men had high normal values, whilst one male with very prolonged disease had low values (Prunty, 1944).

Three cases have recently been studied. The first was a Polish grave-digger aged 36 with gross acromegaly. He excreted up to 22.5 mg. of ketosteroid daily and 18.4 mg. of ketogenic steroid. The second case was that of a man aged 43 who for five years had acromegaly with diabetes. He excreted up to 17.9 mg. of ketosteroid daily and 17.8 mg. of ketogenic steroids. The third patient was a woman aged 39 who 15 years previously had had an acidophil adenoma removed for eye signs. She had lately become very hirsute, with severe acne and pubic hair of male type. She excreted up to 34 mg. of ketosteroid and 12 mg. of ketogenic steroids daily. The last two patients were given corticotrophin according to the standard test. This was unfortunately of poor potency and had given apparently subnormal responses in several patients (see Table II). It had been used, however, in a patient with Cushing's syndrome due to hyperplasia of the adrenal cortex, with strongly positive results (S. E., Fig. 17). Such patients have a high sensitivity to corticotrophin. The two acromegalics gave a ketogenic steroid response which was low normal. The first acromegalic received a known potent batch of corticotrophin (89754) and did not show a brisk response. Clinically all these patients could be regarded as being in the active stage of the disease; although they presumably secreted excessive amounts of growth hormone, it does not follow that they did likewise with some other adrenal-cortical-activating agent. The results merely suggest that growth hormone itself did not appear to be a potent potentiating substance under these conditions of study.

Gonadotrophins

A second substance which has been implicated in having a subordinate effect on the adrenal cortex is the luteinizing hormone. Albright, Smith, and Fraser (1942) advanced complex arguments, based on the clinical relation between ovarian function and adrenal cortical function, that luteinizing hormone exerted an effect on ketosteroid secretion by the adrenal. Experiments based upon the suppression of ketosteroid excretion, which were thought to cause a simultaneous depression of luteinizing hormone and adrenal secretion of ketosteroid, seemed to support this hypothesis (Reifenstein, Forbes, Albright, Donaldson, and Carroll, 1945). We have conducted several experiments using preparations of luteinizing hormone ("pregnyn"; chorionic gonadotrophin) and follicle-stimulating hormone ("gestyl," prepared from pregnant mare's serum; serum gonadotrophin). The first of these was tried in the patient with Simmonds's disease who failed to show an effect with the growth-hormone preparation. No significant changes occurred when, in addition to corticotrophin, 1,000 units of gestyl was given daily for 10 days and on the fifth day 1,000 units of pregnyn daily was started for 10 days. This was in spite of the fact that a small menstrual period occurred, the first for many years.

In a second experiment a seemingly positive result was obtained. A patient with Hodgkin's disease was

under successful treatment with corticotrophin; on 20 "clinical units" daily he excreted 34 mg. of ketogenic steroid and 24 mg. of ketosteroid. The addition of 3,000 units of pregnyn daily (chorionic gonadotrophin) caused, on the fifth and sixth days, a rise in ketogenic steroids to 60 mg. and ketosteroids to 33 mg. There was a prompt fall in the steroid levels after withdrawing gestyl.

A third experiment with pregnyn in a patient with rheumatoid arthritis treated with corticotrophin was negative. Two other experiments have also been made with gestyl (from pregnant mare's serum) with negative results; if anything there was a fall in the ketogenic steroid excretion.

These results are difficult to interpret, and, even if the positive finding can be repeated, it should be noted that both ketogenic steroids and ketosteroids responded to chorionic gonadotrophin, which does not correspond with the theory of Albright. There remains little doubt that there is a complex reciprocal relation between the adrenal cortex and the gonads (Parkes, 1945), and studies in animals suggest that the ovaries may sometimes even replace the adrenal cortex (Hill, 1948). In guinea-pigs gonadotrophin has been found to suppress the response of the adrenal to corticotrophin, using the excretion of ketogenic steroids as a criterion (Clayton and Hammant, 1955b).

Effect of Tri-iodothyronine on Response to Corticotrophin

It has been thought for many years that there is an interdependence of the thyroid and adrenal cortex (Marine, 1930). Claims have been made that cortisone and corticotrophin reduce thyroid function (Hill, Reiss, Forsham, and Thorn, 1950; Wolfson, Beierwaltes, Robinson, Duff, Jones, Knorpp, Siemienski, and Eya, 1951; Lerman, 1955a). In the rabbit, radioactive iodine release by the thyroid is retarded by cortisone (Harris, 1955a). The low ketosteroid excretion in myxoedema has been found to rise when replacement therapy is given (Beierwaltes and Bishop, 1954; Lerman, 1955b). In our experience thyroid therapy has been required to produce significant change in patients with myxoedema secondary to anterior hypopituitarism.

We have been able to conduct a single observation on the effect of tri-iodothyronine on the activity of the adrenal cortex. The same patient with Simmonds's disease was again under treatment with corticotrophin (Fig. 6), and tri-iodothyronine was given intravenously in a single dose of 0.5 mg. and two days later 0.25 mg. was injected. The typical effects of tri-iodothyronine are shown on the B.M.R. and the serum creatine and blood cholesterol levels. After an interval of two days the urine steroids and plasma hydrocortisone levels rose significantly. There were also effects on the level of blood sugar and on sodium retention. It can be concluded from this experiment that conditions exist in which increased thyroid activity can increase adrenal cortical secretion. The likely interpretation seems to be that when a near maximal stimulus by corticotrophin was applied the response was limited by a degree of hypothyroidism, the correction of which produced a good normal response. This effect might not so easily occur in patients treated for myxoedema, since in this situation, although thyroid hormone is raised to adequate levels for response of the adrenal cortex, the rate of endogenous secretion of corticotrophin could be the limiting factor.

Development of Resistance to Corticotrophin

Earlier, when inactivation in muscle or subcutaneous tissue was a common problem, the development of resistance to continued injection also occurred (Forsham, Renold, and Frawley, 1951). It has already been indicated how variable

the response of patients can be to different doses and batches of corticotrophin, and for this reason some direct measurement of corticotrophin activity seems essential before in clinical practice the conclusion is drawn that a patient does not show a symptomatic response to treatment.

This point has been stressed by West (1954) and is illustrated by J.J., a woman under treatment with corticotrophin for rheumatoid arthritis (Fig. 7). After a prolonged period of therapy she no longer responded well clinically. It was known that the batch of corticotrophin in use on day 88 was not very effective in other individuals, so it was changed to a known very potent one (89754) at the same dosage (10 "clinical units" twice daily). As the response at day 109 was poor we repeated the identical batch and dose given at the beginning—that is, corticotrophin (89754) in doses of 20 "clinical units" twice daily. The patient responded again, but to a smaller degree. Nevertheless the intravenous administration of 40 "clinical units" of the same material on day 118 produced a large response, the excretion rate reaching 6.1 mg. per hour. It must be concluded that this patient developed a partial degree of unresponsiveness which could easily be overcome with sufficient stimulus. This experience demonstrates another reason for avoiding assumptions about the dose-response relationship to corticotrophin. West (1956) has also observed under certain conditions a failure of adrenal response to long-acting corticotrophin which is capable of being overcome.

Responses to Operation

The first observations on the effect of trauma in man on adrenal cortical secretion were made by Venning (1945), who found by bioassay an elevation of cortin levels of urine extracts under these conditions. For this and other reasons it may be supposed that operative intervention is a good method of investigating a patient's response to endogenous corticotrophin and possibly of studying quantitative differences in the response from those obtained with exogenous corticotrophin. Post-operatively, the general pattern of events seems to be that there is often a short initial increase of ketosteroids, sometimes followed by a fall for several days (Forbes, Donaldson, Reifstein, and Albright, 1947; Prunty, 1953). When the Porter-Silber method was used the corticoid metabolites were found to

FIG. 6.—Effect of tri-iodothyronine on response to corticotrophin in a patient with Simmonds's disease. Tri-iodothyronine, two doses, was given intravenously. Eosinophils marked ●; plasma hydrocortisone ▲.

FIG. 7.—Long-term response of patient with rheumatoid arthritis to three batches and varying doses of intramuscular corticotrophin (except where marked, i.v.). The numbers on the top line, above the dosages, refer to different batches of corticotrophin.

increase early after operation and to persist for several days, depending on the amount of trauma (Moore, Steenburg, Ball, Wilson, and Myrden, 1955); this also occurs after burns (Wilson, Lovelace, and Hardy, 1955). With the method for ketogenic steroids the response is similar (Fig. 8). This has also been found by Jepson, Jordan, and Levell (1956). The magnitude of the response varies in different patients, but there is a tendency for the corticoid metabolites to increase to a greater extent than the ketosteroids. When methods of blood analysis for corticoids were used a sharp rise was also found immediately after operation, which may to a small extent be enhanced by temporary inability of the liver to metabolize the hormones as rapidly as normally (Sandberg, Eik-Nes, Samuels, and Tyler, 1954).

Adrenal Cortical Virilism and Hirsutism

In 1933 Broster and Vines classified patients with the adrenogenital syndrome into those occurring before puberty and those after puberty. The rise in urine ketosteroids is a much more pronounced feature in the former group, and the ketosteroid levels in these patients are often very high after the adrenarche at puberty (Mason and Engstrom, 1950). In the latter group ketosteroids may apparently be normal, high normal, or just above normal, which is seen to be the case in the group of patients described below. Recently Patterson has found that the pre-pubertal group is divisible into two subgroups according to a high or low level of dehydroepiandrosterone excretion (Broster, Patterson, and Camber, 1953). There seems to be no obvious clinical significance of this finding at the present time. It has been observed that even patients with virilizing adrenocortical tumours, who sometimes have high excretion rates of dehydroepiandrosterone, need not necessarily excrete large amounts of this substance (Mason and Engstrom, 1950; Prunty, 1950). Its excretion will depend on its further conversion by the adrenal and rate of metabolism after secretion.

F.F. 58 ♂

FIG. 8.—Response of "normal male" to lobectomy for bronchial carcinoma. Subsequently he responded normally to a poorly potent batch of corticotrophin.

FIG. 9.—Failure of corticoid synthesis in pre-pubertal virilism with excessive excretion of pregnanetriol and 11-ketopregnanetriol.

The greatest biochemical advances in the field of the pre-pubertal syndrome have been made by Wilkins and his associates, particularly Bongiovanni (Wilkins, Bongiovanni, Clayton, Grumbach, and van Wyk, 1955). Bartter, Albright,

FIG. 10.—Pre-pubertal virilism. Effect of corticotrophin, aldosterone, and cortisone. Note the determination of ketogenic steroids includes pregnanetriol, therefore corticoid metabolites measured by Reddy method (P-S). The latter rises on cortisone therapy due to metabolites derived from it.

Forbes, Leaf, Dempsey, and Carroll (1951) were the first to suggest that there is basically a deficiency in adrenal corticoid synthesis in this condition, and this would theoretically lead to an increased circulation of corticotrophin. The latter has been demonstrated by Morris (1952) and by Sydnor, Kelley, Raile, Ely, and Sayers (1953). The deficiency in synthesis of hydrocortisone and other corticoids is partially due to failure of introduction of the hydroxyl group at position 21 (Dorfman, 1955). A high proportion of ketosteroid is often found with hydroxylation at position 11. Bongiovanni has also found that pregnanetriol is excreted in the urine in large amounts which, it seems likely, originate from 17-hydroxyprogesterone, as also probably does the increased excretion of 11-ketopregnanetriol reported by Finkelstein, von Euw, and Reichstein (1953) (Fig. 9). It is thought that the high secretion rate of corticotrophin stimulates the secretion of excessive adrenal androgen, a raised blood level of which, unlike that of hydrocortisone, does not inhibit further corticotrophin release.

These patients vary in their response to exogenous corticotrophin, but the response shown in Fig. 10 is typical. The corticoid metabolites have here been estimated by a modification of the method of Reddy *et al.* (1952) and show very little response to corticotrophin, whereas there is a distinct rise in ketosteroids from already excessively high levels. The urine of this patient also contained greatly increased amounts of pregnanetriol, the excretion of which was further raised by corticotrophin. The estimation of pregnanetriol is relatively simple and has appeared to be quite specific in this syndrome. It is, however, also included in the steroids estimated by the method of Norymberski *et al.* (1953), giving falsely high values for the level of corticoid metabolites. Wilkins and his group have observed a large series of these cases, and have found that cortisone, as suggested by Bartter *et al.* (1951) and especially if used early enough, has dramatic effects in alleviating the abnormalities. It also has a very powerful effect, as would be expected, on reducing

TABLE III.—*Steroid Excretion and Response to Corticotrophin—Adult Hirsutism*

Patient	Age	Summary	Resting (mg./day)		Response to Corticotrophin	
			Ketosteroid	Ketogenic Steroid	Max. Ketogenic Steroids (mg./day)	K.G.S./K.S. Ratio
1 F. S.	37	10 years' severe hirsutism accompanied by post-encephalitic changes from illness at age 7	20-27	10-4-13-1	27	0-3
2 L. C.	22	Hirsutism at puberty. Menorrhagia. Pseudomucinous cystadenoma removed at age 17, with one ovary. Histology of other ovary normal at 22. Other ovary had to be removed with uterus for menorrhagia. Urine oestrogen 13 µg./day after this	11-7-14-3	3-7-4-1	36	0-8
3 S. J.	28	Hirsute at 21	20-25	2-8-11-8	44	0-9
4 Z. C.	30	Hirsute at puberty. Virilized. Homosexual. Normal ovarian histology. Improved after subtotal adrenalectomy	10-12 (1-4-5-5 post-op.)	6-7-10-3	30	1-1
5 S. B.	21	Hirsute at 16. Severe. Normal E.U.A.	18-19	12-2-14-9	60	1-1
6 O. F.	25	Overweight and hirsute from after puberty. Severe acne ..	15-22	15-4-20-1	74	1-5
7 K. P.	26	Overweight 10 years, severely hirsute 4 years. Few purple striae. E.U.A. normal	17-20	12-4-15-4	50	1-2
8 M. L.	29	Severe hirsutism from puberty. E.U.A. normal	9-12	13-2-17-7	67	1-9
9 J. B.	30	2 years' severe facial hirsutism. A mild case	17-25	9-3-9-7	75	1-4

E.U.A. refers to examination of genital tract under anaesthesia.

the level of the ketosteroid output, and control of the dosage in this way is essential. It similarly depresses the excretion of pregnanetriol.

Post-pubertal Hirsutism

Patients suffering from post-pubertal hirsutism have been the subject of less fruitful research, and indeed are often grouped together under the term "idiopathic hirsutism." There is a striking incidence of the beginning of the disturbance at or soon after puberty, which is at a time when the adrenal cortex has assumed its adult pattern of secretion, especially with regard to ketosteroids (Mason and Engstrom, 1950). It is perhaps odd that they develop the degree of hirsutism that they often do, with little or no increase in their resting ketosteroid excretion. It has been shown that under conditions of increased stimulation with corticotrophin they are then capable of excreting amounts of ketosteroids relatively in excess of the normal. A group of nine patients have been studied.

The basal steroid levels in the urine and the data of response to corticotrophin for these patients are summarized in Table III. There is a general tendency for the ratio of ketogenic steroids to ketosteroids to correlate with their clinical state, Case 1 being the most severe and Case 9 the mildest. There seems to be little relation between this ratio and the basal ketosteroid levels. Dr. Brooks, in our laboratory, has kindly carried out more detailed investigations of the ketosteroid patterns in three of the worst of these patients, using methods he has been developing for this purpose (Brooks, 1956). The β -ketosteroids represent largely dehydroepiandrosterone (Table IV). One of the most virilized individuals (Case 4) showed the highest basal level and, like Case 2, responded well to corticotrophin. The excretion of this substance in Case 1 was poor. The fraction of polar ketosteroids is particularly important, as it represents the ketosteroids with 11-ketone or hydroxyl groups. That they do not form a large proportion of the ketosteroid excretion

TABLE IV.—*Ketosteroid Fractionation in Adult Hirsutism*

Case	Total Ketosteroid (mg./day)	β -Fraction (mg./day)	Polar Ketosteroids (mg./day)
1 { Resting	20-4	0-9	2-7
1 { Corticotrophin	49-2	2-4	8-1
2 { Resting	14-3	2-6	6-5
2 { Corticotrophin	41-3	13-7	6-5
4 { Resting	12-1	5-6	1-2
4 { Corticotrophin	27-6	11-4	4-0
Rheumatoid { Resting	11-4	2-9	<0-1
arthritis { Corticotrophin	18-5	4-0	3-7
Prepubertal { Resting	107	12-1	33-6
adrenogenital { Corticotrophin	182	19-1	56-6
syndrome			

after stimulation with corticotrophin would therefore indicate that the larger part of the ketosteroids is not primarily derived indirectly as corticoid metabolites, but almost certainly by increased ketosteroid synthesis by the gland. The highest proportion of the total ketosteroid as polar ketosteroid after corticotrophin (19%) was found in Case 1. This was appreciably less than in the patient studied with prepubertal adrenogenital syndrome, whose corresponding figure was 31%. The comparison is similar in the case of the resting levels of the polar fraction.

TABLE V.—*Pregnanetriol in Urine*

	Before Corticotrophin (mg./day)	After Corticotrophin (mg./day)
Case 1. Hirsutism	1-5	8-2 (4)*
" 2. "	1-1	6-0 (4)
" 4. "	1-2	1-6 (4)
" 5. "	1-3	4-5 (3)
" 8. "	1-1	7-6 (4)
Adrenogenital syndrome (F) ..	45-0	124-0 (3)
Rheumatoid arthritis (F) ..	1-9	2-3 (3)
" (F)	0-2	4-6 (3)
Sub. nephritis (F)	0-2	3-0 (4)
Hodgkin's disease (M)	—	2-0 (—)
Cushing's syndrome (M)	2-8	2-7/8 hours*

Figure in parentheses is day of corticotrophin administration.
* Equivalent to approximately 8 mg./24 hours.

TABLE VI.—*Comparison of Steroid Responses in Patients with Normal Adrenals and Those with Adult Hirsutism or Prepubertal Adrenogenital Syndrome*

	Normal	Hirsutism	Adrenogenital Syndrome
Resting corticoid metabolites	Normal	Normal	Normal or low
" ketosteroids	"	Normal to high	High to very high
" 11-oxyketosteroids	"	Normal	Often high
" β -ketosteroids	"	Normal or high	Sometimes high
" pregnanetriol	"	Normal	High
Corticoid metabolites response to corticotrophin	Good	Good	Poor
Corticoid metabolites/ketosteroid response to corticotrophin	High	Low	Very low

The excretion of pregnanetriol in these patients is not very different from that found in other individuals (Table V), even after stimulation with corticotrophin. The highest level reached is that in Case 1, which is similar in magnitude to that in the patient with Cushing's syndrome. It cannot therefore be inferred that there is an abnormality in pregnanetriol excretion comparable to that in the pre-pubertal adrenogenital syndrome. Comparison of the hirsute group with normal individuals and the findings in the adrenogenital syndrome are summarized in Table VI.

Time of Adrenarche

It is clear that the inhibition of synthesis of adrenal corticoid, characteristic of the adrenogenital group, is not present in the hirsute group, although they are inclined to excrete more ketosteroid than normal, especially when given corticotrophin. In the same way that the adrenogenital cases excrete more β -ketosteroid, so sometimes do the hirsute ones. The basis of the abnormality of the hirsute patients is still not clear, but a tentative hypothesis may be put forward. One patient in our series, not included among those mentioned, developed hirsutism at the age of 8 and puberty at 13. Yet in adult life she had normal ketosteroid excretion. It is suggested that she had an early adrenarche as mentioned by Talbot, Sobel, McArthur, and Crawford (1952), but which is poorly documented.

We think also that we have encountered an example of the reverse condition of delayed adrenarche in a boy who did not develop puberty until the age of 18, when F.S.H. became detectable in the urine and ketosteroids were still only 4.6 mg. per day. The response of ketosteroids to corticotrophin in particular was poor at the ages of 16 and 18. The latter was confirmed with intravenous corticotrophin. A maximal dose of 40 "clinical units" intravenously was sufficient to stimulate the adrenal more normally. The fact that in younger normal children and in this boy ketosteroids do not rise to adult levels with corticotrophin suggests that there is some factor operating, either in the adrenal cortex or possibly an additional stimulating factor, which brings about the normal pattern after the adrenarche. This factor does not appear to be chorionic gonadotrophin, as suggested by Albright. It would therefore appear that the lesion in hirsute women may depend on an abnormality of the accessory factor which enables the adrenal to respond by secreting excessive quantities of normal ketosteroids. The probable presence of a hypothalamic lesion in Case 1 should be specially considered as a cause of such maladjustment. The relation of the hypothalamus to the adrenal cortex is further discussed.

Treatment with corticotrophin of a group of patients with anterior hypopituitarism over a period of years is reviewed more fully below. For the present it is sufficient to note that, although their responses were good in other respects, their ketosteroid levels were always on the low side of normal. It was not possible to obtain adequate hair growth with the corticotrophin alone; the men required testosterone, and in the women pubic and axillary hair remained below normal standards. This might indicate that in these patients a factor is lacking which enables the adrenal to secrete normal amounts of androgen. The question of the exact nature of the androgenic hormones from the adrenal cannot yet be regarded as entirely settled (Munson, Goetz, Laidlaw, Harrison, and Thorn, 1954; Kappas *et al.*, 1955).

Failure of Cortisone Therapy in Hirsutism

The importance and value of the use of cortisone in suppression of excess ketosteroid secretion in the pre-pubertal adrenogenital syndrome has been mentioned. It seemed worth while to explore its value in hirsute women. The property of suppression of ketosteroids is not confined to patients with the pre-pubertal syndrome (Copeman, Savage, Bishop, Dodds, Kellie, Stewart, Glyn, Henly, and Tweed, 1952; Pond, 1955). We have treated five hirsute women with cortisone (Table VII). Doses of cortisone ranging from 37.5 to 125 mg. daily were given for periods from three months to three years. In none of these patients have there been any really satisfactory changes in the state of the hirsutism. This finding is, however, to be expected, since it has been pointed out by Wilkins *et al.* (1955) from their large experience that well-established hirsutism is slow and difficult to improve in the adrenogenital syndrome. This has also been our experience.

It is of great interest to note that Wilkins *et al.* (1955) reported that one of their adult patients had developed masculine baldness, and with cortisone treatment had a new growth of hair

TABLE VII.—Excretion of Ketogenic Steroids and Ketosteroids After Administration of Cortisone

Case	Sex	Cortisone Acetate Dose* (mg., day)	Ketogenic Steroids (mg./day)	Ketosteroids (mg./day)
A Hypophysectomy; car. prostate	M	25	6.8 (27%)	4.3
B Hirsute	F	25	9.1 (36%)	9.2
C .. (Case 4, Table I:1) after adrenalectomy	F	37.5	14 (38%)	3.6
B Hirsute	F	50	16-20 (32-40%)	7
D .. (Case 5)	F	50	22 (44%)	10
E Attempted hypophysectomy; car. breast ..	F	50	25 (50%)	6.5
A Hypophysectomy; car. prostate	M	50	27 (54%)	8.0
B Hirsute	F	75	36 (48%)	6
F .. (Case 3)	F	100	22 (22%)	6
E Attempted hypophysectomy; car. breast ..	F	100	52 (52%)	5.5
G Hirsute (Case 1)	F	100 i.m.	35 (35%)	5.6
H Lupus erythematosus	F	125	19-44 (15-35%)	10-24
I Iridocyclitis	F	250	93 (37%)	9.7
J Exfoliative dermatitis	F	375	133 (35%)	65
	M	2,000	137-620 (7-31%)	18-44

* Divided oral doses.

on the scalp. We encountered a dramatic and rapid growth of normally pigmented hair on the head in a completely bald woman of 44, who was under treatment for an eye lesion, with large doses of cortisone (375 mg. scaling down to 50 mg. daily). This event prompted its trial in a second patient. This latter had ovarian dysgenesis accompanied by complete baldness, which had been present for several years. Although she improved greatly in other respects, oestrogen therapy had no effect on the scalp until an addition was made of 100 mg. of cortisone daily. In the next three weeks it was possible to discern a definite growth, and several weeks later she had a satisfactory head of hair. The mechanism of this is of interest. The baldness in the second patient could hardly be ascribed to excess of adrenal ketosteroid, as it was in the patient of Wilkins *et al.* (1955).

Several years ago we observed that oestrogen therapy had no effect on the excess ketosteroid excretion in a patient with pre-pubertal adrenogenital syndrome, and this has been found by others. On the other hand, after gonadectomy of a male pseudohermaphrodite there was some suppression of adrenal ketosteroid, but not of the dehydroepiandrosterone fraction (Prunty, 1953). We have administered oestrogen to two of our hirsute patients. In one of these, who was co-operative and obsessional about her condition, we were able to collect and weigh facial hair shavings. Both cortisone and ethinyloestradiol seemed to produce a small depression of ketosteroid excretion (from about 8 to 10 mg. per day to 6 mg. per day). The rate of hair growth, approximately 6.2 mg. daily on 50 mg. of cortisone daily for three weeks, was unchanged. Ethinyloestradiol for a period of up to 120 days was also without effect, in spite of the fact that the patient was quite convinced that it caused improvement. This presumably was due to an improvement in the texture of the skin. Hydrocortisone ointment applied locally also produced no result. On the other hand, the situation in the adrenogenital syndrome is probably somewhat different. A combination of cortisone with oestrogen (0.2 mg. of ethinyloestradiol daily for three in every four weeks) appears to be more effective than the use of larger doses of cortisone alone. Armstrong (1955) has published similar and more convincing evidence of this effect in three patients with this syndrome.

During the course of cortisone therapy we have made observations on the recovery of cortisone metabolites by the method of Norymberski *et al.* (1953) (Table VII). These authors concluded that a recovery of about 40% of the administered dose was obtained. Except when the cortisone dosage is high, the recovery figures are unreliable because of the unpredictable depression of the endogenous corticoid secretion. After subtotal adrenalectomy patient C excreted 4 mg. of ketogenic steroids daily, so that net cortisone recovery must have exceeded 27%. The average

recovery at dose levels above 37.5 mg. of cortisone daily is 40%. Nevertheless patient J, receiving 2,000 mg. of cortisone daily, excreted at times as little as 7% as ketogenic steroid. The ketosteroid figures show that significant amounts begin to appear at dose levels above 100 mg. daily. In patient J the highest ketosteroid level corresponded with the lowest ketogenic steroid level, and vice versa. The metabolism is therefore somewhat unpredictable. It is important to note that in the five hirsute patients in Table VII the ketosteroids were depressed by an average of about 30%. There was therefore no indication of an abnormal increase in ketosteroid excretion in these particular patients when cortisone was given. The fall in recovery after intramuscular as opposed to oral cortisone administration should be noted. This has also been observed during the course of treatment of a patient with the pre-pubertal adrenogenital syndrome. It is probably dependent on the slower rate at which the steroid is taken to the liver.

Observations on a Group of Male Obese Patients

In Table VIII are listed seven obese male patients. Clinically these patients are divisible into at least two categories. Firstly, there are the two juveniles who respond to corticotrophin like the normal children. Secondly, the adults show a gradation in their behaviour. Patients D.C. and L.D. responded like a normal adult with respect to the excretion of ketogenic steroids. The others all gave poor ketogenic steroid responses to corticotrophin (Table II), in spite of the fact that they received the potent batch (89754) given to the hirsute women. In addition they show a gradation with respect to the ketogenic steroid:ketosteroid ratio, the extreme case giving a ratio as low as 0.6. This patient also responded to intravenous corticotrophin with a bigger increase in ketosteroids than ketogenic steroids. The intravenous test in R.C. showed a similar trend, whilst

TABLE VIII.—Obese Male Patients

Patient	Age	Summary	Ketogenic Steroid (mg./day)	Ketosteroid (mg./day)	K.G.S./K.S. Ratio on Corticotrophin
R.P.	12	Always tended to be overweight. Plethoric obese lad with white striae on limbs. No hypertension	7	7	2.7
I.S.	15	5 years' obesity. No signs of puberty	7	3	5.1
R.S.	25	Overweight since age 7. Plethoric. Obesity limbs and trunk. Reddish striae present. No hypertension	16	23	0.6
R.C.	20	40 lb. (18.1 kg.) weight gain in 1 year. Slightly plethoric. Reddish abdominal striae. Mild hypertension and early diabetic sugar curve. Plasma hydrocortisone 6 µg./100 ml.	13	13	1.1
D.C.	25	Obese since age 10. Generalized obesity with reddish striae. Mildly feminine habitus. No hypertension. Plasma hydrocortisone 5 µg./100 ml.	10	8	1.8
L.D.	24	3 stone (19 kg.) weight gain in 4 months. Polio, age 16. Plethoric and rounded face. Moderate reddish striae and acne. No hypertension. Very good response to restricted diet	16	17	1.3
P.H.	22	5 stone (31.8 kg.) weight gain to 17½ stone (111 kg.) in 4 years. A few small striae (pale) with generalized obesity. No hypertension	17	12	2.8

that in D.C. and L.D. was of a more normal pattern. It is to be noted that R.S. had the highest level of resting ketosteroids. The significance of these findings is not clear at the present time. The results show that there are conditions in the male when the ketogenic steroid:ketosteroid response to corticotrophin can resemble that in the abnormal women.

Abnormal Responses to Operation

The normal findings in the response of steroid excretion to the trauma of operation have already been discussed. Amongst the hirsute women it was possible to study the

FIG. 11.—Response to operation (hysterectomy) in a patient with adult hirsutism (L.C., Case 2). Note before operation abnormal pattern of corticotrophin response (see text).

response to hysterectomy (Case 2) and this is shown in Fig. 11. It will be recalled that this patient responded to corticotrophin with a higher ketosteroid excretion than ketogenic steroid excretion. The response to operation shows a similar trend, in general the rise in ketosteroids exceeding that of ketogenic steroids. The response of the adrenal therefore seems to depend upon endogenous conditions in the patient and not merely to the properties of the corticotrophin administered.

Conditions Under Which Ketogenic Steroid Excretion Falls Below That of Ketosteroids After Adrenal Cortical Stimulation

The conditions which tend to cause reversal of the ketogenic steroid:ketosteroid ratio can now be summarized. This has been encountered in the following circumstances: (1) Adult hirsutism. (2) Certain adult obese males. (3) When the corticotrophin stimulus to the adrenal cortex is lessened—that is, when the response is diminishing. An example of this is seen in Fig. 7 at day 40. This presumably indicates that the rate of corticoid secretion falls off first. This tendency has also been observed during the “tail-off” period after intravenous corticotrophin. (4) With different batches of corticotrophin there appears to be a definite tendency with less potent batches of corticotrophin for the ratio to be lower than usual. The effect of a potent batch (89754) is strikingly seen in Fig. 7 at day 91, where there is sudden reversal in the ratio with a small upward trend of the ketogenic steroids.

Forsham *et al.* (1955) noted a difference in the ratio of corticoid metabolites to ketosteroids, particularly with different batches of corticotrophin. This finding indicates that caution must be used in the interpretation of variations in the ratio. It is the reason why comparisons have at present been restricted to those observations made with a known potent corticotrophin (89754) which has been shown to be particularly active in causing an increase in ketogenic steroid excretion rates. While these observations were being concluded, Laidlaw *et al.* (1955) published data similar to ours in adult virilized women. They took care to exclude ovarian pathology as a cause for the disturbance in their patients.

Response of Patients With Hypopituitarism to Corticotrophin Therapy

No doubt largely because of the practical difficulty of administration, the long-term use of corticotrophin in

patients with anterior hypopituitarism is rarely reported. Furthermore, discouraging reports appeared suggesting that complications, such as severe water retention (Whittaker and Whitehead, 1954) or cerebral disturbances, occurred after short periods of administration (Summers and Sheehan, 1951). Robertson and Kirkpatrick (1951) treated a patient for 32 days without ill effect. The report of the longest use of corticotrophin is that of Aber, Chandler, and Hartfall (1954), who gave it to one patient for 160 days. They recommend that as little as 40 "clinical units" of gel, given once weekly as a maintenance dose, may be sufficient. The data presented below indicate almost surely that this is far too infrequent administration of the hormone, and would be likely to be dangerous in a severely affected patient.

Three patients with hypopituitarism have been treated with corticotrophin for two to three years. This enabled us to study the continued response of the adrenal and assess the amount of corticotrophin required in patients in whom it is virtually absent. The first patient (J.L.) was a man aged 28 with typical hypopituitarism and failure of puberty to develop owing to an apparent small chromophobe adenoma (Mills, 1955a). E.R. was a man aged 62 with typical features of about three years' duration, ketosteroids falling to 1.5 mg. per day. The aetiology of his condition has never been established. L.L. was a woman aged 44 with classical Simmonds's disease due to post-partum haemorrhage seventeen years previously. Each patient noticed pronounced subjective improvement on corticotrophin therapy, with weight gain and a feeling of vigour not experienced previously. The effects on the thyroid and hair growth have been discussed. There was reversal of abnormal water diuresis and insulin sensitivity tests.

The response to the hormone over the last twelve to fourteen months in these three patients is shown in Fig. 12. It will be seen that they are sensitive to dosage, one patient in particular (E.R.) giving excessively high ketogenic steroids on 20 "clinical units" of corticotrophin gel twice daily. On the whole, normal levels of ketogenic steroids were produced by doses of rather less than 20 "clinical units" of gel once daily. There were times when weight

gain and fullness of the face became excessive, in spite of the fact that corticotrophin dosage was not otherwise thought to be too large. This, of course, particularly occurred in J.L. when his ketogenic steroids reached 37 mg. daily and in E.R. when they reached 84 mg. daily. But these signs sometimes occurred with ketogenic steroids in the neighbourhood of 20 mg. They suggested reduction of dosage. The effect of poor local absorption is seen in L.L., when for a time 25 "clinical units" daily gave only 1.5 mg. of ketogenic steroid in the urine. This may have been due to local infection at the injection site. The occurrence of a severe cold at another time in this patient did not have the same effect. This investigation showed, therefore, that it was possible to maintain adrenal cortical stimulation with corticotrophin for prolonged periods. It also gave valuable information concerning the quantity required to maintain normal levels of ketogenic steroids, but at the same time the ketosteroid excretion tended to be below normal.

Observations on Cerebral Disturbances in Anterior Hypopituitarism

Anxiety to reduce the dosage of corticotrophin in J.L. ultimately led to a serious state of affairs.

This patient was treated for four months with 10 "clinical units" daily, together with a testosterone implant and thyroid, and was clinically very well, having a ketogenic steroid level of 6.6 mg. daily. He had a very full facial appearance, and the dosage was further reduced to 10 "clinical units" on alternate days. After two months on this dosage his ketogenic steroids, during the day of injection, were 4.9 mg. per day. Shortly after he complained of left temporal headache, which became progressively worse and vomiting began. The nature of the headache, which was very persistent, suggested the possibility of an intracranial lesion. On admission for investigation his ketogenic steroids were only 0.3 mg. per day during the 24 hours after injection of corticotrophin. It seems most likely that this fall was due to inadequate dosage (compare Fig. 11) rather than to any resistance on the part of the adrenal to corticotrophin. He lay curled up in bed and was resistant to interference, with moderately frequent vomiting. He was put on cortisone, 75 mg. intramuscularly daily. At this time his blood pressure fell to 95/65 mm. Hg. Meanwhile investigations of the cerebral condition showed no change in the appearance of the sella turcica and all other investigations were normal. However, the E.E.G. had shown a pattern suggesting a right temporal focal lesion and, six weeks later, a pattern compatible with the presence of a craniopharyngioma. A month after starting cortisone he showed significant improvement and two weeks later had resumed his normal health. At this stage the E.E.G. had almost returned to normal.

A second patient with severe anterior hypopituitarism was a woman aged 54 who twenty-three years previously had developed amenorrhoea following apparently normal parturition. One day she complained of headache and feeling unwell and went to bed, where she lapsed into coma for thirty-six hours. She was admitted to hospital in a state of severe irritability and incontinent of urine and faeces. Tendon and abdominal reflexes were absent. Lumbar puncture was performed with extreme difficulty on account of her irritable condition, and the C.S.F. contained 70 mg. of protein and 6 leucocytes per c.mm. Blood pressure was 75/45 mm. Hg. After two days she responded to a glucose-saline drip. She had a second milder attack which responded to "eucortone," testosterone, and salt. A year later she was admitted in coma with convulsions and stertorous breathing. She had felt vaguely unwell, ate nothing for three days, and lapsed into coma. All tendon reflexes were difficult to elicit. The blood sugar was 45 mg. per 100 ml. In spite of treatment with dextrose and eucortone intravenously she died in two hours. There was a cyst 10 mm. in diameter in the pituitary fossa. No thyroid or adrenal tissue was found and both ovaries were small.

FIG. 12.—Response of three patients with hypopituitarism to long-term corticotrophin therapy (see text).

In a third instance of difficulty with a stuporous patient the findings were even more confusing.

A man aged 47 had had a chromophobe adenoma operated upon five years previously. After a period of good health he returned with malaise, depression, headache, and vomiting. Full neurological investigations were made, and it was considered that he had no recurrence of his pituitary lesion. He was transferred to us by Mr. Harvey Jackson for endocrinological investigation. He was somnolent and would answer only one sentence at a time. He regularly vomited in the early mornings. He had a small stubbly beard, small and soft testicles, and somewhat sparse axillary and pubic hair. Blood pressure was 120/75 mm. Hg. The fundi were normal. The insulin tolerance was normal. Keto-steroids were low, but response to corticotrophin was good. Excretion of radioactive iodine was 10.5% in twenty-four hours and 63% in forty-eight hours. The E.E.G. was abnormal, "consistent with a central space-occupying lesion or biochemical disturbance." It was decided to repeat ventriculography, but before this could be done the patient suddenly died. At necropsy a secondary carcinoma, by meningeal spread, 2 cm. in diameter, was found in the midline above the cerebellum (Fig. 13). This was causing

FIG. 13.—Carcinoma of anterior pituitary. A. Malignant cells in remaining tissue in pituitary fossa after previous operation. B. Section through secondary neoplasm above cerebellum. Several small secondaries were also present on the meninges.

a moderate degree of internal hydrocephalus. The thyroid acini showed flattened epithelium, some atrophy of the adrenal cortical fasciculata had occurred, and there was a decrease in testicular interstitial cells, but spermatogenesis was active but reduced. It must be concluded that the patient had only a mild degree of hypopituitarism, and his symptoms were mostly due to internal hydrocephalus.

Metastasizing carcinoma of the pituitary is exceedingly rare, but has been found to spread by the meninges (Willis, 1952). Local infiltration by carcinomas of the pituitary gland is a rather more commonly recognized condition (King, 1951).

Observations on Cerebral Disturbances in Addison's Disease

The problem of the differentiation of cerebral irritation due to organic cause as distinct from a metabolic one also arises in Addison's disease. I have observed this confusion in three instances, and am indebted to Dr. G. W. Thorn for permission to mention the first two of these.

The first was a man aged 31 who had developed severe Addison's disease nine years previously and who five years later also developed diabetes mellitus (see Thorn and Clinton, 1943). He was readmitted to hospital in a state of Addisonian crisis, on which supervened a diabetic crisis. He was treated for both conditions, but did not resume a normal mental state for five days. Most of the time he was comatose and apathetic, but during practically the whole of this period developed typical fits of Jacksonian epilepsy, starting unilaterally and sometimes becoming generalized. For several days they occurred with a fre-

quency of about once hourly. The strong temptation to perform ventriculography was resisted. On satisfactory restoration to normal metabolic conditions the attacks completely disappeared. It was not possible to correlate these occurrences with any particular biochemical abnormality.

The second patient was a boy aged 12 who complained of recent headaches, drowsiness, and fainting spells. Nystagmus, a degree of neck rigidity, and varying tendon reflexes had been noted. The E.E.G. showed the typical slow-wave activity seen in Addison's disease but suggesting localization in the right frontal region. Trephine of the skull was carried out for exploratory purposes, and this operation precipitated an Addisonian crisis, from which he subsequently recovered.

The third example also concerned a child, a girl aged 10, who developed listlessness and dizzy spells. She complained of having had diplopia and blurred vision a month previously. She was found to have an ataxic gait. She lay curled up on her side, resisted interference, and was very irritable if roused. Questionable meningism was also occasionally present. In spite of the presence of developing pigmentation, the possibility of a posterior fossa neoplasm was considered. The E.E.G. showed high-voltage slow-wave activity and, it was thought, also suggested a posterior fossa lesion. She was not hypoglycaemic. It was nevertheless decided to treat her for Addison's disease, but within an hour before this was done she went into crisis from which she could not be rallied. There was complete atrophy of the adrenal cortex.

These examples of cerebral disturbance in hypopituitarism and Addison's disease draw further attention to a diagnostic problem. In Addison's disease some of the accompanying well-known mental changes were mentioned by Addison himself and later emphasized by Cleghorn (1951). They include apathy, negativism, depression, inactivity, paranoia, and delusions, and are dangers which are usually readily recognized. The occurrence of other phenomena suggestive of the presence of intracranial lesions is more disturbing and must of necessity, for a time at least, be taken more seriously. Engel and Margolin (1941) were first to draw attention to the fairly typical abnormal electroencephalogram with high-voltage diffuse slow activity which occurred in their patient who was having convulsions. Storrie (1955) has described three further somewhat similar cases. He also observed convulsions. His observations did not support the thesis of Engel and Margolin that the abnormalities were the direct result of hypoglycaemia. The comatose states seen in hypopituitarism, have also been recently discussed by Caughey and Garrod (1955). These authors suggested seven possible causes of a comatose condition: post-operative, infections, hypoglycaemia, water intoxication, sodium depletion, hypothermia, and unknown. To this list should be added presence of an expanding intracranial lesion.

In an earlier paper Caughey and Garrod (1954) have stressed the importance of poor adrenal function, while the observations of Sheehan and Summers (1952) certainly indicate that it is not always an easy therapeutic problem to treat these cases of coma with the means available, although they were not then able to employ intravenous hydrocortisone. Our second patient was an example of severe adrenal cortical failure in which crisis was precipitated by mild infection. The disturbance in the first patient with hypopituitarism must be ascribed to slowly developing adrenal cortical failure, but it is important to note the long length of time he took to recover on receiving cortisone therapy. With increasing knowledge of the endocrine changes occurring in hypopituitarism the differentiation of neurosurgical and endocrine causes for cerebral symptoms in these patients will become increasingly more important.

The Hypothalamus and Pituitary Function

The mechanisms governing the control of corticotrophin are at present much debated. One method of regulation is almost certainly by way of hypothalamic control of the activity of the anterior pituitary. The following case is therefore of great interest.

J. G., a man aged 27, weighed 16 st. (101.6 kg.) (Fig. 14), but had exceeded this and was particularly troubled by nocturnal

enuresis, having a daily urine output of 12 litres, containing up to 400 g. of glucose. At the age of 9 he had had "poliomyelitis" and for some time had noticed blurring of vision. He was somnolent by day, but not sufficient to prevent his working. He had gynaecomastia, sparse head and facial hair, but almost normal axillary and pubic hair. His external genitalia were small. Reduction of visual acuity accompanied bilateral papilloedema with numerous small circumscribed retinal haemorrhages. He had left-sided ptosis and signs suggesting old poliomyelitis in the left arm. His voluntary food consumption was equivalent to 7,000 calories daily.

FIG. 14.—Case J.G. Obesity and gynaecomastia associated with chronic encephalitic lesion especially involving hypothalamic region.

The E.E.G. showed bilateral slow high-voltage activity, especially in the frontal regions, paroxysmal in nature. It was increased by overbreathing and stabilized by photic stimulation. Apart from what turned out to be a falsely positive ethyl iodophenyl-undecylate ventriculogram, suggesting a filling defect in the third ventricle, other neurological investigations were normal. The blood sugar was 300 mg. per 100 ml. and liver glycogen only 0.74%. There was never excess ketone in blood or urine. The blood sugar was unresponsive to adrenaline. F.S.H. was absent from urine. All epiphyses were united. After later control of the glycosuria with insulin he concentrated his urine to only S.G. 1014 by dehydration or with "pitressin." The B.M.R. was plus 15%, but thyroid uptake of ^{131}I was only 10%. His response to corticotrophin (89754), both intramuscularly and intravenously, was poor and suggestive of hypopituitarism. At first he required 240 units of soluble insulin daily to control his glycosuria, with a reduction of urine volume to 2 litres. He was then put on a 2,000-calorie diet and given x-ray therapy to the infundibular area, whereupon his insulin requirement fell to 30 units daily. He was discharged on insulin and thyroid, but several months later returned with cerebellar symptoms and signs and died with hyperpyrexia.

At necropsy there were old encephalitic lesions of generalized nature, but specially involving the neighbourhood of the tuber cinereum and mammillary bodies (Fig. 15). There were extensive and more recent haemorrhagic lesions in the pons. The pituitary appeared normal; the thyroid showed vesicles filled with colloid and flattened epithelium (Fig. 16); the adrenals were small, and the fascicula was particularly narrow, with pronounced pigment granules in the inner zone and shrunken reticula; the testicles showed a considerable impairment of spermatogenesis with prominent Sertoli cells, and Leydig cells were reduced in number. Breast tissue showed considerable duct hyperplasia.

Harris (1955b) has carefully reviewed the evidence from experimental work on the hypothalamus and concludes that external stimuli to the anterior pituitary can pass by two pathways. The ordinary day-to-day stimuli are transmitted through nervous pathways, whereas stimuli can also pass by a direct metabolic mechanism. The animal experiments on which this conclusion is based are of two types: section of the pituitary stalk with care to avoid regeneration of the portal vessels, or stimulation of the various parts of the hypothalamic region and production of experimental lesions in this region. Thus stimulation of the posterior part of the hypothalamus in monkeys causes corticotrophin release (Porter, 1954). Pituitary stalk section need only lead to slight atrophy of the anterior pituitary and some atrophy of the adrenal cortex. It is reasonable to suppose that in our patient sufficient stimulation of the anterior pituitary remained by extrahypothalamic pathways to maintain the adrenal in its subnormal state of activity and to give the

somewhat reduced positive response to injected corticotrophin. The decreased uptake of ^{131}I in our patient closely follows Harris's (1955b) finding of reduced thyroid activity in the rabbit with pituitary stalk section.

With regard to gonadotrophin secretion, most experimental work has been done in female animals, but Benoit and Assenmacher (1953) found that isolation of the anterior pituitary of the duck caused decreased testicular function and atrophic changes. The changes in our patient were of partial degree and less severe than those often seen in anterior hypopituitarism (Howard, Sniffen, Simmons, and Albright, 1950). Similarly, the adrenal atrophy had not progressed to the degree often occurring in anterior

FIG. 15.—Case J.G. Typical encephalitic lesion adjacent to mammillary body. ($\times 280$.)

FIG. 16.—Case J.G. A. Quiescent thyroid. B. Testicle. Note the basement membrane, diminished spermatogenesis, prominent Sertoli cells, and reduction of interstitial cells. ($\times 110$.)

hypopituitarism (Sheehan and Summers, 1949). In a series of 60 cases showing endocrinological disturbances as a result of damage in the hypothalamic area meningo-encephalitic lesions were uncommon (Bauer, 1954). The most prominent change was alteration of sexual pattern, precocious puberty, or hypogonadism. In the latter group tubero-infundibular lesions predominated, and when examined the pituitary was usually normal. Patients with encephalitis most resembling ours, but less completely investigated, are those of Moore and Cushing (1935) and of Ayala (1934). They both, however, had diabetes insipidus; in the former, irradiation of the region of the sella reduced urine volume. In monkeys the hypothalamic nuclei are known to be sensitive to irradiation (Arnold, 1954).

Glycosuria is a rarity in such cases and ketosis seems to occur (Paton and Petch, 1954). Long (1940) reviewed the evidence, including that from experiments, that hypothalamic lesions cause glycosuria, and concluded that they do not permanently alter the control of blood glucose. He favoured the view that hypothalamic or brain-stem lesions cause disturbance via the sympathetic system and adrenal medulla. Our patient had no evidence of sympathetic overactivity, except possibly the small increase of basal metabolic rate. His history seemed to indicate the presence of polyuria due to glycosuria for a long period. The complete absence of ketosis, even with severe depletion of liver glycogen, still needs explanation.

Developing Ideas in Relation to Cushing's Syndrome

In typical cases the diagnosis of Cushing's syndrome is simple; in the causative type of adrenal cortical lesion the diagnosis may prove more difficult, and also in the less typical and related cases. It was only recently noticed how poor is the prognosis of the condition; of 32 patients, 17 were dead within five years (Plotz, Knowlton, and Ragan, 1952). In a study of a series of 40 patients, Soffer, Eisenberg, Iannoccone, and Gabrilove (1955) noted significant deviations from the classical form of obesity. Indeed, the facies may not always show the classical "moon" picture, but more nearly suggest benign obesity. The commonest abnormality, present in 80% of patients, is hypertension, which ultimately leads to irreversible systemic damage. The possible presence of a malignant tumour needs serious consideration before becoming inoperable or metastasizing. When appraising the views of our American colleagues it must be remembered that the occurrence of benign and malignant tumours is more frequent in North America than in Europe. Of 127 American cases, 40% had tumours (Sprague, 1953; Soffer *et al.*, 1955; Forsham *et al.*, 1955). European figures are more difficult to obtain, but the incidence in 57 cases appears to be about 7% (Luft, 1955; Querido, 1955).

It was once thought that the level of 17-ketosteroid excretion was a valuable sign (Forbes and Albright, 1951), but this is only sometimes true (Soffer *et al.*, 1955). The use of recent methods of assay of corticoids and their metabolites has been of help in the diagnosis of Cushing's syndrome, but not of the presence of tumours. In a series of 62 patients, 8 gave normal or doubtful values (Sprague, 1953; Soffer *et al.*, 1955). With a different technique Dustan *et al.* (1953) reported 8 out of 11 patients with normal values. It seems undoubted that a few patients with upper normal values will be found (Laidlaw *et al.*, 1955). It may, however, yet prove to be the case that an estimate of the free or unconjugated hydrocortisone and cortisone in the urine is of more definitive value (Gray, 1955; Cope and Harrison, 1955; see also Dustan *et al.*, 1953). Estimates of plasma corticoids and metabolites give a similar picture, and may be

FIG. 17.—Response in urine steroids of patients S.E. and N.R. with non-tumorous adrenal cortices in Cushing's syndrome to corticotrophin. A. Corticotrophin intramuscularly. B. Intravenously. (Compare Fig. 4.)

FIG. 18.—Responses of patient with Cushing's syndrome with non-tumorous gland. Failure of depression of steroids by 9α-fluorohydrocortisone, 10 mg. daily by mouth. Considerable increase in ketogenic steroid excretion after unilateral adrenalectomy (see text).

difficult to interpret because of the possibility of confusion with emotional factors (Bayliss, 1955).

An approach to the differentiation of malignant tumours from adrenal hyperplasia is the use of corticotrophin. Patients with overactive adrenals are very responsive (Fig. 17). Patient S. E. had adrenal hyperplasia and N. R. had adrenals of normal size but prominent zona fasciculata; both had pronounced Cushing's syndrome. Caution is needed in carrying out these tests, for cardiac failure may be precipitated. A rapid increase of blood corticoids or excretion of corticoid metabolites in patients of this type has also been noted by Burnett, Rukes, Kolb, and Forsham (1954) and by Laidlaw *et al.* (1955). The lability of the level of steroid excretion should be noted. In patient S. E. plasma hydrocortisone was 25 μ g. per 100 ml. and in N. R. values of 11 and 24 μ g. per 100 ml. were found.

On the other hand, the response of patients with adrenal cortical carcinoma to corticotrophin is said to be absent, on account of the autonomous nature of the tumour cells. This was not our experience in patient J. H. (Brooks, McSwiney, Mills, Prunty, Robertson, and Wood, 1956), who showed a significant rise of ketogenic steroids and ketosteroids with corticotrophin. The resting output of these steroids was high, but the plasma hydrocortisone was only 8 to 10 μ g. per 100 ml.

A second possible method of differentiation of tumour patients is that suggested by Venning, Pattee, McCall, and Browne (1952). The principle is to suppress cortical secretion of a non-tumorous gland by administration of a substance which will suppress corticotrophin secretion. Cortisone does this, but, since it also gives rise to significant quantities of corticoid metabolites, it is necessary to measure suppression of ketosteroids. This technique has been fairly successful in the hands of Jailer, Gold, and Wallace (1954), but cannot be regarded as very satisfactory (Wilkins, 1955). With the advent of 9 α -fluorohydrocortisone, which is active in very small amounts, the situation has theoretically changed (Thorn, Renold, Morse, Goldfien, and Reddy, 1955), as it produces insignificant amounts of measurable metabolites. Aldosterone is not powerful enough in its corticoid effects to suppress the anterior pituitary to a useful extent (Prunty, McSwiney, Mills, and Smith, 1954). Cope and Harrison (1955) have observed suppression of free hydrocortisone in the urine with 9 α -fluorohydrocortisone, but its effects in the hands of Nabarro and Walker (1955) have been variable. In Fig. 18 is seen its effects in patient N. R. No significant steroid suppression occurred.

Since the description by Conn (1955) of a patient with adrenal cortical tumour a new facet of the problem has appeared, and as many as 13 similar patients are now known (Conn and Louis, 1956; Brooks *et al.*, 1956). So far as can be determined at present, important features are hypertension, tendency to obesity, polyuria, and cramps with muscular weakness. These need not be present simultaneously. These patients also have low serum potassium levels and alkalosis. For many years these blood changes have interested the Mayo Clinic group, headed by E. J. Kepler; a recent survey of 83 patients with Cushing's syndrome has shown that 41% had alkalosis and 13% low serum potassium (Sprague and Power, 1953). These changes occurred in patients with both hyperplasia and tumour. An important additional biochemical finding in the group of atypical cases mentioned may be the presence of excess aldosterone in the urine. This can be present with normal corticoid metabolites and ketosteroids (Conn and Louis, 1956), or with elevation of these substances (Foye and Feichtmeir, 1955; Brooks *et al.*, 1956), or with an apparent increase of corticosterone (Mader and Iseri, 1955). Conn gave the unfortunate term "primary aldosteronism" to the condition to distinguish it from that state in which patients with subacute nephritis also excrete excessive aldosterone. Adrenal cortical tumours need not necessarily be present (*Lancet*, 1955; Conn and Louis, 1956), but adenomas are usual, and occasionally carcinoma (Foye and Feichtmeir, 1955; Brooks *et al.*, 1956). The exact relationship of aldosterone secretion to corticotrophin stimulation is not settled, but the pituitary hormone

does not appear to be essential (Gaunt, Renzi, and Chart, 1955; Falbriard, Muller, Neher, and Mach, 1955).

The lability of secretion of corticoid in these patients with Cushing's syndrome is also shown in the very pronounced rise in ketogenic steroids observed in response to unilateral adrenalectomy. From radiological evidence, I.V.P., and air insufflation it was thought likely that N. R. had a tumour of the left adrenal cortex. At operation it was found that the appearances were due to a large fibrofatty pad of tissue lying above the adrenal, which complicated the surgical procedure kindly carried out by Mr. R. H. O. B. Robinson. After the removal of this adrenal there was a sharp increase in ketogenic steroid excretion (Fig. 18) and the operation was followed by a severe sepsis of the area that was difficult to control. In addition there developed intractable haematemesis from large acute gastric ulcerations, from which the patient ultimately succumbed. This difficulty of healing has also been noted by Jepson *et al.* (1956) in a patient who did not have Cushing's syndrome but spontaneously responded in an exaggerated fashion with respect to the secretion of ketogenic steroids. The occurrence of spontaneous intestinal haemorrhage in Cushing's syndrome is apparently rare, Plotz *et al.* (1952) noting a death rate from this cause of less than 2%.

On the other hand, the occurrence of gastric bleeding in patients undergoing treatment with cortisone is quite well recognized. In Cushing's syndrome, where there is such lability of corticoid secretion, there may be risks in the post-operative period if only unilateral adrenalectomy is performed as a first-stage operation. The type of post-operative steroid response in this patient is also important. It resembles the response to corticotrophin, the ketogenic steroid:ketosteroid ratio being high, and is quite different from the response of the hirsute patient L. C. (Fig. 11).

Adiposity Resembling Cushing's Syndrome

Simpson (1951, 1953) made a study of a group of patients with obesity commencing in childhood who, he felt, had some form of altered adrenal cortical activity. Among the patients reviewed here several have had signs, particularly obesity and red striae, suggesting altered adrenal function (see L. D. and R. S. in Table VIII), but hypertension was mild or absent. Their response to corticotrophin does not show the pattern of Cushing's syndrome; indeed, in R. S. it was quite different. An extreme example is C. L., aged 18, who developed obesity at the age of 10; puberty was normal, but suddenly at 18 he developed very severe red lineae distensae (Fig. 19). He had a full plethoric face, a pre-diabetic sugar curve, and mild hypertension. His tallness seemed to be familial. X-ray studies, serum electrolytes, urinary ketosteroids, and ketogenic steroids were normal. Plasma hydrocortisone was 2.5 and 3.5 μ g.%. He showed a normal (not increased) response to corticotrophin (89754). He improved after pituitary irradiation and with reducing diet. We are inclined to agree with Simpson that there is an underlying abnormality of adrenal cortical function in these individuals which by present methods fails to be demonstrated. Such patients are not uncommon and their treatment remains a challenge.

FIG. 19.—Case C. L. Obesity.

In conclusion, it seems that when allowances are made for varying activity of corticotrophin preparations, and conditions are otherwise standardized, different patterns of response of the adrenal cortex may be discerned. In addition to the hyperactive response seen in certain patients with Cushing's syndrome there are those interesting ones with a low basal activity but potentially very responsive to stimulation. Some mechanism other than the enzymatic lesion of pre-pubertal virilism appears to control the relative production of corticoid and androgen, and this may be disturbed in varying conditions, including hirsutism.

The part played by central nervous control of the anterior pituitary through the hypothalamus must be acknowledged. Information on the interaction of hormones from other glands on the activity of the adrenal cortex is scanty and needs to be increased; probably the thyroid is particularly worthy of further study. The effects of metabolic derangement on the function of the central nervous system, including the mimicry of some manifestations of space-occupying lesions, is of practical importance. With the discovery of aldosterone the problem of differentiating the hormonal abnormalities in Cushing's syndrome and allied states has increased in complexity.

Summary

Relevant knowledge of compounds secreted by the adrenal cortex and their metabolism is reviewed. Experiments are discussed which give information about the action of corticotrophin in man. Difficulties in evaluating the response due to variations in the activity of different batches of corticotrophin are stressed. After making due allowance for these difficulties, it is possible to define a "spectrum" of response in individuals with various pathological entities. Particular attention is drawn to the response of patients with idiopathic hirsutism, and the effects are contrasted with those in hypopituitarism and normal childhood on the one hand, and in adrenal cortical virilism on the other. Comparisons are made between the effect of exogenous corticotrophin and the endogenous stimuli produced by surgical trauma.

A condition of "basal hypo-adrenal corticalism" is defined and its significance discussed. The possibility of synergic action between corticotrophin and a factor associated with growth hormone, gonadotrophins, and thyroid hormones is considered. There seems little doubt that thyroid hormones are of great importance in the responsiveness of the adrenals. Evidence concerning the existence of a potentiating factor to corticotrophin for the stimulation of adrenal androgens is shown to be worthy of consideration. A similarity in behaviour of certain adult male obese subjects to that of hirsute women is noted and contrasted with others bearing a superficial resemblance to patients with Cushing's syndrome.

Attention is drawn to clinical problems in relation to alterations of cerebral activity in patients with Addison's disease and hypopituitarism and to difficulties which arise when space-occupying intracerebral lesions may be present. A patient with long-standing encephalitis has been studied, with particular reference to the secondary endocrinological phenomena which ensued, and the significance of these is discussed. Recent advances and ideas in relation to the diagnostic problems in Cushing's syndrome and in patients with adrenal cortical tumours, particularly when hypocalcaemia and renal symptoms are present, are considered and supplemented with additional observations.

ADDENDUM.—Since this manuscript was prepared additional confirmatory evidence has been obtained in another

patient in support of the effects of tri-iodothyronine on corticotrophin stimulation in hypopituitarism.

I should like to thank those colleagues whose assistance made much of this work possible, and especially Drs. R. V. Brooks, B. E. Clayton, R. R. McSwiney, and I. H. Mills, and many house-physicians; Professor W. G. Barnard and his staff for assistance in morbid anatomy; Miss P. Seimssen and the nursing staff of Holden Metabolic Ward; and the many people who gave technical assistance. Lastly, my thanks are due to the Governors of St. Thomas's Hospital for some financial assistance from endowment funds.

REFERENCES

- Aber, G. M., Chandler, G. N., and Hartfall, S. J. (1954). *British Medical Journal*, **1**, 1.
- Addison, T. (1855). *On the Constitutional and Local Effects of Disease of the Suprarenal Capsules*. London.
- Albright, F., Smith, P. H., and Fraser, R. (1942). *Amer. J. med. Sci.*, **204**, 625.
- Apert, M. (1910). *Bull. Soc. Pédiat. Paris*, **12**, 501.
- Armstrong, C. N. (1955). *Lancet*, **1**, 1051.
- Arnold, A. (1954). *J. clin. Endocr.*, **14**, 859.
- Astwood, E. B., Raben, M. S., Payne, R. W., and Grady, A. B. (1951). *J. Amer. chem. Soc.*, **73**, 2969.
- Ayala, G. (1934). *Rev. neurol. (Paris)*, **41**, 975.
- Ayres, P. J., Simpson, S. A., and Tait, J. F. (1956). *J. Endocr. (Proc.)*. In press.
- Bartter, F. C., Albright, F., Forbes, A. P., Leaf, A., Dempsey, E., and Carroll, E. (1951). *J. clin. Invest.*, **30**, 237.
- Bauer, H. G. (1954). *J. clin. Endocr.*, **14**, 13.
- Bayliss, R. I. S. (1955). *British Medical Journal*, **1**, 495.
- and Steinbeck, A. W. (1953). *Biochem. J.*, **54**, 523.
- (1954). *British Medical Journal*, **1**, 486.
- Beierwaltes, W. H., and Bishop, R. C. (1954). *J. clin. Endocr.*, **14**, 928.
- Benoit, J., and Assenmacher, I. (1953). *Arch. Anat. micr.*, **42**, 334.
- Bishop, P. M. F. (1955). *Proc. roy. Soc. Med.*, **48**, 1032.
- Bongiovanni, A. M. (1954). *J. clin. Endocr.*, **14**, 541.
- Brooks, R. V. (1956). Unpublished.
- McSwiney, R. R., Mills, I. H., Prunty, P. T. G., Robertson, K. M., and Wood, F. J. Y. (1956). In preparation.
- Broster, L. R., Paterson, J., and Camber, B. (1953). *British Medical Journal*, **2**, 1288.
- and Vines, H. W. C. (1933). *The Adrenal Cortex*. Lewis, London.
- Bruce, H. M., Parkes, A. S., and Perry, W. L. M. (1952). *Lancet*, **1**, 790.
- Bulloch, W., and Sequeira, J. H. (1905). *Trans. path. Soc. Lond.*, **56**, 189.
- Burnett, R., Rukes, M., Kolb, F., and Forsham, P. H. (1954). *J. clin. Endocr.*, **14**, 775.
- Bush, I. E. (1953). *Ciba Foundation Colloquium on Endocrinology*, **7**, 210.
- Callow, N. H., Callow, R. K., and Emmens, C. W. (1938). *Biochem. J.*, **32**, 1312.
- (1940). *J. Endocr.*, **2**, 88.
- Callow, R. K. (1938). *Proc. roy. Soc. Med.*, **31**, 841.
- Cater, D. B., and Stack-Dunne, M. P. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 31.
- Caughey, J. E., and Garrod, O. (1954). *British Medical Journal*, **2**, 554.
- (1955). *Proc. roy. Soc. Med.*, **48**, 884.
- Christy, N. P., Wallace, E. Z., and Jailer, J. W. (1955). *J. clin. Invest.*, **34**, 899.
- Clayton, B. E. (1954). Unpublished.
- and Hammant, J. E. (1955a). *J. Endocr.*, **13**, v.
- (1955b). *Nature (Lond.)*, **176**, 401.
- and Prunty, F. T. G. (1951). *Analyst*, **76**, 474.
- Cleghorn, R. A. (1951). *Canad. med. Ass. J.*, **65**, 449.
- Collip, J. B., Anderson, E. M., and Thomson, D. L. (1933). *Lancet*, **2**, 347.
- Conn, J. W. (1955). *J. Lab. clin. Med.*, **45**, 3.
- and Louis, L. H. (1956). *Ann. intern. Med.*, **44**, 1.
- Cope, C. L., and Harrison, R. J. (1955). *British Medical Journal*, **2**, 457.
- Copeman, W. S. C., Savage, O., Bishop, P. M. F., Dodds, E. C., Kellie, A. E., Stewart, J. W., Glyn, J. H. H., Henly, A. A., and Tweed, J. M. (1952). *Ibid.*, **1**, 397.
- Crooke, A. C., and Callow, R. K. (1939). *Quart. J. Med.*, **8**, 233.
- Cushing, H. (1932). *Bull. Johns Hopk. Hosp.*, **50**, 137.
- Daughady, W. H., Jaffe, H., and Williams, R. H. (1948). *J. clin. Endocr.*, **8**, 166.
- Dedman, M. L., Farmer, T. H., Morris, P., and Morris, C. J. O. R. (1953). In *The Suprarenal Cortex*, by J. M. Yoffey, p. 19. Butterworth, London.
- Dixon, H. B. F. (1955). *Proc. roy. Soc. Med.*, **48**, 903.
- Dorfman, R. I. (1949). *Ann. N.Y. Acad. Sci.*, **50**, 551.
- (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 112.
- and Ungar, F. (1953). *Metabolism of Steroid Hormones*. Burgess Publ. Co., Minneapolis.
- Dustan, H. P., Mason, H. L., and Corcoran, A. C. (1953). *J. clin. Invest.*, **32**, 60.
- Eik-Nes, K., Sandberg, A. A., Nelson, D. H., Tyler, F. H., and Samuels, L. T. (1954). *Ibid.*, **33**, 1502.
- Engel, G. L., and Margolin, S. G. (1941). *Arch. Neurol. Psychiat. (Chicago)*, **45**, 881.
- Evans, H. M. (1923-4). *Harvey Lectures*, p. 212.
- Falbrriard, A., Muller, A. F., Neher, R., and Mach, R. S. (1955). *Schweiz. med. Wschr.*, **85**, 1218.
- Ferrazzini, V. F., Borth, R., and Mach, R. S. (1952). *Helv. paediat. Acta*, **7**, 60.
- Finkelstein, M., von Euw, J. V., and Reichstein, T. (1953). *Helv. chim. Acta*, **36**, 1266.
- Forbes, A., and Albright, F. (1951). *J. clin. Endocr.*, **11**, 926.
- Donaldson, E. C., Reifstein, E. C., and Albright, F. (1947). *Ibid.*, **7**, 264.
- Forsham, P. H., Raimondo, V. D., Island, D., Rinfret, A. P., and Orr, R. H. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 279.
- Renold, A. E., and Frawley, T. F. (1951). *J. clin. Endocr.*, **11**, 757.
- Foye, L. V., and Feichtmeir, T. V. (1955). *Amer. J. Med.*, **19**, 966.
- Fraser, R. W., Forbes, A. P., Albright, F., Sulkowitch, H., and Reifstein, E. C. (1941). *J. clin. Endocr.*, **1**, 234.
- Gaunt, R., Renzi, A. A., and Chart, J. J. (1955). *Ibid.*, **15**, 621.
- Geschwind, I. I., and Li, C. H. (1952). *Endocrinology*, **50**, 226.
- Gordon, D., Horwitt, B. N., and Segaloff, A. (1954). *J. clin. Endocr.*, **14**, 297.
- Gray, C. H. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 527.

- Harrington, C. R., and Barger, C. (1927). *Biochem. J.*, **21**, 169.
- Harris, G. W. (1955a). *Ciba Foundation Colloquium on Endocrinology*, **8**, 531.
- (1955b). *Neural Control of the Pituitary Gland*, p. 150. Arnold, London.
- Hartman, F. A., and Brownell, K. A. (1930). *Science*, **72**, 76.
- Hayes, E. H., and White, W. F. (1954). *Recent Progr. Hormone Res.*, **10**, 265.
- Hechter, O., Jacobsen, R. P., Jeanloz, R. W., Levy, H., Marshall, C. W., Pincus, G., and Schenker, V. (1950). *Arch. Biochem.*, **25**, 457.
- Hellman, L., Bradlow, H. L., Adesman, J., Fukushima, D. K., Kulp, J. L., and Gallagher, T. F. (1954). *J. clin. Invest.*, **33**, 1106.
- Henriques, S. B., Henriques, O. B., and Selye, H. (1949). *Endocrinology*, **45**, 153.
- Hill, R. T. (1948). *Ibid.*, **42**, 339.
- Hill, S. R., Reiss, R. J., Forsham, P. H., and Thorn, G. W. (1950). *J. clin. Endocr.*, **10**, 1375.
- Howard, R. P., Sniffen, R. C., Simmons, F. A., and Albright, F. (1950). *Ibid.*, **10**, 121.
- Jailer, J. W., Gold, J. J., and Wallace, E. Z. (1954). *Amer. J. Med.*, **16**, 340.
- Jenkins, D., Forsham, P. H., Laidlaw, J. C., Reddy, W. J., and Thorn, G. W. (1955). *Ibid.*, **18**, 3.
- Jepson, R. P., Jordan, A., and Levell, M. J. (1956). *Brit. J. Surg.*, **43**, 390.
- Kappas, A., Dobriner, K., and Gallagher, T. F. (1955). *J. clin. Invest.*, **34**, 1559.
- Kendall, E. C. (1949). *Ann. N.Y. Acad. Sci.*, **50**, 540.
- King, A. B. (1951). *Bull. Johns Hopk. Hosp.*, **89**, 339.
- Laidlaw, J. C., Reddy, W. J., Jenkins, D., Haydar, N. A., Renold, A., and Thorn, G. W. (1955). *New Engl. J. Med.*, **253**, 747.
- Lancet, 1955, **1**, 1167.
- Lerman, J. (1955a). In *The Thyroid*, by S. C. Werner, p. 595. Cassell, London.
- (1955b). *Ibid.*, p. 716.
- Li, C. H. (1953). In *The Suprarenal Cortex*, by J. M. Yoffey, p. 1. Butterworth, London.
- Evans, H. M., and Simpson, M. E. (1943). *J. biol. Chem.*, **149**, 413.
- Liddle, G. W., Island, D., Rinfret, A. P., and Forsham, P. H. (1954). *J. clin. Endocr.*, **14**, 839.
- Long, C. N. H. (1940). *Res. Publ. Ass. nerv. ment. Dis.*, **20**, 486.
- Luft, R. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 523.
- Mader, I. J., and Iseri, L. T. (1955). *Amer. J. Med.*, **19**, 976.
- Marine, D. (1930). *Amer. J. med. Sci.*, **180**, 767.
- Mason, H. L., and Engstrom, W. W. (1950). *Physiol. Rev.*, **30**, 321.
- Migeon, C. J., and Plager, J. E. (1955). *J. clin. Endocr.*, **15**, 702.
- Mills, I. H. (1955a). *Proc. roy. Soc. Med.*, **48**, 313.
- (1955b). Unpublished.
- Moore, F. D., Steenburg, R. W., Ball, M. R., Wilson, G. M., and Myrden, J. A. (1955). *Ann. Surg.*, **141**, 145.
- Moore, R. A., and Cushing, E. H. (1935). *Arch. Neurol. Psychiat. (Chicago)*, **34**, 828.
- Morris, C. J. O. R. (1952). *Ciba Foundation Colloquium on Endocrinology*, **3**, 372.
- and Williams, D. C. (1953). *Ibid.*, **7**, 261.
- Munson, P. L., Goetz, F. C., Laidlaw, J. C., Harrison, J. H., and Thorn, G. W. (1954). *J. clin. Endocr.*, **14**, 495.
- Murray, G. R. (1891). *British Medical Journal*, **2**, 796.
- Nabarro, J. D. N. (1954). *Lancet*, **2**, 1101.
- and Walker, G. (1955). *British Medical Journal*, **2**, 617.
- Neher, R., and Wettstein, A. (1955). *Acta endocr. (Kbh.)*, **18**, 386.
- Nelson, D. H., and Samuels, L. T. (1952). *J. clin. Endocr.*, **12**, 519.
- Norymberski, J. K., Stubbs, R. D., and West, H. F. (1953). *Lancet*, **1**, 1276.
- Parkes, A. S. (1945). *Physiol. Rev.*, **25**, 203.
- Paton, A., and Petch, C. P. (1954). *British Medical Journal*, **1**, 855.
- Patterson, J., McPhee, I. M., and Greenwood, A. W. (1942). *Ibid.*, **1**, 35.
- Perloff, W. H., Lasché, E. M., Nodine, J. H., Schneeberg, N. G., and Vieillard, C. B. (1954). *J. Amer. med. Ass.*, **155**, 1307.
- Peterson, R. E., Wynsgaarden, J. B., Guerra, S. L., Brodie, B. B., and Bunim, J. J. (1955). *J. clin. Invest.*, **34**, 1779.
- Pincus, G., and Romanoff, E. B. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 97.
- Plager, J. E., and Samuels, L. T. (1952). *Fed. Proc.*, **11**, 383.
- Plotz, C. M., Knowlton, A. I., and Ragan, C. (1952). *Amer. J. Med.*, **13**, 597.
- Pond, M. H. (1955). *Acta endocr. (Kbh.)*, **19**, 1.
- Porter, R. W. (1954). *Recent Progr. Hormone Res.*, **10**, 1.
- Prunty, F. T. G. (1944). M.D. Thesis, Cambridge.
- (1950). *J. clin. Path.*, **3**, 87.
- (1953). *J. Endocr.*, **9**, 244.
- Brooksbank, B. W. L., Clayton, B. E., and McSwiney, R. R. (1951). *Ibid.*, **7**, lxxv.
- Clayton, B. E., and McSwiney, R. R. (1951). *Ibid.*, **7**, xxii.
- and Mills, I. H. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 324.
- McSwiney, R. R., Mills, I. H., and Smith, M. A. (1954). *Lancet*, **2**, 620.
- Querido, A. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 524.
- Kassenaar, A. A. H., and Cats, A. (1955). *Ibid.*, **8**, 309.
- Reddy, W. J., Jenkins, D., and Thorn, G. W. (1952). *Metabolism*, **1**, 511.
- Reifenstein, E. C., Forbes, A. P., Albright, F., Donaldson, E., and Carroll, E. (1945). *J. clin. Invest.*, **24**, 416.
- Robertson, J. D., and Kirkpatrick, H. F. W. (1951). *Lancet*, **2**, 54.
- Rolleston, H. D. (1936). *The Endocrine Organs in Health and Disease*. Oxford Univ. Press, London.
- Rosenberg, I. N., Cleroux, A. P., Raben, M. S., Payne, R. W., and Astwood, E. B. (1951). *Arch. intern. Med.*, **88**, 211.
- Rummell, P. M., Baer, L. S., Hollister, L., and Kolb, F. (1951). *J. clin. Endocr.*, **11**, 872.
- Sampson, H. (1897). *Phil. Trans. B*, **19**, 80.
- Sandberg, A., Eik-Nes, K., Samuels, L. T., and Tyler, F. H. (1954). *J. clin. Invest.*, **33**, 1509.
- Sarett, L. H. (1946). *J. biol. Chem.*, **162**, 601.
- Sayers, G. (1950). *Physiol. Rev.*, **30**, 241.
- White, A., and Long, C. N. H. (1943). *J. biol. Chem.*, **149**, 425.
- Sayers, M. A., Sayers, G., and Woodbury, L. A. (1948). *Endocrinology*, **42**, 379.
- Sheehan, H. L., and Summers, V. K. (1949). *Quart. J. Med.*, **18**, 319.
- (1952). *British Medical Journal*, **1**, 1214.
- Simpson, S. A., Tait, J. F., Wettstein, A., Neher, R., von Euw, J., Schindler, O., and Reichstein, T. (1954). *Helv. chim. Acta*, **37**, 1163.
- Simpson, S. L. (1951). *Proc. roy. Soc. Med.*, **44**, 453.
- (1953). *Ibid.*, **46**, 39.
- Smith, P. E. (1930). *Amer. J. Anat.*, **45**, 205.
- Soffer, L. J., Eisenberg, J., Iannaccone, A., and Gabrilove, J. L. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 487.
- Sprague, R. G. (1953). *Proc. roy. Soc. Med.*, **46**, 1070.
- Hayles, A. B., Power, M. H., Mason, H. L., and Bennett, W. A. (1950). *J. clin. Endocr.*, **10**, 289.
- Mason, H. L., and Power, M. H. (1951). *Recent Progr. Hormone Res.*, **6**, 315.
- and Power, M. H. (1953). *J. Lancet*, **73**, 217.
- Stack-Dunne, M., and Young, F. G. (1951). *J. Endocr.*, **7**, lxxi.
- (1954). *Ann. Rev. Biochem.*, **23**, 405.
- Steiger, M., and Reichstein, T. (1937). *Helv. chim. Acta*, **20**, 817, 1164.
- Stolz, F. (1904). *Ber. dtsch. chem. Ges.*, **37**, 4149.
- Storrie, V. M. (1955). *Bull. New Engl. med. Cent.*, **1**, 229.
- Summers, V. K., and Sheehan, H. L. (1951). *British Medical Journal*, **2**, 564.
- Swingl, W. W., and Pfiffner, J. J. (1930). *Science*, **72**, 75.
- Sydney, K. L., Kelley, V. C., Raile, R. B., Ely, R. S., and Sayers, G. (1953). *Proc. Soc. exp. Biol. (N.Y.)*, **82**, 695.
- and Sayers, G. (1952). *J. clin. Endocr.*, **12**, 920.
- Talbot, N. B., Butler, A. M., and Berman, R. A. (1942). *J. clin. Invest.*, **21**, 559.
- Sobel, E. H., McArthur, J. W., and Crawford, J. D. (1952). *Functional Endocrinology from Birth through Adolescence*. Harvard Univ. Press, Cambridge, Mass.
- Thompson, R. E., and Fisher, J. D. (1953). *Endocrinology*, **52**, 496.
- Thorn, G. W., and Clinton, M. (1943). *J. clin. Endocr.*, **3**, 335.
- Dorrance, S. S., and Day, E. (1942). *Ann. intern. Med.*, **16**, 1053.
- and Forsham, P. H. (1949). *Recent Progr. Hormone Res.*, **4**, 229.
- Renold, A. E., Morse, W. I., Goldfien, A., and Reddy, W. J. (1955). *Ann. intern. Med.*, **43**, 979.
- Venning, E. H. (1945). *Josiah Macy Jr. Foundation Conference on Metabolic Aspects of Convalescence*, October 15-16, p. 133.
- (1949). *Ann. N.Y. Acad. Sci.*, **50**, 553.
- Pattee, C. J., McCall, F., and Browne, J. S. L. (1952). *J. clin. Endocr.*, **12**, 1409.
- Wendler, N. L., Graber, R. P., Jones, R. E., and Tishler, M. (1950). *J. Amer. chem. Soc.*, **72**, 5793.
- West, H. F. (1954). *Ann. rheum. Dis.*, **13**, 56.
- (1956). *Ibid.*, **15**, 124.
- Wettstein, A. (1954). *Experientia (Basel)*, **10**, 397.
- Whittaker, S. R. F., and Whitehead, T. P. (1954). *British Medical Journal*, **2**, 265.
- Wilkins, L. (1955). *Ciba Foundation Colloquium on Endocrinology*, **8**, 523.
- Bongiovanni, A. M., Clayton, G. W., Grumbach, M. M., and van Wyk, J. (1955). *Ibid.*, **8**, 460.
- Willis, R. A. (1952). *The Spread of Tumours in the Human Body*. Butterworth, London.
- Wilson, H., Lovelace, J. R., and Hardy, J. D. (1955). *Ann. Surg.*, **141**, 175.
- Wolfson, W. Q., Beierwaltes, W. H., Robinson, W. D., Duff, I. F., Jones, J. R., Knorpp, C. T., Siemienski, J. S., and Eya, M. (1951). *Proceedings of the Second Clinical ACTH Conference*, **2**, 95. Churchill, London.

EFFECTS OF HEAVY AND REPEATED MALARIAL INFECTIONS ON GAMBIAN INFANTS AND CHILDREN

EFFECTS OF ERYTHROCYTIC PARASITIZATION

BY

IAN A. MCGREGOR, L.R.C.P.&S.Ed., D.T.M.&H.
H. M. GILLES, M.D., B.Sc., D.T.M.&H.
J. H. WALTERS, M.D., M.R.C.P.
A. H. DAVIES, M.B., Ch.B.

AND

F. A. PEARSON, B.Sc., M.R.C.S., D.T.M.

From the Medical Research Council Laboratories,
Gambia, British West Africa

Since 1947, when the Medical Research Council established its laboratories in the Gambia, many workers investigating the importance of non-parasitic diseases in the African population have found great difficulty in assessing the contribution to the general picture of the inevitable malarial infection. McGregor and Smith (1952) and Walters and Waterlow (1954) found the twin threads of parasitization and malnutrition exceedingly complex and difficult to unravel. In no section of the community is this complication more in evidence than in infancy and early childhood—when plasmodial infection is severe and almost universal. This paper describes the result of an investigation in the Sukuta area of the Gambia that was undertaken to determine precisely the effects, both immediate and long-term, of repeated and heavy malarial infections on infants and young children.

Sukuta is one of the principal villages in the Kombo St. Mary District of the Gambia, and as such it possesses