A great deal of trouble and anxiety, not only to the patient, relatives, and doctor, but also the hospital staff, could be saved if we can get a higher proportion earlier.

Summary and Conclusions

An account is given of a series of cases of acute appendicitis treated conservatively with some illustrative cases.

During the course of over 30 years of surgery, a good deal of which has been emergency surgery, I have gradually been tending more and more to conservative treatment in cases of advanced appendicitis.

For many years I have believed it best to treat appendix abscesses conservatively. For more than four years I have believed it best to treat all cases of acute appendicitis over 24 hours old conservatively.

It is probably wise to treat all cases of acute appendicitis under 24 hours old by an emergency appendicectomy, and this is our usual custom.

One sometimes wonders whether it would not be a sound procedure to treat all cases of acute appendicitis conservatively. They seem to settle down quite nicely, and some never seem to have any further trouble: the appendix has wizened.

We should then be left with appendicectomy for recurrent acute appendicitis, and for chronic appendicitis—the “grumblers” with faecaliths in the appendix, with kinks, and with adhesions.

Looking into the future, one cannot help feeling that our successors will be more conservative in their outlook in this matter, and may look back on us as having been too “appendicectomy-minded.”

I acknowledge the assistance of my three registrars, Mr. K. Drummond, Mr. R. Kotariya, and Mr. R. Mansharamani, without whose help, skill, and co-operation this work would not have been possible.

EXFOLIATIVE CYTOLOGY IN THE DIAGNOSIS OF LUNG CANCER EXAMINATION OF ONE LABORATORY'S RESULTS

BY

F. HAMPSON, D.M.

From the Department of Pathology, Grimsby General Hospital

Although much of the early work on the recognition of cells exfoliated from tumours was done in this country, little use is made here to-day of the techniques of exfoliative cytology in the diagnosis of cancer, though there have been signs recently of a renewed interest in the subject (Osborn, 1953; Philips, 1954).

After I had spent several months in Dr. Papanicolaou's department a service in cytology was offered in this laboratory for some four and a half years. The majority of requests have been for the examination of sputum. A review of the results obtained might be of some value in assessing whether exfoliative cytology has any contribution to make towards the diagnosis of lung cancer.

Material and Method

Specimens of sputum have been sent to the laboratory from a wide area, most of them coming from chest clinics and chest hospitals. In 1955, for example, 567 specimens out of a total of 855 arrived from such

sources. The remainder were from general hospitals and general practitioners.

All specimens, whether sent to the laboratory by post or not, had been collected by getting the patient to cough directly into a jar containing 70% alcohol. This routine, which is necessary only for specimens that cannot be examined at once, has proved very satisfactory in practice. Any possible slight loss in quality has been preferred to the risk of specimens being spoilt altogether by delay in sending or examining material not so treated.

It is not intended to describe the laboratory procedures that have been followed. No new methods have been developed, and the techniques and the reporting of results have followed closely those described by Papanicolaou (1954). Specimens have been placed into one of five classes: (1) absence of atypical or abnormal cells; (2) atypical cytology but no evidence of malignancy; (3) cytology suggestive of, but not conclusive for, malignancy; (4) cytology strongly suggestive of malignancy; and (5) cytology conclusive for malignancy. For the purpose of this report, specimens which had been reported originally as class 3 have been called “suspicious” and those which had been placed in class 4 or 5 have been called “positive.”

Results

Table I shows the outline of the results obtained for the years 1952–5. For the years 1952, 1953, and 1954 the number of specimens examined, the number of patients from whom they came, and the number and percentage of patients diagnosed cytologically as suspicious and positive are given.

TABLE I

Year	Total Specimens	Total Patients	Suspicious or Positive	
			No.	%
1952	200	107	10	9.4
1953	444	232	27	11.6
1954	743	329	44	13.4
1955	855	351	42	11.9

For the year 1955 a more detailed analysis is possible. A request for information was sent to the doctors concerned in respect of the 42 patients diagnosed as suspicious or positive and full information was obtained about 40 of them. The smears from 23 of these 40 patients were reported as positive and from 17 as suspicious (Table II).

TABLE II

	Final Diagnosis of Cancer	Final Diagnosis Not Cancer	Histological Evidence of Cancer	Cancer Cases with Positive Smears and Negative Biopsies
Positive group (23 cases)	23	0	11	2
Suspicious group (17 cases)	12	4 and ?1	5	2

During 1955 all cases from which positive smears were obtained were finally diagnosed as cancer. Histological confirmation in all cases was not available. Several patients were too ill for further investigations and bronchoscopy was refused in one case. The final diagnosis of cancer was taken from the opinion expressed by the doctor concerned in his answers to the questionnaire. In some cases, therefore, the final diagnosis of cancer rests on clinical findings, plus radiology, smears, and histology; in others, one or more of these pieces of evidence will be lacking, but in no case was the doctor in any doubt about the diagnosis. In the suspicious group it is seen that four

cases for certain and a possible fifth turned out to be non-malignant, whilst 12 were confirmed as malignant either by histology or on clinical grounds, or both. In 10 of these 40 cases the doctor concerned said that the smear result had been the first objective evidence of malignancy. The series includes two cases reported as positive where a biopsy was negative; two cases reported as suspicious also had negative biopsies. In one of these four cases operation showed an epidermoid carcinoma. In all the others the diagnosis was of inoperable carcinoma, and two patients have died whilst one was alive six months after diagnosis was made.

Discussion

The results obtained over the past few years have shown that exfoliative cytology is of real value in the diagnosis of lung cancer. It is not suggested that it should replace any of the existing diagnostic methods but that it is a most useful additional aid. In the first place, the method is simple and practicable for patient and clinician. No preparation of the patient is necessary, he does not have to attend hospital, and additional specimens can be readily obtained. For the laboratory the method has the advantage that no elaborate equipment is required, although it is certainly true that experience is needed before confident results can be given.

It has been found that reliance can be placed on positive results. In 1955 all of 23 cases reported positive were finally diagnosed as malignant, and in no case was any doubt expressed on this point by the clinician concerned. In four of the 17 suspicious cases the clinician was satisfied that the diagnosis was non-malignant, and in a fifth case the patient died before a firm diagnosis was made.

It is clear, therefore, that if a confident diagnosis of malignancy is made on cytological grounds there is a very high probability of proving malignancy, and that most of the cases regarded as suspicious will also prove to be malignant. Further experience will probably reduce the number of specimens placed in this category. Negative results are of less significance. It has not been possible to ascertain the number of malignant cases on which negative results were returned, but there were undoubtedly instances of this. Apart from inadequate examination or inaccurate appraisal of cells seen, negative smears may be found in cancer cases for several reasons: it may be that no tumour cells were being exfoliated at the time of the examination; the tumour may not communicate with a bronchus, or, even if it does, the bronchus itself may have become obstructed by pressure so that no escape of cells can take place (Osborn, 1953).

The results presented and the answers to the questions asked of the clinicians show that the method is of value in the early diagnosis of cancer, and this is of especial importance in this disease. In 1955, 40 cases were regarded as positive or suspicious, and in 10 of these the clinician stated that the positive report issued after cytological examination constituted the first objective evidence of malignancy. It is here that the extreme simplicity of the method—from the patients' point of view—is so important. It is a simpler matter to arrange for cytological examinations of sputum than to arrange for other special examinations to be performed, and it seems reasonable that earlier recourse should be had to cytological methods for that reason. It is undoubtedly true that small early tumours produce recognizable exfoliated cells, and "carcinoma *in situ*" of the bronchus has been diagnosed by cytological methods (Papanicolaou and Koprowska, 1951). In the case described by these authors it is stressed that positive cytological findings were repeatedly obtained in spite of negative biopsies, and the diagnosis of carcinoma *in situ* was finally established at the post-mortem examination following death from pneumonia.

It is difficult in such a small series to attempt to compare the value of cytological diagnosis with that of other special methods of examination. However, even among these few cases two reported as positive and two as suspicious were

negative according to the biopsies. These four cases were subsequently regarded as undoubtedly malignant. In one case the diagnosis was proved at operation; the other three cases were found to be inoperable. Herbut (1951) referred to 500 cases of lung cancer in which cytological methods gave positive results in 88% and bronchoscopic examination a positive result in only 31%; in 29% of the proved cases cytology gave positive results and bronchoscopy was negative. Again, Benioff (1951), referring to 162 proved cases that came to operation, showed that in 24% of these cytology provided the only pre-operative evidence of malignancy. The conclusion seems justified that cytological methods will disclose more positive cases than will bronchoscopy.

The objection is sometimes raised that it is a time-consuming method of investigation. This is true, but it is also true of many other laboratory procedures which, nevertheless, are in common use. Over a four-year period more than 10% of patients from whom specimens have been sent to this laboratory have been regarded as showing definite evidence of malignancy or as affording strong grounds for suspicion. This compares well with the percentage of positive material obtained in many other laboratory investigations. The actual examination of the slides takes somewhat longer to perform than does an examination for tubercle bacilli; and here, where the preliminary screening is done by an experienced technician, some seven to eight minutes is spent in examining a negative smear.

Summary

The incidence of "positive" and "suspicious" smears from sputum suspected of containing malignant cells is surveyed for the years 1952–5.

For the year 1955 the results are followed in greater detail and the accuracy of the cytological diagnoses is assessed.

It is concluded that the method is of great value in the diagnosis of lung cancer, and especially in early cases; that the positive results are reliable; and that in accuracy it compares favourably with bronchoscopic examination.

The method is extremely convenient for the patient and clinician, and demands no elaborate laboratory equipment.

My thanks are due to Mr. E. C. Darnell for expert technical assistance and to all those clinicians who provided information about their cases.

REFERENCES

- Benioff, M. A. (1951). Symposium on Exfoliative Cytology, New York City. Abstracts of papers. (Unpublished.)
- Herbut, P. A. (1951). Ibid.
- Osborn, G. R. (1953). *Applied Cytology*. Butterworth, London.
- Papanicolaou, G. N. (1954). *Atlas of Exfoliative Cytology*. Commonwealth Fund, Harvard.
- and Koprowska, I. (1951). *Cancer*, 4, 141.
- Philips, F. R. (1954). *Brit. J. Cancer*, 8, 67.

A medical manuscript written very legibly in Anglo-Saxon before the Norman Conquest was bought for £550 at Sotheby's on December 11 by the Wellcome Trust. Although the text covers only one side of a single leaf, there was brisk bidding from both English and American buyers. The manuscript contains recipes for making medicines and salves for "heartache," disease of the lungs and the liver, and for tumours. It was incorporated in the binding of a book printed in Antwerp in 1558 which later came into the library of Lord Robartes. This fragment must have belonged to the vast quantity of mediaeval manuscripts which were thrown out of the monasteries at the time of the Dissolution and used for all kinds of menial purposes. As Bishop Bale wrote in 1549, "Some they sold to the grossers and sopesellers, and some over see to the bokebynders, not in small nombre, but at tymes whole shyppes fulle, to the wonderynge of the foren nacyons."