REFRESHER COURSE FOR GENERAL PRACTITIONERS

THE DIAGNOSIS OF EARLY PHTHISIS

BY

JAMES L. LIVINGSTONE, M.D., F.R.C.P.

Physician, King's College Hospital and Brompton Hospital

On the early diagnosis of pulmonary tuberculosis depends not only the prognosis of the patient but also the prevention of spread of infection to others. Mass radiography in this and other countries has shown that at least four in every thousand of the apparently healthy young adult population examined for military service are suffering from this disease, and the figure is probably higher in the older age groups.

First Principles

The following principles should be borne in mind.

(1) In every case of phthisis there is a stage in development of the disease when there are no clinical symptoms or signs whatever.

(2) The culture of *Mycobacterium tuberculosis* from sputum, laryngeal swab, or gastric lavage is the most delicate and certain method of diagnosis, and the presence of *M. tuberculosis* indicates "active" disease.

(3) The medical man should not only suspect but take active steps to exclude pulmonary tuberculosis in all contacts with an "open" case, and in patients who have had primary pleurisy (with or without effusion), erythema nodosum, tuberculous adenitis, or other symptoms of a primary lesion.

(4) When one or more cases of "primary" tuberculosis are found in the home, school, office, etc., a careful search should be made for the "open" case who has been acting as a carrier, and who is usually entirely unaware of the diagnosis.

(5) Haemoptysis never comes from a "small vein in the pharynx."

When the first or asymptomatic stage of the disease proceeds to the second or "active" stage the progress is usually slow and insidious over several months, during which time the lung lesion enlarges, cavitates, or spreads. In those fortunate patients whose lesion involves the pleura and causes pain, or breaks through into a bronchus to produce haemoptysis, the diagnosis may be made earlier, for these symptoms take the sufferer to the doctor. In others the lesions may be widespread before toxæmic or pulmonary symptoms are marked. The onset of the second stage may be acute, particularly in young adolescents and in races with a poor resistance to the disease. In such cases there is extensive bronchogenic spread on one or both sides and areas of atelectases from tuberculous disease of the bronchi.

Clinical Examination

In the investigation of a suspected case of early phthisis a careful history must be taken, which should include the family history and the possibility of contact with "open" cases.

Symptoms due to Toxaemia.—(1) Lassitude, particularly in relation to the week-end amusements—e.g., hiking, football, or dancing. (2) Loss of appetite, fullness after meals, and "dyspepsia." (3) Menstrual irregularities, especially oligomenorrhoea. (4) Loss of weight. (5) Abnormal sweating.

Symptoms due to the Pulmonary Disease.—(1) Stained sputum or haemoptysis. (2) A "cold in the head and chest" followed by a persistent productive cough. (3) Vague chest pains. (4) Huskiness. (5) Persistent wheezing on one side. (6) Dyspnoea on active exertion.

Physical Signs.—(1) Pallor, often visible on the palate. (2) Early clubbing of the fingers, as shown by a lilac colour of the nail-bed and shiny thickening of the soft tissues at the base of the nails. (3) Slight diminution of movement and air entry on one side and post-tussive rales in the upper zones, especially posteriorly. (4) Occasionally, unilateral persistent wheezing may indicate tuberculous bronchitis. (5) On laryngoscopy, pallor and swelling of the aryepiglottic folds may be present.

Gross physical signs may be found in more advanced disease, but there is often a remarkable absence of signs in the presence of extensive radiological disease. When a patient has been in bed for a week or two the auscultatory signs frequently disappear completely.

Radiological Examination

A radiograph of good quality is an essential part of the investigation. If opacities are seen in the middle or lower zones a lateral film is helpful both in showing the anatomy of the lesions and as a standard for future comparison.

It must be remembered, particularly in "contacts," that a minimal lesion may not be visible soon after exposure to infection, and that subsequent films at intervals of one month and then three months may be required in some cases before excluding the diagnosis. An opacity caused by a non-tuberculous infection usually regresses or disappears within one month, whereas a tuberculous lesion resolves more slowly.

The radiological appearances in pulmonary tuberculosis vary according to the type and extent, but four characteristic lesions may be seen in the early case, alone or in combination.

(1) Discrete round foci—Assmann's foci—which are seen in the upper part of the lung fields as sharply defined round or oval shadows from 5 to 25 mm. in diameter; there is no enlargement of the related hilar lymph nodes.

(2) Early infiltration—"Frühinfiltrat." This is shown by fine mottling consisting of a group of comma-shaped opacities, usually in the upper or mid zones.

(3) Cavitation.

(4) Areas of segmental collapse of part of a lobe, usually showing as dense triangular opacities with the base towards the chest wall and along the lines of the interlobar fissures. This appearance is associated with tuberculous disease of the bronchi.

Bacteriological Examination

If sputum is present three separate morning specimens should be examined by stained smear for *M. tuberculosis*. If this examination is negative a twenty-four-hours collection should be sent to the laboratory for culture of

FIG. 1. — Assmann's focus in the right first space, 20 mm. in diameter.

M. tuberculosis. The result will not be known for two to six weeks. Every laboratory should be equipped to undertake culture of this organism.

If sputum is not being produced there are two methods of obtaining specimens for culture.

Laryngeal Swab Culture.—The doctor should wear a face-mask. An ordinary throat swab is moistened with normal saline and the wire is bent to about 140 degrees with sterile forceps. The tongue is depressed with a spatula and the patient made to breathe deeply through the mouth; the swab is placed behind the epiglottis and the patient told

FIG. 2.—Early infiltration in the right first space.

History.—Female aged 18. Contact with father for two years. No symptoms. Normal temperature and pulse rate and normal B.S.R. Went through routine course of sanatorium treatment for six months with little change in the x-ray.

FIG. 3.—Infiltration in right upper and middle zones with atelectasis above the inter-lobar fissure. Early infiltration left mid-zone.

History.—Female aged 18. December, 1948: dry cough with occasional greenish sputum. February, 1949: feverish cold with increased cough. March: loss of appetite and nausea. Developed jaundice and was in bed for five weeks. April: cough increased. Temperature 104° F. (40° C.). Given sulphonamides. A few days later: haemoptysis. Diagnosed as broncho-pneumonia. Referred to chest clinic in May: sputum positive.

to cough. If he is unable to do so, the swab is pushed downwards in the direction of the glottis, and this often provokes a cough reflex. The wire is straightened with the forceps and the throat swab replaced in the test-tube. It is advisable to take two laryngeal swabs, which are sent to the laboratory with a request for culture of *M. tuberculosis*.

FIG. 4.—Infiltration and cavitation at the left apex, with calcified scars at the right apex.

History.—Female aged 21. 1948: several colds, slight lassitude, and loss of weight. May, 1949: persistent cough with trace of greenish sputum; T.B.-positive. Family history: negative. Contact: nil.

Gastric Lavage.—This can be done perfectly well in the patient's home or doctor's surgery by the district nurse or anyone with a little practice. Obviously it is essential to do it soon after the patient wakes in the morning and before anything has been taken by mouth. A Ryle's tube is swallowed, and some 60 ml. of saline introduced into the stomach and sucked back with the syringe. The stomach washings are sent to the laboratory for culture and direct-smear examination. As rubber and glassware are difficult to free from *M. tuberculosis* despite careful cleansing and boiling, it occasionally happens that the direct smear reveals bacilli not obtained from the patient. Confirmation by culture should therefore always be done.

These investigations may prove the diagnosis of "active" pulmonary tuberculosis within a few days or up to a month, and it must be stressed that the presence of bacilli or cavitation indicates "active" disease in spite of the absence of symptoms or signs. During the "active" stage progress of the disease may be rapid and insidious, and the essential treatment is immediate bed rest for at least three months, during which time the physician can assess progress and decide on collapse therapy or other methods of treatment.

Unless strict instructions are given the patients rarely have proper bed rest. By this is meant lying in bed with two pillows and being allowed up for toilet and bath once a day; a chamber-pot or bed-bottle should be used for the bladder. The temperature and pulse rate should be recorded twice daily on a chart.

Is the Lesion "Active" or Not?

The radiograph may show an opacity though the bacteriological findings are negative, and the clinician is then faced with two problems: (i) What is the cause of the lesion? (ii) If it be tuberculous, is the lesion "active," quiescent, or healed?

The following routine should be carried out.

(1) The clinical examination should be repeated and the patient instructed to keep a temperature-and-pulse chart, while leading a normal life, and to record his weight every week.

(2) A further chest film should be taken a month after the first one. If the original opacity has markedly decreased or disappeared it is probably due to a non-tuberculous infection; if stationary or increased in size it is more likely to be due to tuberculosis or neoplasm, and cavitation may take place in both these conditions.

(3) The bacteriological examination should be repeated.

(4) The erythrocyte sedimentation rate may be raised, indicating toxic absorption.

(5) If the radiograph suggests segmental atelectasis a bronchoscopy should be considered.

(6) In patients under 35 years of age a tuberculin-jelly test may be done: if negative, the Mantoux test (1/100 strength O.T.) should be carried out. Should this be negative, it is strong evidence against the lesion being tuberculous.

When these examinations prove negative it may be assumed that the disease is inactive, but the patient should be kept under observation and serial films taken of the chest every three months for at least one year. Any progression, "softening," or cavitation in the lesions is an indication for a further search for *M. tuberculosis*: tomography is helpful when there is doubt whether cavitation is taking place.

In persons over 40 an early bronchial carcinoma may cause dyspnoea, haemoptysis, or other symptoms simulating phthisis. Until the tumour is big enough to block a bronchus the chest radiograph will show no abnormality. This is an important point, and a negative film should not deter the physician from requesting a bronchoscopy when symptoms are suggestive.

The early diagnosis of phthisis calls for the doctor's clinical acumen and knowledge of his patient's living conditions. The investigations required are relatively simple, and should be available to every general practitioner. The culture of secretions for *M. tuberculosis* is a much more delicate method than the direct smears. When bacilli are found the diagnosis of "active" phthisis is established, and the patient should be put to bed forthwith. Further assessment and treatment call for the opinion of a specialist.

ROCKEFELLER FOUNDATION INTERNATIONAL HEALTH DIVISION

The International Health Division of the Rockefeller Foundation continues its world-wide assistance to public health projects. The report, just issued, for 1948 describes work on the control of malaria, yellow fever, and other virus diseases in both hemispheres.

World-wide interest has been aroused by the anopheline eradication campaign in Sardinia, which is a combined project of the Italian Government and the Foundation. It is described as a comprehensive effort to push forward the frontiers of public health practice. Sardinia has long been highly malarious, and an attempt has been made to eliminate *Anopheles labranchiae* by house-spraying and larvicidal campaigns. By the end of 1948, after a little more than two years' continuous work, considerable progress had been made towards species eradication.

The next phase, which may be continued through 1950, calls for an intensive search for surviving adult mosquitoes and larvae, together with strict control and re-treatment of any zones harbouring them. Effective quarantine measures are also being instituted to guard against the risk of reinfestation from the island of Corsica, ten miles distant, where, incidentally, the French Government is organizing a control project, again with the help of the Foundation. In the house-spraying operations in Sardinia a plant has been used which is capable of turning out 561,000 litres of D.D.T. solutions a month.

Comment is made in the report on the excellent morale of the large labour force enlisted locally. Men from all parts of the island who were engineers, butchers, barbers, soldiers, or casual labourers have come into an organization totally strange to them in method and purpose and have given it their energy and good will to a degree far beyond that demanded by an ordinary conscientious application to duty.

Yellow Fever and Q Fever

The yellow fever research institutes at Lagos, in Nigeria, and at Entebbe, in Uganda, have been hard at work. At Lagos research has been continued on the "scratch vaccine" recently developed by French workers from the dried brains of mice infected with a neurotropic strain of yellow fever virus. The vaccine has been tested out in the little Nigerian town of Kumba, where the entire population of some 4,000 has been successfully vaccinated by the new technique. There were no serious reactions, and an immunity rate to yellow fever of approximately 92% was achieved.

In Uganda research on the epidemiology of yellow fever has continued. The identification of the principal insect vectors and vertebrate hosts has been further studied to determine how the virus is maintained. Other work on yellow fever has been undertaken in Central and South American countries.

Reference is made to Q fever, which was first recognized as a disease affecting man in Queensland, and which is now known to be endemic in widely scattered areas of the world, including