REFRESHER COURSE FOR GENERAL PRACTITIONERS

MEASLES

BY

J. V. ARMSTRONG, M.D., D.P.H.

Physician Superintendent, Brook General Hospital, Woolwich

Measles is still a serious disease in this country, particularly in young children up to 2 or 3 years of age. Although its virulence has declined in a remarkable fashion in the last twenty years and deaths have fallen to a record low level, it is still capable of causing extensive and permanent damage, especially to the lungs, middle ear, and eyes. Before this decline measles probably killed more children than any single infectious condition except enteritis, and though hygiene and better nourishment have had considerable effect in controlling the disease an increase in its virulence is always possible.

Aetiology

The causative virus gains access to the body through the respiratory tract by droplet infection from another sufferer. The disease is intensely infectious, especially in its early stages, and all ages are susceptible. Young infants are immune if their mothers have had the disease, and they do not lose this maternal immunity completely until they are about nine months old, after which time unmodified measles may be expected. If the mother has not had measles her infant will be susceptible from birth.

Like most virus diseases measles has a long incubation period, of from ten to twenty-one days, twelve days being the average for an unmodified attack; the immunity produced usually lasts for life and second attacks are rare. In modified measles the incubation period may be longer.

The measles virus seems to have the property of lowering the resistance of the respiratory mucous membrane and rendering it susceptible to attack by secondary bacterial invaders, particularly the streptococcus and pneumococcus.

Clinical Features

In spite of the rash being such a prominent feature measles should be regarded as essentially a respiratory disease in which the whole respiratory tract may be involved. In the prodromal stage, which usually lasts three to four days but may be longer, the upper respiratory tract bears the brunt of the attack. The condition then simulates an acute cold, and coughing, sneezing, and lacrimation, associated with general misery, are prominent features. It may sometimes be accompanied by a catarrhal laryngitis, which is often mistaken for laryngeal diphtheria. The temperature will almost certainly be raised and a marked degree of conjunctivitis may be present.

Koplik's Spots.—A careful inspection of the mouth, an essential part of the examination of any child whatever disease is suspected, will probably reveal the presence of Koplik's spots. These spots are pathognomonic of measles and resemble grains of salt on a reddened buccal mucosa. Sometimes they are so numerous as to become confluent in places. They are usually found opposite the molars and can be seen only in good daylight or with a lamp giving proper colour values. They are practically invisible with ordinary electric light, and the small pocket torch is useless. The spots rapidly clear on the appearance of the true rash (for this reason they are usually looked for

too late), though a reddened and granular appearance of the mucous membrane often persists for a short time. In addition, some degree of stomatitis is always present and the mouth may become foul.

Prodromal Rashes.—These occur occasionally and may take the form of a fine macular rash on the trunk suggesting rubella, or of a scarlatiniform eruption falsely indicating scarlet fever. The scarlatiniform type of prodromal rash is occasionally the cause of the double diagnosis of scarlet fever and measles being made. The two diseases do not usually occur together, though as previously stated the streptococcus is an early secondary invader in measles, particularly in the mouth, nose, and middle ear, and on the fifth or sixth day the patient may present a typical peeled streptococcal tongue as in scarlet fever.

Respiratory catarrh may occasionally be accompanied by appendicular catarrh and the signs of acute appendicitis. In such cases operative intervention may be necessary. I saw four examples last year, all of which came to operation.

The Rash.—The true rash of measles begins as a few fine macules behind the ears and rapidly spreads on to the face and then to the trunk and limbs, finally reaching the feet. The lesions are often papular as well as macular and coalesce to form irregular, dusky red, blotchy patches. Sometimes large areas of skin, especially on the trunk, are covered by a confluent eruption somewhat suggesting scarlet fever. On the other hand, the rash may remain macular and discrete, particularly in modified measles.

When the rash is very heavy it may contain obvious haemorrhagic elements. This is not necessarily a sign of a severe attack (true haemorrhagic measles is very rare), and in most patients with a heavy eruption staining due to altered blood pigment is left behind when the rash fades. A branny desquamation occurs also. With the appearance of the true rash the temperature, which usually falls to normal at the end of the prodromal period, rises again. In uncomplicated cases it falls in three to four days.

In ordinary cases the rash usually subsides by the end of a week; the patient rapidly becomes convalescent and may be allowed up. He may safely be regarded as free from infection by the end of ten days' illness and can be considered cured by the end of the second week.

Complications

Complications involving the respiratory tract are the most common and are generally due to secondary bacterial infection.

Otitis.—Infection may spread from the nasopharynx to cause otitis media, the most common complication of measles at the present time. It is usually suppurative, but with early treatment by chemotherapy is unlikely to be followed by mastoid disease. Deafness may occur.

Laryngitis.—Secondary laryngitis developing as the true rash fades is a much more serious condition than the early catarrhal laryngitis of the prodromal stage. It may become obstructive and is nearly always either diphtheritic

or streptococcal in origin; it must be treated as diphtheritic until proved otherwise.

Bronchitis.—This is a common and sometimes severe complication that may be followed by bronchopneumonia, often of the interstitial variety, which is particularly likely to give rise to subsequent fibrosis and bronchiectasis. Lobar pneumonia is not a usual complication in young children, and empyema after either form of pneumonia is at present rare.

Enteritis.—This complication is most likely in infants and may be a danger to life. It may be present in the early stages as a toxic manifestation, but if it persists is probably due to secondary infection. All the classical signs of infantile gastro-enteritis may be present, with dehydration and liver damage.

Eye Lesions.—Conjunctivitis is a normal accompaniment of measles, but corneal ulceration may occur very rapidly, particularly in debilitated children, and have serious consequences. Opacities that seriously impair sight may result, and perforation of the anterior chamber and disorganization of the eye have been known to follow in the worst cases.

Encephalitis.—Post-infective encephalitis may occur in measles as in other virus diseases. It usually shows itself at the end of the first week. The damage may be minimal, but in anything up to 30% of cases it is severe enough to cause death. At least half of the cases that survive seem to make a complete recovery, but the remainder may develop epileptic fits and mental deterioration. In the early stages meningeal irritation may be present, with irritability or convulsions. Stupor or complete unconsciousness may supervene, and during the slow recovery stage spasticity of the limbs is a pronounced feature. Myelitis may coexist.

Associated Diseases.—Measles can be deadly in association with other diseases. Laryngeal diphtheria has already been mentioned, and the combination of measles and whooping-cough greatly increases the risk of severe pneumonia and subsequent fibrosis. Tuberculosis of the lungs is not so common after measles as was once thought, although it may be lit up by a bad attack of pneumonia in a previously infected subject.

Diagnosis

A history of exposure to measles should always be sought. The combination of pyrexia, catarrhal signs, and a generalized morbilliform rash is enough strongly to suggest measles, and if Koplik's spots are seen the diagnosis is established. In the prodromal stage the disease has to be distinguished from the ordinary cold, from laryngeal diphtheria, and, more rarely, from the catarrhal stage of whooping-cough. In the latter disease catarrhal signs, including laryngitis, may be pronounced. The prodromal rashes have already been mentioned as suggesting rubella or scarlet fever, but confusion with these two diseases may also occur when the true rash appears. The rash of rubella is pink, however, and the macules are smaller and usually discrete. In addition all the superficial lymph glands are enlarged, including the occipital, post-cervical, and, most typical of all, the mastoid.

Scarlet fever should not be confused with measles, as there is no true rash on the face. In some cases the erythema of scarlet fever may be blotchy enough on the limbs to suggest measles, although evenly spread on the trunk. The tongue may be similar in both diseases, but catarrhal signs and stomatitis will be lacking in scarlet fever.

Smallpox may be confused with measles at two stages of its development. There may be a morbilliform prodromal

rash, but it is the papular stage of the true rash, particularly when it is profuse on the face, which causes the greatest difficulty. The eruption of smallpox is more shotty and it is not associated with catarrhal signs or fading Koplik's spots; its distribution is different, for the rash shows a predilection for the extremities and tends to be scanty on the trunk. The subsequent development of papules into vesicles settles the issue.

The rose rash of the enteric group, which usually appears about the sixth day of disease, is sometimes profuse enough, particularly in paratyphoid fever, to suggest measles or rubella. The slow onset, the typical "staircase" rise of temperature, and the absence of catarrhal signs are against measles, and blood culture followed later by stool culture and Widal tests will confirm the diagnosis of enteric fever.

Another rash which is occasionally mistaken for measles is the haemorrhagic eruption of cerebrospinal fever, if it is at all profuse. This disease has a fulminating onset and there is usually early development of meningeal signs.

The rash produced by the sulphonamide group of drugs may bear a close resemblance to measles. Koplik's spots will be absent, however, and the eyes are usually clear. A history of taking the drug can usually be obtained, and such a history should always be sought in any doubtful case of measles. The barbiturates may give morbilliform rashes; penicillin rashes are usually urticarial. Serum rashes resembling measles are seldom seen since the production of refined antitoxin, but sensitization to other foreign proteins may produce rashes of morbilliform type. These include areas of circinate or marginate erythema that are never seen in measles. Food rashes are either of this nature or urticarial.

Treatment

There is no specific treatment for measles, but patients with mild attacks recover quickly if kept warm in bed for a few days. Plenty of fresh air must be insisted upon. The diet in the early stages should consist mainly of milk with as much bland fluid, such as sweetened fruit drinks, as the child can be persuaded to take. A simple linctus may be necessary to control the cough if troublesome.

In the more severe cases, particularly if home conditions are not good, hospital treatment should be sought early. In such cases bacterial complications may be expected and a prophylactic course of a sulphonamide or, better still, penicillin should be given. For a child aged 2 years, for example, 0.5 g. of sulphadimidine thrice daily or 50,000 units of penicillin twice daily should be given for three or four days. The sulphonamide group should not be administered unless a sufficient fluid intake can be assured.

The Eyes.—There is usually severe photophobia, and the patient should be screened from strong light. The eyes should be bathed often and ung. hydrarg. ox. flav. smeared along the margin of the lids. Regular and careful inspection of the cornea is essential in order to detect the first sign of ulceration. This complication calls for specialist attention, but if this is not available the pupil should be dilated with atropine drops and hot bathings carried out two-hourly. "Albucid" or penicillin eye ointment should be instilled also.

The Mouth and Ears.—The hygiene of the mouth is particularly important, as there is always some degree of stomatitis and streptococci and other organisms abound there. In spite of these precautions otitis media may occur. It should be vigorously treated with penicillin systemically, and, if suppurative, by local drops into the ear. Failure of the discharge to clear up in a reasonable time is an

indication for operation, to remove the risk of a subsequent flare-up and to prevent the hearing being permanently affected. Deafness occurring in a young child will call for special education.

Laryngitis.—The early laryngitis of the prodromal stage usually needs no treatment, and signs of obstruction are slight or absent. The late laryngitis needs vigorous treatment. As soon as signs show themselves the patient should be given 20,000 units of diphtheria antitoxin intramuscularly. A course of penicillin should also be started—100,000 units six-hourly for the first twenty-four hours, and then twice daily. If there is much obstruction a steam tent may be helpful, and preparations for operative intervention should be put in hand. Tracheotomy is the operation of choice; the pneumonia which once followed so often can sometimes be avoided by vigorous chemotherapy. Intubation may induce ulceration of the larynx (if the condition is streptococcal ulceration may already be present) and lead to subsequent stenosis.

Bronchopneumonia.—Cases of bronchopneumonia should be nursed in the fresh air as much as possible, but guarded against chills. If more than slight cyanosis is present an oxygen tent is indicated. Full doses of sulphadimidine and penicillin should be given: for a child of two years, 0.5 g. of sulphadimidine and 50,000 units of penicillin six-hourly are advisable. The fluid intake must be adequate. If restlessness occurs small doses of potassium bromide or brandy may help. The latter is also of value if there is cardiac distress. Other useful adjuncts are tepid sponging, if the temperature is high, and kaolin poultices to the chest. If an empyema develops, frequent aspiration of the fluid, with penicillin replacement, may suffice without operative intervention being necessary.

Enteritis.—Infants who show signs of enteritis should immediately be given half-strength glucose-saline by mouth two-hourly; after this, third-strength "Benger's food" may be a useful stepping-stone back to normal feeds. If vomiting is a feature and dehydration develops, an intravenous drip transfusion of 5% glucose-saline, alternating with half-strength plasma, may be necessary.

Encephalitis.—There is no certain treatment for post-infective encephalitis. Convalescent serum from recovered cases of encephalitis has been tried, and thought by some to be of use. A lumbar puncture will be necessary for diagnostic purposes.

Convalescence.—A bad attack of measles will debilitate a child seriously, and a plentiful diet with adequate vitamins is necessary in the convalescent period. The child should be kept under observation, and if the lungs have been affected an x-ray examination should be made at the end of six months.

Terminal Disinfection.—Measles virus is short-lived and no special disinfection is necessary. Infected clothes and bedding should be washed, and the patient's room given a thorough cleaning.

Prevention

Isolation.—The prodromal stage of measles is so infectious that any child suspected of the disease should, if possible, be isolated from other children at once.

Passive Immunization.—If exposure to infection is known to have taken place a decision will have to be made whether protection by immune human serum should be undertaken. Such serum is very difficult to obtain at present, and for this reason the ordinary healthy child of 3 years or over will usually have to be allowed to take his chance. With infants and young children, especially those

in a poor state of health or suffering from some other disease, every effort should be made to obtain enough serum for their protection. Whatever type is used it must be given in the first five days after exposure, the earlier the better: it is passive immunity which is being conferred, its duration is only about three weeks, and another exposure after this period may reinfect the child.

The following sera can be used:

	Protective dose up to 3 years	Protective dose after 3 years
Gamma globulin ..	2.5 ml. intramuscularly	Not determined
Convalescent serum	5 ml. "	(Age $\times$ 2) ml.
Placental globulin ..	7.5 ml. "	(Age $\times$ 3) ml.
Adult serum ..	10 ml. "	(Age $\times$ 4) ml.

These sera, although scarce, can sometimes be obtained by application to a hospital or to one of the large commercial chemists.

Of the four types mentioned gamma globulin is probably the best. It is at least twice as strong as convalescent serum, and does not carry the risk of the other human sera—homologous serum jaundice. It must not be given intravenously.

Convalescent serum is obtained by bleeding a measles patient about the fourteenth day of disease, when the antibodies are at their highest level. It is at least half as strong again as placental globulin and twice as strong as serum from adults who have had measles in infancy (adult serum). No serum reaction may be expected from any of these, as they contain human, not horse, protein; but placental globulin may cause a rise in temperature.

If sera are not available whole blood from an adult with a known history of measles can be given intramuscularly, the dose required being double that of adult serum. The risk of giving Rh-positive blood to an Rh-negative female child must be guarded against. In any case the method is not satisfactory as it produces an artificial haematoma, with its attendant dangers.

Attenuation.—If serum is available, attenuation rather than protection is a more satisfactory plan for children over 3 years. Half the full protective dose of serum should be given in the first five days after exposure. This is more economical than a full protective dose between the sixth and eighth day, though this will probably produce the same effect. The mild attack which results—if it is not too mild—will probably confer active immunity for life. With such a modified attack, however, though the child is not very ill, he is infectious and a risk to other children.

Notification.—Measles and whooping-cough were made notifiable in England and Wales in 1939. The information thus made available is of value to the public health authorities in the control of epidemics and in gathering epidemiological data.

In a booklet entitled *The Marriage Relationship* the Society of Friends emphasizes the importance of a religious background to life if marriage is to be successful. Surveying the causes of the breakdown of marriages, the report draws attention to inadequate preparation for marriage and lack of education in parenthood, housing, war and post-war conditions, and weakened belief in the permanent nature of marriage. Marriage guidance councils and other bodies are doing much helpful work, and "it should be increasingly realized that many of the difficulties encountered by young married couples could have been prevented by a talk with a suitably qualified doctor or some other experienced person before marriage." The booklet is obtainable for 9d. (postage 2d.) from the Friends Book Centre, Euston Road, London, N.W.1.