A CASE
OF
RHEUMATIC PERICARDITIS AND EXTREME
DILATATION OF THE HEART

WITH AN
INVESTIGATION INTO THE MICROSCOPY
OF RHEUMATIC HEART DISEASE

BY
F. J. POYNTON, M.D.

COMMUNICATED BY DR. NORMAN MOORE

Received February 14th—Read May 9th, 1899

THIS case, for the use of which I am indebted to the kindness of Dr. Sidney Phillips, affords both clinical and pathological evidence in support of the statements made by Dr. Lees and myself in two papers read before this Society in June, 1898. The first, by Dr. Lees, emphasised the importance of acute dilatation of the heart in rheumatic fever; the second, a joint paper, dwelt upon the frequency of its occurrence in the rheumatism and chorea of childhood.

These views we supported by clinical observations

extending over a considerable period of time, and some of the results obtained were illustrated by cardiac tracings thrown on the screen by a lantern. Though personally assured of the truth of our clinical investigation, we felt it necessary to adduce pathological evidence in its support. This was to some extent supplied in an appendix to the papers, by an analysis of 150 cases of fatal rheumatic fever in children under twelve years of age.

In the discussion that followed, the great size of some of our cardiac tracings was criticised. Dr. Ewart, on the other hand, accepted in part the frequency of a considerable increase of the præcordial dulness, but considered the explanation to be an evanescent pericardial effusion.

This case is now brought forward because it deals to some extent with these difficulties, and demonstrates clearly the extreme dilatation that may occur in rheumatic pericarditis with no fluid in the pericardium and almost no valvulitis. The results of the microscopic investigations of the heart, which I have made in this case, seem to me to clearly indicate a morbid condition of its muscular structure, which goes far to explain the great dilatation.

The case is an example of aggravated cardiac rheumatism, minor degrees of which are of common occurrence. There will be, therefore, no need for lengthy clinical details, and those facts which are concerned with the condition of the heart will alone be described.

Arthur G—, aged 15, was admitted to St. Mary's Hospital on August 25th, 1898, for shortness of breath. In December, 1897, he passed through an attack of rheumatic fever in which several joints were swollen and very painful, but for which he did not keep his bed. In February, 1898, he had another attack of rheumatic fever, which lasted for two months. Since this last attack he had suffered from occasional pains over the heart on exertion. Two days previous to his admission to the hospital a small nodule appeared over the right patella.

On admission, on August 25th, his temperature was 100° F., pulse 120, and respirations 30 ; he was very pale and emaciated. The cardiac area was increased, the limits being *one* finger-breadth to the right of the right sternal margin, and *one and a half* to the left of the vertical nipple line ; upward, the limit was the second left intercostal space. The action of the heart was excited, but no pericardial friction was detected. At the apex there was a loud, harsh, systolic murmur conducted to the axilla, and at the base an accentuated pulmonary second sound. The next day pericardial friction was heard, and this on the following days became loud and general. Upon August 30th the area of cardiac dulness had increased, extending now *two* finger-breadths instead of *one*, to the right of the right sternal margin, and still *one and a half* beyond the left vertical nipple line. The loud friction quite obscured the murmur. Upon September 2nd the friction was not so loud, but the cardiac dulness reached the *right nipple line* and extended *two* finger-breadths outside the left nipple line. Upon September 3rd the friction was loud and general, and the cardiac dulness extended *one finger-breadth beyond* the right vertical nipple line, and about *three* beyond the left. There was dulness in the fifth right intercostal space (Dr. Rotch's sign). I was unable, even with the greatest care, to detect any curving inward of the right margin of the cardiac dulness at its lower limit. Over the front of the chest there was doubtful pleural friction on both sides. Upon September 7th the dulness extended *one inch* beyond the right vertical nipple line, and there was systolic retraction of the right intercostal spaces. Pericardial friction was still well marked. The liver extended two inches below the costal margin. Cough, vomiting, restlessness, and all the signs of acute cardiac failure were apparent, and the patient died the following day. The temperature since the day of admission had never risen above 99.5°.

The *post-mortem* was made twenty-four hours after

death. During life the areas of cardiac dulness had been confirmed by Dr. A. G. Butler, Dr. Phillips's house physician. After death I verified the dulness as external to the right nipple, and far outside the left. This was also confirmed by a post-mortem assistant. Though the loud, general, and continuous pericardial friction negatived the idea of a large effusion, there seemed every probability that there would be a distinct increase in the amount of fluid, and I removed the chest wall with especial care, in order to obtain some of this fluid in Pasteur pipettes for bacteriological purposes. In this I was disappointed, for the pericardium was generally adherent, and the enlargement almost entirely due to the dilated heart, and, to a very minor extent, to the thickening of the pericardium. The area of dulness to the right of the sternum was due to the greatly dilated right auricle. In front, over a limited area, the pericardial adhesions were firm, but at the sides and behind the adhesions were quite recent, and the pericardium in places swollen to the thickness of half an inch. The recent adhesions were in the form of flakes of lymph, in the interstices of which was a little fluid. The two lungs were pushed aside by the immense heart, and were adherent to the pericardium by recent adhesions. The valvular changes were extremely slight, consisting only in some slight thickening of the mitral segments, not causing the least stenosis, and there was not any recent valvulitis. The valve rings were dilated, and the cavities of the heart crammed with pale and dark clot, and in the right auricular appendix the clot was found firmly adherent. Both ventricles seemed slightly hypertrophied. The weight of the heart with the pericardium, but empty of blood, was twenty-six ounces. The muscle was streaked with pale striæ, and had on section a pale purple colour. The other viscera showed changes compatible with a recent and acute cardiac failure.

I made numerous sections of the pericardium, heart-walls, blood-clot, and nodules, and will briefly describe

the salient points in the results obtained. (Some of the sections are shown under the microscopes.) In addition I have taken sections from other examples of rheumatic heart disease, from a case of advanced alcoholism, Addison's disease, and from a case of cardiac fibrosis. These appear to me to support the views that are held in this paper, and they will be alluded to as the points they illustrate are considered.

In the case above described, a section through the inflamed pericardium showed extreme vascular dilatation and plastic inflammatory exudation into the pericardial cavity. In the wall of the left ventricle there was a similar but slighter capillary distension, and both under the pericardium and far away from it there was a free exudation of small cells between the muscle-fibres. The muscle itself showed great loss in striation, and many of the fibres showed granules in the proximity of the nucleus, not of a fatty nature. In sections through the right ventricle, the most striking changes were beneath the pericardium, but they were visible throughout the entire thickness of the ventricle wall. Another series of sections were fixed in Hermann's osmic fluid. Both beneath the pericardium and throughout the heart-wall there was fatty degeneration of the muscle-fibres. Sections through the right auricular appendix and *ante-mortem* clot showed externally pericarditis, then hyaline change in the muscle, and coagulation necrosis in the blood-clot.

Thus it will be seen that the changes in the cardiac wall were very general, affecting muscle, vessels, and interstitial tissues. The early commencement of the interstitial changes are, however, better seen in some sections taken from a case of pericarditis of earlier date. In this case, published in the 'Lancet,' July 23rd, 1898, death was caused by a rare complication, extensive thrombosis of the large veins of the neck and upper extremities. The pericardial friction in this case was detected only the day before death. The sections show numerous foci of inflammatory exudation scattered through

the walls of the ventricle, and not localised to that part of the myocardium lying immediately beneath the pericardium. In this case also during life there was great dilatation.

Though these morbid changes are distinct and easily recognisable, the parenchymatous changes are by no means as profound in these rheumatic hearts as in some other conditions. Thus, in some sections taken from the left ventricle of a case of advanced alcoholism, it is difficult to recognise at first sight the muscular tissue, the changes are so extreme. Yet during life I especially noted that there was no appreciable dilatation, a point to which allusion will again be made. Again, though the alterations in the interstitial tissues of the heart are quite definite in these acute cases of rheumatism, they are more extreme in the sections of a fibroid heart taken from a case that terminated in the sudden death of the patient. Yet though the microscopic changes in these fatal cases of acute rheumatism were not extreme, they were general and diffuse.

The sections next described are also from a case of cardiac rheumatism, but with a different clinical history. They were taken from the left ventricle of a man aged twenty-nine. He had suffered in childhood from rheumatic fever, and five years before his death had been known to have organic, mitral, and aortic disease. He finally died in another attack of rheumatism, complicated with pneumonia. These sections show fibrosis of the ventricle—evidence, I believe, of the previous rheumatic attacks, and in addition multiple foci of small-celled exudation scattered through the ventricle—evidence of the recent illness. It is noteworthy that macroscopically this muscle appeared natural, and that there was not any recent valvulitis or pericarditis, but an adherent pericardium of old standing.

It remains now to consider shortly the bearing of these details upon the question of rheumatic dilatation. The case itself proves conclusively that great dilatation,

greater than any that was represented in our tracings last June, may occur in rheumatic fever with plastic pericarditis, but without valvulitis of any appreciable severity. It also proves that Dr. Rotch's sign may be obtained when there is no pericardial effusion. Such a case as this, taken together with the clinical evidence we brought forward in June, makes it probable that dilatation of less degree may occur without pericarditis, and that the myocardial changes are not a *sequela* of the pericarditis. If pericarditis is present the changes are concurrent, and they probably occur in the myocardium independently of any pericarditis.

It is, I believe, usually considered that the pericarditis is the starting-point of the myocardial inflammation, and is the explanation of the dilatation and severe symptoms that appear in the worst cases of cardiac rheumatism. Several observers have, however, held that the myo- and pericardial changes are concurrent, and part of the same rheumatic process. Thus, Dr. Cheadle, when writing of chronic pericarditis, suggests this in the Harveian Lectures on the "Rheumatic State in Childhood" in 1888, and the same opinion was expressed by the late Dr. Sturges in the term rheumatic carditis. Dr. Theodore Fisher, both at the meeting of this Society in June and again at Edinburgh in July, stated that he had observed pathological changes in the myocardium apart from any pericarditis.

The microscopic sections, I think, support the view that when pericarditis is present, the myocardium is also affected concurrently, for the changes in the heart-walls commence by numerous scattered foci, some of them far from the pericardium, and the changes in the muscle are general.

This view is also supported by the recent advances of pathology, especially in the investigation of cardiac muscle, for these impress the fact that micro-organisms gaining access to the body give rise to toxins which circulate in the system. These, doubtless, by delicate

chemical reactions injure the tissues, and usually have, to some extent, a specific action on certain classes of tissues. Thus the rheumatic poison appears to especially attack tissues which are in frequent movement or exposed to friction, for example, the heart, the joints, the tendons, the skin, and voluntary muscles. It is possible that these tissues have a certain chemical relationship which renders them thus liable. However this may be, when after death from rheumatic morbus cordis, acute changes are found throughout the heart, it is most probable that if rheumatism be caused by a toxine, these changes arose concurrently in a general infection of its structures.

Allowance, I think, must also be made for the influence of the surrounding tissue pressures. The pericardium, with a cavity between its two layers, and with numerous vessels and lymphatics lying in the deeper part of its visceral portion, will, in reacting to the rheumatic poison, permit free exudation into its sac, and after death the result is striking and at once appreciated. But in the ventricle wall the firm and contracting muscle will probably allow but little exudation, yet the morbid changes may in the gravity of their results be even greater. Post mortem we may find little macroscopically, and when we do find changes we are naturally inclined to consider that they are less frequent in their occurrence, and commence after the onset of the pericarditis. We have many proofs that the morbid results produced by a poison may be extreme, yet the obvious morbid changes in the tissues themselves may be slight, as, for example, in acute osteomyelitis or in tetanus.

Another point is one upon which Dr. Lees has frequently laid stress, and to which he recently called attention when introducing the discussion upon rheumatic heart disease in childhood at the Edinburgh meeting in July, 1898. It is that in rheumatism, dilatation is frequently very marked, yet the clinical symptoms, provided the patient be at rest, are often remarkably slight.

At first sight this is difficult to realise, but I believe

the explanation to lie in the fact that, the function of the cardiac muscle is damaged in a peculiar way rather than destroyed by the rheumatic poison.

A tissue so complex as the cardiac muscle is liable to a variety of pathological changes, and these probably have a different morbid significance. Some of these changes may be far more detrimental to the vital properties of the muscle than others. Those that are the most destructive may cause death from syncope without appreciable dilatation, as occurred in the case of alcoholism referred to above, and also in the case of Addison's disease. Those changes which are less destructive may cause perversion or impairment of function, ending perhaps in death, but with great dilatation of the heart during life.

Clinically, the first definite evidence of advanced fatty degeneration of the heart may be sudden death; and so we may expect that in rheumatism, with morbid changes of a less severe type, the symptoms due to the dilatation may be but slightly marked.

It is, I think, also legitimate to consider what would have happened had the case recorded in this paper made a temporary recovery. The pericardium, we know, would have become less hyperæmic, have shrunk, and finally have become generally adherent. It is more difficult to follow the fate of the muscle, but probably much would have been replaced by fibrous tissue. The sections through the ventricle of the case of recurrent rheumatism support this. If the boy had then died in a relapse, twelve months later, there would have been a temptation to lay much stress upon the mechanical effect of the obvious adherent pericardium, and little upon the loss of vital power due to the damaged muscle.

Finally, in rheumatic children it is certain that it is the active rheumatism that usually kills them. It is very probable that in adults also active rheumatism is of more importance than might be thought; and that the explanation of some cases of organic heart disease, which

fail to react to treatment, and in which the symptoms far outweigh the clinical signs, is not the adherent pericardium so often found after death, but this active rheumatism injuring the cardiac muscle. The evidences of this rheumatism may be slight, and possibly we do not even yet recognise fully its manifestations. I have been struck with the fact that not infrequently there is in these unfavourable cases a history of recent rheumatism, and now and again such definite symptoms as arthritis develop during their stay in hospital. These symptoms could then hardly be due to a fresh infection, and are more probably an evidence of active rheumatism which has existed from the time that the cardiac breakdown brought them to the hospital.

Since this paper was communicated in February, I have obtained microscopic sections of the cardiac muscle in three cases, each of which show changes that bear upon the questions considered in this paper. I am indebted to Dr. Cheadle, Dr. Lees, and Dr. Phillips for leave to give a brief outline of their histories.

Dr. Cheadle's case was that of a man of thirty, who had suffered from rheumatic fever in childhood, and was admitted for aortic and mitral incompetence. The symptoms were most urgent, quite overshadowing the physical signs. The sections of the left ventricle showed much fibrosis.

Dr. Lees' case, that of a girl of eighteen, was one of acute carditis, almost precisely similar to the case recorded, but even more convincing because there was no marked thickening of the pericardium. The fatty changes throughout the heart-wall were quite remarkable.

Dr. Phillips' case was that of a woman of twenty-three, who died of "infective endocarditis," probably of rheumatic origin. There were mural vegetations upon the endocardium of the left ventricle. The ventricle-wall showed *acute* inflammatory changes, but the pericardium was completely adherent from a previous rheumatic attack.

I am indebted to Dr. Sanders for all the sketches that have been made of the sections that were shown.

(For report of the discussion on this paper, see 'Proceedings of the Royal Medical and Chirurgical Society,' Third Series, vol. xi, p. 126.)

DESCRIPTION OF PLATE IX,

Illustrating a Case of Rheumatic Pericarditis and Extreme Dilatation of the Heart (F. J. POYNTON).

FIG. 1*a*. Diagram of pericardium and wall of the left ventricle (rheumatic pericarditis).

- a*. Points to the pericardium.
- b*. Points to the position in the ventricle from which the sketch shown in Fig. 1*b* was made.

FIG. 1*b*. Section of a portion of the left ventricle (rheumatic pericarditis).

- c*. Points to distended capillaries.
- d*. Points to cellular exudation between the muscle-fibres.

FIG. 2*a*. Diagram of pericardium and wall of the left ventricle (rheumatic pericarditis).

- a*. Points to the inflamed pericardium.
- b*. Points to the position in the ventricle from which the sketch shown in Fig. 2*b* was taken.

FIG. 2*b*. Section of a part of the left ventricle (rheumatic pericarditis).

- c*. Points to a capillary cut longitudinally.
- d*. Points to a free cellular exudation between the muscular fibres.

DESCRIPTION OF PLATE X,

Illustrating a Case of Rheumatic Pericarditis and Extreme Dilatation of the Heart (F. J. POYNTON).

FIG. 3. A section of a part of the left ventricle just beneath the pericardium (rheumatic pericarditis).

- a*. Points to the dark dots which represent fatty changes in the muscle-fibres. Hermann preparation.

FIG. 4. A part of the same ventricle wall as that represented in Fig. 3, but close to the endocardium.

- a*. Points to the fatty changes in the muscle-fibres. Hermann preparation.

FIG. 5*a*. Section through pericardium and wall of the left ventricle in a case of early rheumatic pericarditis, fatal by severe venous thrombosis.

- a*. Points to the pericardium.
- b, c, d*. Point to foci of cellular exudation spreading from the blood-vessels.

FIG. 5*b*. One of the foci seen in Fig. 5*a*, more highly magnified.

- a*. Points to cellular exudation between the muscular fibres.

Fig. 3.

Fig. 4.

Fig. 5a.

Fig. 5b.

DESCRIPTION OF PLATE XI,

Illustrating a Case of Rheumatic Pericarditis and Extreme Dilatation of the Heart (F. J. POYNTON).

FIG. 6. Section of a part of the left ventricle from a case of recurrent rheumatism.

- a.* Points to cellular exudation between the muscular fibres.
- b.* Points to a fibrous strand, spreading from the vessels and implicating adjacent muscular fibres.

FIG. 7. Section of a part of the same left ventricle as Fig. 6, under higher magnification.

- a.* Points to cellular exudation.
- b.* Points to hyaline change in muscle-fibres.

FIG. 8. Section of a part of the left ventricle from a case of chronic alcoholism.

- a.* Points to a muscle-fibre still recognisable.
- b.* Points to a nucleus of a muscle-fibre.
- c.* Points to fatty changes in a muscle-fibre.
- d.* Points to spaces left by the complete destruction of the muscle-fibres. Hermann preparation.

Fig. 6.

Fig. 7.

Fig. 8.

DESCRIPTION OF PLATE XII,

Illustrating a Case of Rheumatic Pericarditis and Extreme Dilatation of the Heart (F. J. POYNTON).

FIG. 9. Section from the left ventricle of a man aged 30, who had suffered from rheumatic fever in childhood.

- a.* Points to the muscle-fibres.
- b.* Points to coarse strands of fibrous tissue traversing the ventricle.

FIG. 10. Section of a papillary muscle from the same case as Fig. 9.

- a.* Points to fibrous strands between the muscle-fibres.
- b.* Points to the muscle-fibres.
- c.* Points to fibrosis around a blood-vessel.