THE AFTER-RESULTS IN FORTY CONSECUTIVE CASES
OF
VAGINAL HYSTERECTOMY PERFORMED
FOR CANCER OF THE UTERUS

BY
ARTHUR H. N. LEWERS, M.D.LOND.,
OBSTETRIC PHYSICIAN TO THE LONDON HOSPITAL

Received October 26th—Read November 13th, 1900

THE series of forty cases recorded in this paper includes all the cases in which I have performed vaginal hysterectomy for cancer of the uterus up to the end of April, 1899, when I completed my fortieth case. Up to the year 1893 I had for the most part treated cases of cancer of the cervix that appeared suitable for operation by the supra-vaginal amputation of the cervix, and reserved total extirpation of the uterus for cases of primary cancer of the body of the uterus.

Since that time, however, I have been more and more convinced that, to perform the supra-vaginal amputation of the cervix in such a way as to give the patient the utmost chance that the disease will not recur, is, in the majority of cases, a more difficult matter than to perform

the operation of vaginal hysterectomy. I have for some time past only performed the supra-vaginal amputation of the cervix for very early cases of cancer of the vaginal portion, in which there is a limited and superficial growth. Up to April, 1899, I have had thirty-three cases of the supra-vaginal amputation of the cervix for cancer, but I do not propose to deal with these cases in this paper.

The Cases that remain Free from Recurrence.

Among the forty cases of vaginal hysterectomy there are fourteen cases in which the disease has not recurred. In two of these, Nos. 3 and 4, in the series, although the clinical evidence convinced me that they were really examples of cancer, this cannot be proved by examination of the specimens. In the remaining twelve cases there can be, I think, no doubt whatever on this point. The specimens have been carefully examined and are here for inspection to-night, as also are sections of the growth in each case. This appears to me an important point.

My object in the present paper has been to inquire more especially into the *after-results* in the cases of cancer of the uterus in which I have performed vaginal hysterectomy. In such an inquiry it is essential, first of all, to prove that, in those cases in which the disease has not recurred, the growth was really malignant. In order to satisfy this condition I have in each case had an independent report on the growth by Mr. Targett of the Clinical Research Association, and his report will be found *verbatim* in the account of each of the cases. The necessity of a careful microscopical examination of the growth in each case is almost self-evident, and when the microscopical examination confirms the opinion based on the clinical features and physical examination, there can, I think, be no hesitation in accepting the result as fully proved.

In three cases in this series, Nos. 3, 4, and 25, the microscopic examination did not confirm the diagnosis of malignancy based on the physical examination. In

Nos. 3 and 4, though I have excluded them as "non-proven," I feel little doubt that they were cases of cancer. I have myself met with an example of cancer of the cervix in which the appearance of the growth under the microscope would certainly not have warranted a diagnosis of cancer, but in the case in question, one treated by the supra-vaginal amputation of the cervix, the malignant nature of the growth was proved by recurrence. As regards *Case No. 25* in this series, the clinical features and the physical examination led me to believe without any hesitation that the case was one of malignant ulcer of the cervix, but on examination of the cervix by the microscope Mr. Targett reported that the case was one of tubercular ulceration. To make quite certain he examined two portions, each taken from a different part of the cervix. I think this case is an important one, as clinically it exactly resembled one of cancer of the cervix. Tubercular ulceration of the cervix as a primary condition must be extremely rare. Very little is to be found about the subject in works on gynaecology, and it has not been regarded as a condition ever likely to give rise to difficulty when the diagnosis of cancer is being considered. Yet in my case (No. 25) primary tubercular ulceration of the cervix produced a condition identical, as far as physical examination was concerned, with that resulting from malignant ulceration. I have seen a very large number of cases of cancerous ulcer of the cervix, certainly some hundreds, and I am sure that in the case in question the clinical features of the cervix were precisely similar to those produced by the ulceration of a cancerous growth. The occurrence of such a case emphasises the importance of always examining growths supposed to be cancer under the microscope before accepting them as really malignant. Again, one has occasionally heard of a case pronounced by a competent authority to be one of cancer of the cervix, where after a long time the patient gradually improved, and ultimately made a good recovery to the surprise of every one con-

nected with the case, and no doubt to the discredit of the gynæcologist who had made the diagnosis. It seems to me, in the light of the case above referred to, No. 25, that not improbably some such cases may have really been suffering from tubercular ulceration of the cervix, which presumably may occasionally get well spontaneously. However that may be, in Case No. 25, believing it to be a typical example of cancer of the cervix, I performed vaginal hysterectomy, and the patient remains quite well. It is more than four years since the operation, but I have, of course, excluded the case from Table II, the list of cases proved to be cancer of the uterus, and treated by vaginal hysterectomy, which remain free from recurrence at the present time.

To return to the question of the after-history of patients suffering from cancer of the uterus and treated by vaginal hysterectomy, in addition to proving the malignancy of the growth in each case, there is the necessity of ascertaining the state of the patient after the operation. It is not an easy thing to do. To have any real value inquiries have to be made from time to time over long periods; for instance, two cases in this paper have been kept under observation for more than seven years. The evidence of non-recurrence may be obtained either by actual vaginal examination, or by seeing the patient at intervals and observing that she continues in good health, or by hearing at intervals from her medical attendant that she continues well, or in some cases by hearing from the patient herself at intervals that she continues well. No doubt the most satisfactory thing is to examine the patient; in eight out of the twelve cases in Table II I have myself examined the patients from time to time. In some cases, however, when the patient continues well, she may not be unwilling to come to see you, but she will object to be examined, as she has nothing the matter. Two cases (Nos. 7 and 24, Table I) are in this category. In some cases one can only hear by letter from the patient, or her doctor, from time to

time. This evidence is not so unsatisfactory as it might appear, and for this reason: in those cases in which recurrence takes place the downward progress to a fatal issue is generally fairly rapid—a matter of months; so that if inquiries made at long intervals—once a year, for instance—result in a favourable report, there can be no reasonable doubt that the disease has not recurred.

Numerous cases of vaginal hysterectomy, and series of cases of vaginal hysterectomy for cancer, have been published from time to time; but I am not aware of any considerable series of such cases in which the proof both of the malignancy of the growth, and especially of the condition of the patients many years after the operation, has been given so rigorously, and in such detail, as in the present series of cases. As regards the after-histories especially, it will be seen that I have given the date on which I either examined the patient, or saw her without examining her, or heard from herself by letter, or heard from her usual medical attendant by letter.

As regards the Intervals that have elapsed since the Operation without Recurrence in the Twelve Cases referred to.

In one case (No. 7) more than seven years have passed, and the patient is known to be well.

In one case (No. 8) seven years have passed without recurrence.

Both 7 and 8 were examples of primary carcinoma of the body of the uterus.

In one case (No. 15) six years passed without recurrence.

In one case (No. 19) five and a half years have passed without recurrence.

In one case (No. 23) five years have passed without recurrence.

In one case (No. 24) nearly five years have passed without recurrence.

In one case (No. 26) nearly four years have passed without recurrence.

In one case (No. 29) more than three years have passed without recurrence.

In one case (No. 32) two years and ten months have passed without recurrence.

In one case (No. 33) two and three-quarter years have passed without recurrence.

In one case (No. 36) more than two years have passed without recurrence.

In one case (No. 39) more than one year has passed without recurrence.

Omitting No. 39, it will be seen, therefore, that there are eleven cases out of the forty remaining free from recurrence from two to seven years after the operation—that is 27·5 per cent.

This result appears deserving of close attention, since at the present time some gynæcologists have been so much discouraged by their experiences as to doubt if it is worth while to operate on cases of cancer of the uterus at all.

The comparatively high percentage of patients remaining free from recurrence in my series is, I believe, largely due to the careful selection of cases. My practice has been generally to examine the cases carefully under anæsthesia, and only to operate when the disease has appeared, so far as physical examination can decide, to be limited to the uterus.

As regards the Cases in which Recurrence is known to have taken place.

The disease is known to have recurred in 18 cases.

In 12 of them recurrence took place during the 1st year.

In 2 " " " " 2nd "

In 2 " " " " 3rd "

In 2 " " " " 4th "

The 40 cases may be divided into groups as follows:

In 12 cases there has been no recurrence.

In 18 cases there has been recurrence.

4 cases died.

In 2 cases the nature of the disease cannot be proved by examination of the specimens.

In 3 cases I have no definite information as to whether the disease recurred or not.

In 1 case the disease proved to be primary tubercular ulceration of the cervix, and not malignant disease.

40

It will be seen that there were no recurrences later than during the fourth year after operation; so that when a patient has passed through four complete years without recurrence, there seems to be some probability that she may remain permanently well.

Mortality of the Operation.

There were three deaths from the operation in the series of forty cases. This represents a mortality of $7\frac{1}{2}$ per cent. There was also one case which died of pneumonia three weeks after the operation. My colleague, Dr. Schorstein, who made the post-mortem examination, did not, however, consider the pneumonia to be of septic origin, and the peritoneal wound at the floor of the pelvis was healed and normal, so that I do not think this death is properly to be attributed to the operation. Still I think it right to mention it. As regards the cause of death in the three fatal cases, in Case 2 death was not due to any of the special dangers of vaginal hysterectomy, such as hæmorrhage, peritonitis, etc. It seemed that the patient could not stand a severe operation. She had a fatty heart, and died of œdema of the lungs forty-eight hours after the operation. In one case, No. 13, death was due to hæmorrhage; and in one case, No. 18, peritonitis was the cause of death. In that case forceps had

been used, as in the large majority of my cases, for securing the broad ligaments, and they were removed at the end of forty-eight hours. This interval is certainly sufficient as regards security from hæmorrhage, but it may not be always sufficient to prevent the breaking down of the recent adhesions that shut off the general peritoneal cavity from the region of the wound when the forceps are being removed. Case No. 18 was doing perfectly well till the forceps were taken off. While this was being done she suddenly complained of sharp pain in the abdomen, and from that time began to do badly, symptoms of peritonitis appeared, and she died on the seventh day. I believe the cause of the peritonitis was that the adhesions shutting off the general peritoneum were broken down in taking off the forceps, and the region of the wound, which is always in a foul, sloughy condition at this stage of the case when pressure forceps have been used, was thus put into communication with the peritoneum. Since that case I have generally left the forceps on for either three or four complete days in order to allow time for the protecting adhesions to become firm.

SOME POINTS IN THE TECHNIQUE OF THE OPERATION.

The mode of securing the broad ligaments.—Various instruments, special forms of pressure forceps, and special kinds of clamp have been invented and used for securing the broad ligaments. I have never used any special instrument for this purpose, but have always used Wells's pressure forceps, large or small, according to circumstances. In a few of my earlier cases (Nos. 1, 2, and 3) I used no pressure forceps, but relied entirely on ligatures. Since then I have always used Wells's forceps; occasionally a ligature or two have been used as well, but not often.

The posterior vaginal wall.—The cut edge of the posterior vaginal wall often bleeds rather freely. As a rule, after opening Douglas's pouch I pass sutures with a

Hagedorn's needle from the vaginal wall near the cut edge to the peritoneum and back again, so as to be able to tie the cut edge in some two or three sections. This at the same time avoids a wide separation of the peritoneum from the posterior vaginal wall. There is much loose tissue between these structures, and they separate easily to a considerable extent. This is avoided by suturing in the manner described.

Use of the cautery.—Although I now prefer vaginal hysterectomy to supra-vaginal amputation of the cervix, except in early cases of cancer of the vaginal portion, partly because it is an easier operation and partly to avoid the chance of subsequent dysmenorrhœa, which occasionally follows supra-vaginal amputation owing to contraction of the cicatrix, yet it is well to bear in mind that it is more especially in the direction of the vagina and of the connective tissue round the cervix that cancer of the cervix tends to spread. Only rarely does it spread to the body of the uterus, at least till such a late period in the case that complete removal of the diseased tissues has become obviously impossible.

I believe it is a good plan when operating for cancer of the cervix, after the cervix has been freed, to apply Paquelin's cautery to the cut surfaces adjacent to the growth. I have done this in all except a few of my earlier cases.

SOME COMPLICATIONS OF VAGINAL HYSTERECTOMY.

Secondary hæmorrhage.—In only one case (No. 39) have I found bleeding occur *immediately after* removing the forceps. The bleeding was not much, but I thought it better to expose the upper part of the vagina with a Sims's speculum, and secure the bleeding point with another pair of Wells's forceps, which was left for two days longer. No subsequent trouble occurred. It is clearly an unusual accident.

Secondary hæmorrhage at a later period occurred in Case 28 ; this was actually at the end of the third week, and required large pressure forceps to control it. In all cases where forceps have been used the tissues at the upper part of the vagina by the time the forceps are taken off, and until some two or three weeks after the operation, are in a sloughy condition. If secondary hæmorrhage should occur at any time during this period, great care is necessary when applying forceps to stop it to take up nothing more than is absolutely required to secure the bleeding point. The vaginal portion of the cervix—the usual familiar landmark—is of course absent, and a portion of the bladder, or one of the ureters, may be easily picked up. Fortunately secondary hæmorrhage, either when the forceps are taken off or subsequently, is very uncommon. I have only met with it in the two instances mentioned in the series of forty cases.

Injuries to ureters.—The only case in which a ureter was injured was No. 28, the case in which secondary hæmorrhage occurred at the end of the third week. As mentioned above, pressure forceps were applied and left on for two days. Not long after removing them the patient began to suffer from incontinence of urine. When the raw surface of the top of the vagina had healed, this incontinence still continued. I thought at first it might be due to a vesico-vaginal fistula. When, however, the bladder was filled with milk, while the vagina was fully exposed with a Sims's speculum, none of the milk came through into the vagina, but the flow of urine still continued. It was thus evident that the incontinence was due to a ureteral fistula. The patient was unwilling to have anything done at that time, and I saw nothing of her for several months. She then one day came up to the ward to tell me that she was quite dry, and had been so for some time. I examined her, and found no sign of incontinence. I have seen her several times since, and there has never been any sign on examination of anything wrong with either of the kidneys.

In Case 4 the patient suffered from hæmaturia for some days. The urine was bloody when drawn off with the catheter. This condition gradually disappeared, and the urine became normal. The hæmaturia may possibly have been caused by reflex congestion of the kidneys, due to a portion of a ureter, or of the bladder, having been nipped by the pressure forceps. She had no incontinence of urine, and remains quite well, nearly ten years since the operation, so that no permanent damage could have been done.

Injuries to ureters have not been, as far as I have been able to ascertain, so very rare in the history of vaginal hysterectomy, but the case mentioned (No. 28) is the only one in which the accident has happened in my own practice. Nor have I met with any cases where the ureters have been injured among my thirty-three cases of supra-vaginal amputation of the cervix—an operation in which there is practically the same chance of injuring the ureters as in vaginal hysterectomy.

Injuries to the bladder.—I have not had a case where the bladder has been accidentally injured, either among my vaginal hysterectomies or among my cases of supra-vaginal amputation of the cervix. I have, however, seen a bad case of vesico-vaginal fistula after a vaginal hysterectomy, said to have been performed in South America.

Cases suitable for vaginal hysterectomy.—When there is evidence that the disease has spread beyond the anatomical limit of the uterus in any direction, it is, I believe, hopeless to expect any permanent benefit from vaginal hysterectomy, even if it is possible to perform the operation. In many cases it is at once evident on examination that the growth has spread far beyond the uterus. No difficulty arises in coming to a conclusion in such cases. In other cases, where, on an ordinary examination, the mobility of the uterus appears to be unimpaired, it is not so easy to decide. The uterus may be freely movable, yet the base of the bladder may be

involved. Again, the mobility at an ordinary examination may seem normal, yet when an attempt is made to draw the cervix down to the vulva a check is felt at one side or the other in the region of the broad ligaments, or of the utero-sacral ligaments, due to the extension of the malignant growth in the corresponding direction. This extension can generally be best appreciated by examining *per rectum* while the cervix is being drawn down with a volsella.

Cases of this kind are certainly unsuitable for radical operation. It may be possible to remove the uterus, but that is of little use if a portion of broad ligament or utero-sacral ligament, infiltrated with the malignant growth, is left behind.

In most of these doubtful cases the best plan is to make a careful examination under anæsthesia before deciding as to operation.

For my own part, I have never felt it to be justifiable to perform an operation like vaginal hysterectomy as a palliative operation,—in other words, to perform it where one feels certain that the patient has not even a chance of obtaining any permanent benefit.

On the other hand, there may be a very extensive growth, and yet, if it does not seem to have extended beyond the uterus, operation is advisable. One must not be too ready to reject cases as hopeless. Case No. 22 in this series is a good example of this. The growth from the cervix was a very large one of the “cauliflower” type, and yet she certainly remained well for more than three years after removal of the uterus. I am not even certain that she had recurrence, as, though she is dead, she died in America a comparatively short time after I last examined her, and there was then no sign of recurrence.

Again, No. 19 had been told at another hospital that her case was too advanced for operation, yet she remains well more than five years after the operation. In No. 15 also, though the growth is obviously a large one,

the patient remained free from recurrence for six years, when she died of phthisis. No. 23 is another instance in point; the growth is a large one, but she remains well five years after operation. These cases, then, Nos. 15, 22, and 23, go to show that the size of the growth from the cervix is relatively of small importance, and that the essential point is whether there is any extension to the tissues outside the uterus itself.

I have arranged the cases in two tables.

Table I is a list of the forty vaginal hysterectomies for cancer which I had performed up to April, 1899, with some of the more important details of the cases.

Table II is a list of the twelve cases in which the disease is known not to have recurred.

Notes of the cases in Table II are also given.

Conclusions.—The conclusions that appear to follow from a consideration of the facts in this paper are—

1. That in a certain proportion of cases patients suffering from cancer of the uterus may be relieved by operation for periods of many years; in some cases for so long a time, seven years and upwards, that there seems some probability that the relief may be permanent.

2. That the proportion of cases in which this result can be expected must remain very small so long as cases generally only seek advice at a late stage of the disease; and that consequently—

3. The great desideratum is early diagnosis. Improvement in this direction depends to some extent on a better appreciation, on the part of women themselves, of the early symptoms of the disease, and especially of the significance of bleeding after the menopause, or of bleeding occurring at an earlier time of life between the menstrual periods.

Early diagnosis, of course, also depends partly on the profession. Especially important is the general recognition of the gravity of the symptom just mentioned. It is equally important also to bear in mind that patients suffering from cancer of the uterus may, and generally do,

for a relatively long period *look* quite well. They may be well nourished, or not infrequently even excessively fat ; and, as regards the aspect of the face, they may appear to be in perfect health.

Notes of the twelve cases (Table II) in which the disease has not recurred.

CASE 7. MRS. D— (with two illustrations).—The patient was 53 years of age. She had been married twenty-one years ; she had had no children, but two miscarriages, each at about three months, the last thirteen years before I saw her. I was asked to see her in consultation with Dr. Godson and Dr. Stott, of 1, Highgate Road, N.W. The symptoms were continued hæmorrhage for seven or eight months, with occasional profuse losses of blood and severe pain in the hypogastric region. On examination the uterus was found to be moderately enlarged and freely movable. The vaginal portion of the cervix was normal. The sound occasioned very free bleeding. It was agreed that the best course would be to dilate the cervix and examine the endometrium. This was accordingly done. Two specially prepared laminaria tents were inserted, and the next afternoon the dilatation of the cervix was completed with Hegar's dilators under anæsthesia. There was a hard, irregular condition of the inside of the body of the uterus. Nothing could be detached either with the finger or with a curette. The diagnosis rested on the symptoms, and on the fact that the condition of the endometrium could not be classified as belonging to any non-malignant state. I advised that vaginal hysterectomy should be performed, and I performed the operation on February 1st, 1893, assisted by Dr. Godson, Dr. Stott, and Dr. Sequeira, who was resident accoucheur at the London Hospital at that time. Mr. F. W. Braine gave the anæsthetic. The patient made a good recovery. A section from the uterus was made by the Clinical Re-

search Association, and reported on by Mr. Targett as follows :—"The growth of the uterus (Mrs. D—) is a columnar-celled carcinoma of the body, of the villous type. It has deeply invaded the muscular substance."

A drawing of the uterus, and of the sections, accompany this paper. The patient came to see me on June 8th, 1900, just to show me that she was quite well. The interval in this case without recurrence is, therefore, seven years and four months. Her present medical attendant is Dr. Andrew Brown, of 27, Lancaster Road, Belsize Park.

CASE 8 (with two illustrations).—L. S—, a married woman aged 57, was admitted under my care into the London Hospital on May 1st, 1893. She was sent up to me by Dr. Best, of Dover.

Previous history.—She was born at Blackburn, in Lancashire. During her infancy her family removed to Crayford, in Kent. Her circumstances have always been comfortable. She had scarlet fever in childhood, and cholera in 1849.

Menstruation.—The catamenia began between fourteen and fifteen. There was some irregularity for the first ten months, but after that she was very regular every four weeks, and did not lose very much. She always suffered a great deal of pain after the periods. She was married at the age of thirty-three, and had six children, but no miscarriages. The last confinement was fourteen years ago.

Present illness.—About two years ago the periods began to be irregular. She would "see nothing" for two or three months, and then a period would come on with an excessive loss, containing clots. During the last eighteen months she has been losing blood constantly, and for the same time has suffered a good deal of pain, more severe at night, in the lower part of the abdomen. She was closely questioned as to the time when the various symptoms arose, and it became clear that as long ago as

two years there was a red discharge between the periods, and that for the greater part of the last two years she had had a good deal of pain in the back and lower part of the hips. She said also that she always had suffered a great deal of pain during her periods from the time they began. As she grew older the pain at the periods was not so severe until about two years ago, when it recommenced. The pain during this two years has been constantly present, and has increased in severity.

Present state (May 1st, 1893).—She was a stout, heavy woman, and, as regards the appearance of the face, she did not look very ill. There was no cachexia. Her weight on admission was 12 st. 13½ lbs.

Nothing abnormal could be felt on examination of the abdomen.

On vaginal examination there was a considerable descent of the anterior vaginal wall, which protruded from the vulva. There was an extensive rupture of the perinæum, reaching back to the edge of the sphincter. There was also slight prolapse of the rectal mucous membrane. Through the speculum the vaginal portion of the cervix appeared healthy. The uterus was fairly movable, and in the normal position. The body of it was somewhat enlarged, and the sound passed 3¼ inches, causing rather free bleeding.

She was kept under observation in the hospital during the whole of May, and during that time there was a constant red vaginal discharge, and she complained of pain about the lower part of the abdomen, more severe at night.

On May 31st, at 5 p.m., two specially prepared laminaria tents were inserted into the uterus with strict antiseptic precautions, and on the following day (June 1st), at 2 p.m., she was anæsthetised with ether. The tents were removed, and it was found that the cervix admitted the index finger easily. The uterus was drawn down to the outlet with a volsella. The finger was passed into the uterus, and detected a hard, irregular condition of the

endometrium ; this was particularly marked at the upper part and to the right. I could not classify the condition as resembling any non-malignant pathological state, and I came to the conclusion that the case was one of primary carcinoma of the body of the uterus. I should have mentioned that nothing could be scraped away from the endometrium with a curette.

Operation (June 8th, 1893).—The patient was anaesthetised with ether, and placed in the lithotomy position. The vulva and vagina were thoroughly disinfected with perchloride lotion 1 in 1000. Having drawn down the cervix, the anterior fornix was incised first, and the bladder separated from the cervix as high as the internal os. The posterior fornix was then similarly incised, and Douglas's pouch opened. The cervix was cleared laterally as high as the level of the internal os, pressure forceps being used to control the bleeding where necessary. The utero-vesical pouch of the peritoneum was then opened. The remaining lateral attachments were clamped piece by piece, the left broad ligament being first secured and divided. The uterus was then, as it were, delivered, the left side of it first, so that at this stage it was hanging by the right broad ligament. The latter was then secured with pressure forceps and divided. The uterus was then removed. The vagina was filled with iodoform gauze, the upper part of the plug projecting for an inch or so into Douglas's pouch.

The patient's progress after the operation was very satisfactory. The highest temperature in the first week was 100.8° on the evening of June 11th. On one occasion in the second week (June 16th) it reached 101.4° . After June 20th the temperature was practically normal, only once after that date reaching 100° . The pressure forceps were removed on the afternoon of June 10th.

The patient went home on July 21st, 1893.

I have seen her every summer since the operation, and have examined her. So far she has remained free from

any sign of recurrence, and appears to be in perfect health.

Sections for examination under the microscope were prepared by the Clinical Research Association, and reported on by Mr. Targett as follows:—"This growth from the body of the uterus is a columnar-celled carcinoma, in which the tubular arrangement of the cells is well preserved. The tissues have shrunk in hardening, but the sections are sufficiently good for microscopic drawings."

A drawing of the uterus and drawings of the sections accompany this paper.

The last time the patient was examined was on July 19th, 1900, at the London Hospital. There was no sign of recurrence, and she was quite well.

CASE 15 (with one illustration).—Mrs. J—, a married woman aged 57, was seen by me in consultation with Dr. Howse, then of Barking Road, E., but now of Reading, in February, 1894.

She had had thirteen children, the last fifteen years ago, and two miscarriages some time before the last confinement. The complaint when I saw her was that she had been losing blood from the vagina for the last eight weeks. Before this appeared there had been a yellow vaginal discharge for nine months. She had had no pain. She thought she had lost flesh, and said her appetite was bad. The menopause occurred at fifty-one.

On examination there was found a large "cauliflower"-like growth springing from the vaginal portion of the cervix, forming a ring of soft projecting growth round the os uteri, and bleeding readily on examination.

It was decided to make a thorough examination under anæsthesia, with the understanding that, if the conditions were found favourable, the uterus should be removed. Accordingly, on February 27th, 1894, the patient was anæsthetised. It was then found that the uterus could be drawn easily down to the vulva, and that the growth did not seem to have spread beyond the limits of the cervix.

There was a ring of apparently healthy vaginal portion outside the malignant growth all the way round it. This can be well seen even now in the specimen.

The uterus was accordingly removed, the steps of the operation being much the same as in the preceding case, except that Paquelin's cautery was very freely applied to the cut edges of the vaginal fornices and neighbouring tissue from which the cervix had been detached. The patient made a very good recovery.

I heard of her from time to time from Dr. Howse. She remained free from recurrence. He saw her several times in the course of her last illness (phthisis), and there was no return of the cancer up to the time of her death, which occurred in February, 1900, and was due to phthisis. Dr. Howse had attended her many times in former years for lung trouble.

The interval since the operation without recurrence is therefore in this case *six years*.

Sections of the growth for the microscope were prepared by the Clinical Research Association, and reported on by Mr. Targett as follows:—"The second portion of this growth (Mrs. J—) has been examined, but the tissue is so necrotic that the sections are not worth drawing. The nature of the growth is undoubtedly squamous-celled epithelioma, but owing to decomposition the cells are much vacuolated, and the tissue has therefore an unnatural spongy appearance. It is not worth the expense of a drawing." I may add that the uterus had been more than five years in spirit before Mr. Targett had a piece of the growth for sections.

A drawing of the uterus accompanies this paper.

CASE 19 (with two illustrations).—E. S—, a married woman, 52 years old, was admitted under my care into the London Hospital on October 9th, 1894. She had had two children, the last one twenty-six years ago, but had never had any miscarriages.

For the last eight years she had been a monthly nurse.

She has always enjoyed good health, and there is no history of any previous illness. She has always had a comfortable home. She has been a teetotaler all her life, and has always lived in or near the East End of London. Her two children were born within three and a half years of her marriage.

Menstrual history.—The catamenia appeared when she was thirteen years old, and she was always regular every four weeks except when pregnant. She always suffered very much from headache at the menstrual periods. The menopause occurred five years ago, *i. e.* in 1889. After the menopause she “saw nothing” till the end of June, 1894, when, on awaking one morning, she found that she had lost a considerable quantity of blood during the night. Throughout the whole of July, 1894, she continued to have a series of losses, which occurred irregularly, the discharge being blood-stained and offensive. Occasionally it ceased altogether for two or three days. In August there was much less discharge, but since then, and up to the present time, it has continued more or less. She has been able to continue at her work.

She consulted Dr. Howse, then of Barking Road, now of Reading, in September, 1894, and afterwards was an in-patient at the Samaritan Hospital for four days. She said she was told there that the disease was too far advanced for any operation to be advisable.

A month before her admission to the London Hospital, *i. e.* October, 1894, she first noticed a pain in the left iliac and in the hypogastric regions. It was of a burning character, not relieved by lying down, and not worse at night. The pain has gradually become more severe. A week before her admission she began to feel a pricking and shooting sensation in the left breast. She said she had not been getting thinner.

State on admission.—She was fairly well nourished, but rather sallow. On October 11th (1894) she was examined under anæsthesia. On vaginal examination there was no obvious projecting growth, the only sugges-

tive point on gentle examination being that the external os was unusually patulous. On pressing the finger firmly into the os a soft friable growth could be felt occupying the cervical canal. The uterus was drawn down to the vulva with volsellæ, and came down easily. The vaginal fornices seemed to be free from any infiltration. On examining bimanually no extension of the disease into the broad or utero-sacral ligaments could be made out.

Operation (October 13th, 1894).—The patient was anæsthetised with ether, and placed in the lithotomy position. Vaginal hysterectomy was then performed. Wells's forceps were used for securing the vessels. The vagina was packed with iodoform gauze, and the upper part of the gauze plug was pushed into the pouch of Douglas. The gauze and forceps were removed ninety-six hours after the operation, but no vaginal douches were given till the 24th, when she was ordered vaginal douches of weak iodine water twice daily. The temperature after the operation did not rise above $100\cdot4^{\circ}$, and the patient made an uninterrupted recovery.

Examination of the uterus after removal.—The body of the uterus was not appreciably enlarged, and, so far as could be judged by the naked eye, the disease had not extended into it. An interesting point, however, was that at one part of the fundus there was a white nodule of circular shape, the size of a pea, in the thickness of the uterine wall. Microscopical examination of this nodule subsequently showed that it was a small fibromyoma.

Sections of the cervix were prepared by the Clinical Research Association, and were reported on by Mr. Targett as follows :—"The substance of the cervix uteri is infiltrated with branching processes of squamous-celled epithelioma. There are a few cell-nests, and some of the larger processes show granular degeneration of the central cells. The section is suitable for drawing."

Two drawings of the section accompany this paper.

I saw this patient last on March 15th, 1900, at the

London Hospital, and examined her. The scar was quite healthy, and she was herself perfectly well. The interval since the operation in this case is, therefore, five years and a half.

CASE 23 (with three illustrations).—Mrs. H—, 55 years of age, was brought to see me by Dr. Garry Simpson, of East Acton, on May 17th, 1895. She had had six children, the last seventeen years previously, and two miscarriages, both before the date of the last confinement.

The menopause occurred at the age of fifty. Since then she had “seen nothing” till August, 1894. Since that time there has been a constant red discharge from the vagina, and for the last two or three weeks the discharge has been offensive. She felt ill and worn out at the time when the vaginal discharge commenced. For some time she has suffered from continuous pain in the lower abdomen; the date of its commencement could not be fixed. Of late she has been getting rapidly thinner.

On examination of the abdomen nothing abnormal could be felt. *On vaginal examination* a large “cauliflower” growth was felt springing from the vaginal portion of the cervix; the uterus seemed freely movable, and the body of the uterus seemed considerably larger than normal in a woman past the menopause. She went into Fitzroy House for the purpose of having the uterus removed, and during the time—a few days—she was there before the operation the temperature at night was 100°, or thereabouts, as a rule.

Operation (June 1st, 1895).—Vaginal hysterectomy was performed in a manner similar to that already described in the preceding cases. At one stage of the operation the whole field became suddenly inundated with horribly foetid pus. This came from the body of the uterus, and had been let out by a laceration about the junction of the cervix and body caused by traction with

the volsella. The pus had previously been pent up in the body of the uterus by occlusion of the os internum, constituting the condition known as *pyometra*. In spite of the fact that the whole region of the wound was fouled with this foetid pus the convalescence was quite uninterrupted.

This patient writes to me every year on the anniversary of her operation. As she is now living in Australia, I do not actually get the letter for some weeks afterwards. I have had a letter from her this year (1900) dated June 11th, and in it she says that she still continues quite well. The uterus shows well the enormous distension of the cavity of the body by the pus referred to above, and it also shows an extensive cauliflower-like growth from the cervix extending up the cervical canal to the internal os. A portion of the cervix was sent to the Clinical Research Association for microscopical examination, and Mr. Targett reported on it as follows:—"The growth from the cervix uteri is a squamous-celled epithelioma of the papillary type. There is a considerable amount of keratoid change in the epithelial processes."

A drawing of the uterus and two drawings of the section accompany this paper.

CASE 24 (with two illustrations).—Mrs. A. H—, a married woman aged 48, was admitted under my care into the London Hospital on the 21st of August, 1895, at the request of Dr. Aspland.

Previous history.—She had been married twenty-one years, but had been a widow for fifteen years. She had never been pregnant. The catamenia appeared when she was about fifteen years old. She had dysmenorrhœa while she was single, but suffered much less from this after her marriage. The menopause occurred about three years ago.

Present illness.—She has not been well for some time. For the last seven or eight months she has had a yellow vaginal discharge, which has been mixed with blood more

often than not. She had no pain when the discharge began, but for the last two months she has had pain in the left iliac region occasionally. It has gradually become more severe, and is also felt at times across the lower abdomen, and in the lower part of the back. It was clear, however, that a blood-stained discharge, and a feeling of not being in good health, were for some months the only symptoms. She said on admission that she felt weaker than formerly, but did not think she had lost flesh. Her appetite latterly had not been good.

State on admission.—Nothing abnormal was detected on examination of the abdomen. On vaginal examination some blood-stained discharge was seen about the vaginal orifice, and in the vagina. It was not offensive. The uterus was found to be strongly retroverted; the vaginal portion of the cervix was rather soft, but otherwise normal. The uterus was easily replaced with the sound, which passed three and a quarter inches. On bimanual examination the uterus was found to be freely movable, and the body of it uniformly enlarged.

On August 23rd two small laminaria tents were put into the cervix, and on the next day the dilatation of the cervix was completed under anæsthesia with Hegar's dilators up to No. 19.

On passing the finger into the uterus a growth about the size of a walnut was found projecting from the posterior wall. It had a papillary surface, and was very friable, portions being easily broken off with the finger.

Operation (August 30th, 1895).—Vaginal hysterectomy was performed, the steps of the operation being similar to those adopted in the preceding cases. The pressure forceps and gauze were removed on the fourth day, and the patient made an uninterrupted recovery, leaving the hospital on the 23rd of September.

A portion of the growth was sent to the Clinical Research Association for microscopical examination, and Mr. Targett reported on it as follows:—"The growth from the body of the uterus is a soft glandular-celled

carcinoma, which has extensively infiltrated the muscular substance. The growth is of the columnar-celled type, though the tubular arrangement of the cells is not recognisable in the section."

I had a letter from this patient in May, 1900, to say that she was still quite well. The interval since the operation is therefore nearly five years.

CASE 26 (with two illustrations).—E. S—, a married woman aged 35, was admitted into the London Hospital under my care on April 6th, 1896. She was sent to me by Dr. Dunlop, of Purfleet.

She had been married sixteen years, and had had eight children, the last eight months old. She had had one miscarriage five years previously. Menstruation had always been natural.

History of the present illness.—She has found herself getting thinner lately, and has had a bad appetite. About three months ago she consulted a doctor on account of a yellow vaginal discharge, which she had had for some months. During the last two months the discharge has increased in quantity, and has been sometimes blood-stained. She has sometimes felt a pain in the lower abdomen and back, but pain did not seem to have been a marked feature in her present illness.

On April 10th she was examined under anæsthesia. On vaginal examination a "cauliflower-like" growth was seen springing from the vaginal portion of the cervix. The cervix could be easily drawn down to the outlet of the vulva with a volsella, and there was a good margin of apparently healthy tissue round the growth. The body of the uterus was not enlarged. The case seemed a suitable one for radical operation.

Operation (April 16th, 1896).—The patient was anæsthetised with the A.C.E. mixture. Vaginal hysterectomy was then performed, the method adopted being similar to that described in the preceding cases. The cut margin of the vaginal wall round the cervix was thoroughly

seared with Paquelin's cautery. The patient made an uneventful recovery, and went home on May 23rd, 1896.

A portion of the growth was sent to the Clinical Research Association for microscopical examination, and Mr. Targett reported on it as follows:—"This portion of the cervix uteri is thickly infiltrated with a soft squamous-celled epithelioma. There are no typical cell-nests, but a considerable number of cell-inclusions."

I have seen this patient several times since the operation; the last time I examined her was on October 12th, 1899, when there was no evidence of recurrence. I have also seen her on March 15th, 1900, when she said she was quite well, but I did not examine her on that occasion.

The interval since the operation in this case is, therefore, nearly four years.

CASE 29.—E. W.—. This was the case of primary sarcoma of the body of the uterus which formed the subject of a communication read before the Obstetrical Society of London, and which was published in vol. xxxix, p. 246, of the Society's 'Transactions.'

I only refer to it here for the purpose of recording her subsequent progress. I have seen her this year (1900), more than three years since the date of her operation, and she remains free from any sign of recurrence.

CASE 32.—Mrs. B—, aged 61, was sent to me by Dr. Pope, of Kington, Hereford. She had been married many years, and had had two children, the last twenty-eight years ago. I saw her on August 19th, 1897. She gave a history that nine years ago she had a fall, and that there had been more or less discharge, occasionally blood-stained, from the vagina since.

For the last two months there had been a constant discharge of blood every day. About a week before I saw her she had had a flooding, and was said to have lost about a pint of blood.

The menopause was said not to have occurred. She had not been getting thinner.

On August 20th she was examined under anæsthesia. There were several mucous polypi hanging from the cervix. The uterus was considerably enlarged, and was freely movable. As I did not think the mucous polypi could have accounted for the flooding mentioned above, I dilated the cervix and found a growth in the endometrium, a portion of which was detached for examination, and sent to the Clinical Research Association.

Mr. Targett reported on it as follows :—"The tissue from the body of the uterus is extensively infiltrated with a soft columnar-celled carcinoma. The tubular arrangement of the cells is generally well preserved, but in certain parts of the section the growth has undergone caseation."

The patient was accordingly advised to have the uterus removed.

Operation (September 4th, 1897).—Vaginal hysterectomy was performed as in the preceding cases ; the patient made a good recovery. I examined her last on July 7th, 1900. She was quite well, and there was no sign of recurrence. The interval since the operation is, therefore, two years and ten months.

CASE 33.—E. R—, a married woman aged 38, was admitted under my care into the London Hospital on September 4th, 1897.

She had been married nineteen years, and had had six children, the last three years ago, and two miscarriages, the last nine years ago.

The catamenia appeared when she was sixteen. The periods lasted four days, and were preceded by severe pain for three days, relieved on the appearance of the flow. She was never regular every four weeks till the last four years, but since the last confinement she has been fairly regular till recently.

Present illness.—This began nine weeks ago, when bleeding occurred, and continued for nearly five weeks.

After an interval of three or four days she had a flooding, and the loss of blood continued for eight days. She has had no pain, but has been getting thinner the last six weeks, and her appetite during that time has been poor.

On examination (September 5th, 1897) there was seen some blood about the external genitals. There was slight prolapse of both vaginal walls. There was marked enlargement of the anterior lip of the cervix, which was three or four times as thick from before backwards as the posterior lip. The growth on the anterior lip was raised, and in shape like a horseshoe with the convexity forwards. The growth bled easily on examination, and the sound pressed against it readily penetrated its substance. The body of the uterus was of normal size, and the uterus was freely movable.

Operation (September 6th, 1897).—Vaginal hysterectomy was performed, as in the preceding cases. She made a good recovery, and left the hospital on October 1st, 1897. Sections of the growth were prepared by the Clinical Research Association, and reported on by Mr. Targett as follows:—"The substance of the cervix uteri is extensively invaded with thick branching processes of squamous-celled epithelioma."

I last examined this patient on the 21st of June, 1900, at the London Hospital. She was then quite well, and there was no sign of recurrence. The interval since the operation without recurrence is therefore two years and nine months.

CASE 36.—E. M—, a married woman aged 46, was admitted into the London Hospital under my care on November 27th, 1897. She was sent to me by Dr. Honnywill, of Sutton, Surrey.

She had been married twenty-seven years, and had had four children, the last nineteen years ago. She had a flooding in the last week of August, 1897; the loss of blood continued for three weeks; there was then an interval of three weeks without any loss. The bleeding then began

again, and since then she has been losing blood almost constantly. She thought she had been getting stouter.

On examination it was found that the anterior lip was the seat of a growth about equal to half a crown in area. The growth was raised, and had an overhanging margin. The whole anterior lip from before backwards measured $1\frac{3}{8}$ inches, the posterior lip similarly only $\frac{3}{8}$ of an inch. The sound pressed against the growth readily penetrated its substance. Two small hard nodules were felt under the mucous membrane, one to the right of, and the other above the growth. The uterus was freely movable, and there was no evidence of extension of the disease to the broad ligaments or utero-sacral ligaments.

Operation (December 2nd, 1897).—Vaginal hysterectomy was performed, the method adopted being the same as in the preceding cases. Paquelin's cautery was used to sear the cut edges of the vaginal walls. There was slight pyrexia for the first fortnight after the operation, the highest point reached being 101° on the evenings of December 6th and 7th. After the 7th the temperature was normal. The patient did well, and went home on December 29th.

A portion of the cervix was sent to the Clinical Research Association for examination, and Mr. Targett reported on it as follows:—"The growth in the cervix uteri is a malignant adenoma, that is a columnar-celled carcinoma in which the cells preserve their tubular arrangement. The tissues are deeply invaded, and there is much small-celled growth around the proliferating tubules."

I last examined the patient on March 22nd, 1900, at the London Hospital. There was no sign of recurrence, and she expressed herself as feeling quite well. The interval since the operation is therefore two years and nearly four months.

CASE 39.—Mrs. J. R—, a married woman aged 36, was admitted under my care into the London Hospital on February 4th, 1899. She had had eight children, the last

eleven months prior to her admission. She had also had two miscarriages.

Her chief symptom had been a red vaginal discharge, which had been present fifteen months. At times the bleeding has been severe. She had also had pain in the left iliac region for ten months. She was suckling at the time of her admission.

On February 9th, 1899, the following note was made:—
“On the vaginal portion of the cervix is a raised and slightly irregular crescentic growth. It involves chiefly the left side of the vaginal portion immediately adjacent to the external os, and about two thirds of the posterior lip. It has an overhanging margin. About the middle of the anterior lip a little nodule, the size of a shot, is felt under the mucous membrane, the surface of which over the nodule is healthy. There is another similar nodule to the left of the first. There are also two nodules to the right under the mucous membrane of a similar character. The sound pressed against the crescentic growth enters its substance to the depth of an inch. The uterus is freely movable, retroverted, and its body not enlarged. There is no evidence that the growth has extended beyond the limits of the cervix.”

Operation (February 16th, 1899).—Vaginal hysterectomy was performed in the same way as in the preceding cases, Paquelin's cautery being used to sear the cut edges of the vaginal wall. The forceps were removed on the 20th. Slight bleeding occurred, so that the region of the wound was exposed with Sims's speculum. A pair of Wells's small pressure forceps was used to secure the bleeding point. This pair was left on two days, and then removed. No further bleeding occurred. The temperature only rose above 99·5° on two occasions; one was on the evening of the 20th, and the other on the evening of the 22nd, when the temperature was 100·2° and 100° respectively. The patient went home on the 15th March.

A portion of the growth was sent to the Clinical Research Association for microscopic examination, and

Mr. Targett reported on it as follows :—"The cervix of J. R— is very extensively invaded with squamous-celled epithelioma. Some of the epithelial processes are large, and the central cells have undergone granular degeneration. In others there are small cell-nests."

I examined this patient on July 26th, 1900, and found no sign of recurrence. The interval since the operation in this case is therefore one year and five months.

TABLE I.—*Forty Consecutive Cases of Vaginal Hysterectomy performed for Cancer of the Uterus.*

No.	Name.	Disease.	Date of operation.	Place.	Immediate result.	Remarks.	Age.	Remote result.
1	Mrs. R.	Cancer of body	Mar. 1st, 1886	London Hospital	Recovered	Ligatures only used to secure broad ligaments	58	Recurrence early in 1887. Died July 7th, 1887, 16 months after the operation. (Recurrence during the first year.)
2	Mrs. M. M.	Cancer of cervix	Oct. 4th, 1886	London Hospital	Died 48 hours after operation	The body of the uterus was moderately distended with pus (pyometra) which was set free in the course of the operation. There was no local complication such as hæmorrhage or peritonitis. The wall of the right ventricle was very thin and the heart fatty. The patient died from œdema of lungs and heart failure. Ligatures only to broad ligaments	62	—
3	Mrs. W.	Cancer of body	Nov. 7th, 1889	London Hospital	Recovered	Large soft "cauliflower" mass protruding through widely dilated os on admission. This was removed immediately, and vaginal hysterectomy performed on a subsequent occasion. None of the original "cauliflower" mass was kept for examination, as it was expected that the uterus would show more of the malignant tissue. This was not the case; so that, although on clinical grounds I am certain the soft friable mass removed was malignant, I cannot prove it by microscopical examination. Ligatures and pressure forceps to broad ligaments	42	Seen and examined Oct. 5th, 1893, <i>i. e.</i> nearly 4 years after operation, when there was no recurrence. Not seen since.
4	Mrs. D.	Cancer Oct. 23rd, 1890	Oct. 23rd, 1890	London Hospital	Recovered	Definite "cauliflower" growth on posterior lip of cervix. An interesting	41	This patient has been regularly seen up to and including the present

5	Mrs. J. T.	Cancer of body	March 17th, 1892	London Hospital	Recovered	feature was the occurrence of hæmaturia soon after the operation, lasting 5 days. Blood casts were present. Pressure forceps used to broad ligaments. Clinical signs and symptoms seemed conclusive as to malignancy, but the microscopic examination of the growth was not conclusive on this point Soft papillary growth projecting in cavity of body of uterus	44	This patient remained free from recurrence for more than 2½ years after the operation. Within the next 6 months, however, recurrence was ascertained by examination to have taken place. The case is fully recorded in Lond. 'Obst. Soc. Trans., vol. xxxvi, p. 374. (Recurrence during the third year.)	year 1900. There has been no sign of recurrence.
6	Mrs. B.	Cancer of cervix	June 16th, 1892	Private; with Dr. Russell, of Upton Park	Recovered	Little of the growth to be seen at the os uteri. The whole of the cervical canal up to the internal os was affected by the malignant growth; the cervix, in fact, converted into a thin shell lined with the growth	59	Good health for 2½ years after operation, then recurrence took place in the scar. (Recurrence during the third year.)	
7	Mrs. D.	Cancer of body	Feb. 1st, 1893	Private; with Dr. Stott, of Highgate, and Dr. Godson	Recovered	An example of "hard" cancer of the body of the uterus—the rarer variety	53	Still quite well—more than 7 years since the operation.	
8	Mrs. S.	Cancer of body	June 8th, 1893	London Hospital; patient sent to me by Dr. Best, of Dover	Recovered	An example of "hard" cancer of the body of the uterus	57	Still remains quite well. Last examined at the London Hospital July 19th, 1900. No sign of recurrence.	

No.	Name.	Disease.	Date of operation.	Place.	Immediate result.	Remarks.	Age.	Remote result.
9	Mrs. M.	Cancer of cervix	July 20th, 1893	London Hospital	Recovered	—	35	Remained well till the autumn of 1894, i. e. for a little more than a year after operation, then recurrence took place. (Recurrence during the second year.)
10	Mrs. C.	Cancer of cervix	Aug. 4th, 1893	London Hospital	Recovered	—	38	Died with recurrence Feb. 23rd, 1895, 18 months after the operation. (Recurrence during the second year.)
11	Mrs. T.	Cancer of body	Aug. 21st, 1893	London Hospital	Recovered	Soft mushroom-like growth. Perineum had to be incised. After one side of uterus freed it was brought into the vagina by forceps like small midwifery forceps	54	Came up to the hospital and examined Nov., 1893, 3 months after operation. Quite well and no sign of recurrence. All efforts to trace her since failed, as she had left the former address.
12	Mrs. K.	Cancer of cervix	Aug. 28th, 1893	London Hospital	Recovered	Soft mushroom-like growth affecting both lips. I thought at the time of operation that the left broad ligament and left utero-sacral ligament were somewhat infiltrated, and that probably therefore removal of the uterus had not removed all the diseased tissue	42	She was seen and examined on Nov. 30th, 1893, when there was distinct evidence of recurrence in the top of the vagina. I think most probably that the whole of the cancerous tissue had not been removed by the hysterectomy. (Recurrence during the first year.)
13	Mrs. C.	Cancer of cervix	Feb. 1st, 1894	Private	Died	Deep laceration of cervix left side, with firm cicatricial band of tissue running from it. Impossible to draw uterus down properly. A large vessel, probably the uterine artery, retracted out of reach, and though it was ultimately secured, a fatal amount of blood had been lost before this could be done	48	—
14	Mrs. F.	Cancer of cervix	Feb. 22nd, 1894	London Hospital	Recovered	A case of large "cunlifer" growth	52	Recurrence took place in August, 1894. (Recurrence during the first year.)

15	Mrs. J.	Cancer of cervix	Feb. 27th, 1894	Private; with Dr. Howse, then of Barking Road, E., now of Reading	Recovered	An extensive "cauliflower" growth. It was not till the patient was under an anæsthetic that the operation seemed worth trying	57	This patient remained free from any sign of recurrence till she died in February, 1900. Dr. Howse saw her several times in her last illness. She died of phthisis.
16	Mrs. B.	Cancer of cervix	Aug. 13th, 1894	Private; sent to me by Sir J. Williams	Recovered	The case was an example of large "cauliflower" growth of the cervix, and there was a round nodule the size of a cobnut in the left broad ligament. As the uterus could be easily drawn down it seemed worth trying whether removal of the uterus and also of the nodule would be effectual	43	I believe recurrence took place not very long after the operation, but I had not the opportunity of examining the case after she left London. (Recurrence during the first year.)
17	Mrs. S.	Cancer of cervix	Aug. 16th, 1894	London Hospital	Recovered	An example of the nodular and infiltrating form of the disease, the whole cervix being much thickened by it. Little or no growth projecting from the vaginal portion. At the operation, though the uterus was removed, I felt sure some infiltrated tissue was left, especially at the base of the bladder	32	(Recurrence during the first year.)
18	Mrs. R.	Cancer of cervix	Aug. 30th, 1894	London Hospital	Died on 7th day of peritonitis	An example of a large "cauliflower" growth. This patient was doing quite well till the pressure forceps were removed at the end of the second day; while this was being done she complained of sudden severe pain, and within a few hours symptoms of peritonitis appeared. I believe in removing the forceps the general cavity of the peritoneum was reopened by breaking down adhesions, and so infection occurred. Since this case I always leave on the forceps at least three days, and often four.	39	—

No.	Name.	Disease.	Date of operation.	Place.	Immediate result.	Remarks.	Age.	Remote result.
19	Mrs. S.	Cancer of cervix	Oct. 13th, 1894	London Hospital; sent to me by Dr. Howse, then of Barking Road, E., now of Reading	Recovered	Cancer involving the substance of the cervix; little projecting growth. It is interesting to know that the patient had been told at another hospital that the case was too far advanced for operation to be advisable.	52	Has been seen at frequent intervals at the hospital since operation, and has remained quite well; last examined March 15th, 1900.
20	Mrs. N.	Cancer of cervix	Dec., 1894	London Hospital; sent to me by Dr. Winn, of Hackney	Recovered	The removal of the uterus probably did not remove all the tissue infiltrated by the malignant growth.	45	(Recurrence during the first year.)
21	Mrs. G.	Cancer of cervix	Feb. 14th, 1895	London Hospital; sent to me by Dr. Cory, of Bourne-mouth	Recovered	Large "cauliflower" growth.	26	Recurrence took place within a few weeks of the operation. (Recurrence during the first year.)
22	Mrs. de B.	Cancer of cervix	Apr. 25th, 1895	London Hospital; sent to me by Dr. Cursham Corner	Recovered	Large "cauliflower" growth, especially affecting the posterior lip of cervix. Was doubtful at time of operation if all infiltrated tissue was removed.	35	Seen and examined on several occasions, the last time on June 16th, 1898, when the scar was quite healthy, and no sign of recurrence anywhere. Heard that she went to America, and died either at the end of 1898 or early in 1899; no

23	Mrs. H.	Cancer of cervix	June 1st, 1895	Fitzroy House; sent to me by Dr. Garry Simpson, of Acton	Recovered	Large "cauliflower" growth, the part of the growth in the cervical canal just reaching internal os. There was a very large pyometra, the pus being horribly offensive	55	<p>information as to cause of death. She was therefore known to be quite well over three years after operation, and if recurrence had been the cause of death it is very improbable that there should have been no sign of it on June 16th, 1898, if she died a few months after. It seems at least equally likely that she may have died from some independent cause.</p> <p>She has remained well since the operation. She has gone to Australia, but writes to me on each anniversary of her operation. The last letter, dated June 11th, 1900—five years after the operation,—says she was then quite well.</p>
24	Mrs. H.	Cancer of body	Aug. 30th, 1895	London Hospital	Recovered	A large papillary growth projected into the cavity of the endometrium	48	Well, May 18th, 1897; also heard from her in May, 1900, that she was quite well.
25	Mrs. T.	Primary tubercular ulceration of cervix; believed at the time of operation to be cancer of the cervix	Jan. 30th, 1896	London Hospital; sent to me by Dr. Haslip and Dr. Howell	Recovered	—	36	Heard she was quite well, November, 1899, when I operated on a friend of hers.

No.	Name.	Disease.	Date of operation.	Place.	Immediate result.	Remarks.	Age.	Remote result.
26	Mrs. S.	Cancer of cervix	Apr. 16th, 1896	London Hospital; sent to me by Dr. Dunlop, of Aveley, Purfleet, Essex	Recovered	Large "cauliflower" growth of cervix	35	Seen and examined January 11th, 1899; a granular mass, size of raisin, at upper part of vagina. Seen recently, the mass is just the same, and evidently not malignant. She was quite well. Quite well March 15th, 1900.
27	Mrs. A. W.	Cancer of cervix	Sept. 7th, 1896	London Hospital	Recovered	Nodule of malignant growth size cob-nut or thereabouts, posterior lip; anterior lip appeared quite healthy. Supra-vaginal amputation done first, then on slitting up cervix the growth found to extend up to internal os, so body of uterus was then also removed	45	She was well up till about October, 1899, over three years since her operation. Then she began to fail, and came to the hospital in February, 1900, with a mass size of fetal head in hypogastric region, fairly movable. Evident recurrence in the scar, and under the vaginal walls for some distance down. The glands in right groin much enlarged. She was looking cachectic; there was no projecting growth in region of scar, and only slight bleeding on examination. Very slight bleeding in October or November, 1899; for one day was the only spontaneous bleeding. (Recurrence during the fourth year.)
28	Mrs. I.	Cancer of body	Nov. 12th, 1896	London Hospital	Recovered	Secondary hemorrhage in third week after operation requiring application of Wells's forceps. Subsequent incontinence of urine found on filling the bladder not to be due to any vesical fistula, but to a ureteral fistula. This persisted for some three or four months and then healed spontaneously	49	Recurrence in the recto-vaginal septum during the first half of 1900. (Recurrence during the fourth year.)

29	Mrs. W.	Sarcoma of body	Feb. 11th, 1897	London Hospital	Recovered	See 'Obstet. Trans.' for full report of case	24	Quite well March 24th, 1900.
30	Mrs. E. S.	Cancer of cervix	June, 1897	London Hospital	Recovered	—	About 60	Recurrence took place about a year after the operation. (Recurrence during the first year.)
31	Mrs. B.	Cancer of cervix	Aug., 1897	London Hospital; sent to me by Dr. Francis, of Leytonstone Road	Recovered	—	60	Recurrence took place and she died May 19th, 1898. (Recurrence during the first year.)
32	Mrs. B.	Cancer of body	Sept. 4th, 1897	Private; sent by Dr. Pope, of Kingston, Hereford	Recovered	There were some mucous polypi attached to the lower part of cervix. The malignant growth was in the body	61	Last examined July 7th, 1900. No sign of recurrence.
33	Mrs. R.	Cancer of cervix	Sept. 6th, 1897	London Hospital	Recovered	—	38	Well June 21st, 1900.
34	Mrs. B.	Cancer of cervix	Sept. 23rd, 1897	London Hospital	Died three weeks after the operation of pneumonia	The only case I have had in which cancer of the cervix affected a pro-cident uterus. This patient was mentally feeble and very difficult to manage after the operation. At the post-mortem the pneumonia was said to be not septic, and the parts in the neighbourhood of Douglas's pouch were looking healthy	53	—
35	Mrs. H.	Cancer of cervix	Oct. 5th, 1897	Private; with Dr. McMillen, of Brixton	Recovered	A rather advanced case	52	Died of recurrence July, 1898, the scar remaining unaffected to the last. (Recurrence during the first year.)

No.	Name.	Disease.	Date of operation.	Place.	Immediate result.	Remarks.	Age.	Remote result.
36	Mrs. E. M.	Cancer of cervix	Dec. 2nd, 1897	London Hospital; sent to me by Dr. Honnywill, of Sutton	Recovered	—	46	Examined and found quite well March 22nd, 1900, more than three years since the operation.
37	Mrs. D.	Cancer of cervix	May 12th, 1898	London Hospital; sent to me by Dr. Jackson, of King's Lynn	Recovered	A rather advanced case with large conical ulcer	38	Recurrence took place March, 1899. She became much worse June 5th, 1899, and died soon after. (Recurrence during the first year.)
38	Mrs. S.	Cancer of body	May 14th, 1898	Fitzroy House; Dr. Johnston, of Maidstone	Recovered	—	About 50	—
39	Mrs. R.	Cancer of cervix	Feb. 16th, 1899	London Hospital	Recovered	Bleeding on taking off forceps, requiring application of one pair more for another two days	36	Quite well July 26th, 1900.
40	Mrs. E. K.	Cancer of cervix	April 27th, 1899	London Hospital	Recovered	The disease affected both the vaginal and supra-vaginal portions of the cervix	36	Recurrence early in 1900. (Recurrence during first year.)

TABLE II.—*Twelve Cases of Vaginal Hysterectomy for Cancer of the Uterus in which the Disease is known not to have recurred.*

No.	No. in Table I,— a consecutive series of 40 vagi- nal hysterecto- mies for cancer.	Disease.	Date of operation.	Place of operation.	Interval since operation without recurrence.	Microscopic characters of growth.
1	7. Mrs. D.	Cancer of body	Feb. 1st, 1893	Private; with Dr. Stott, of Highgate Road, and Dr. Godson London Hospital; sent to me by Dr. Best, of Dover	7 years and 4 months. I saw her on 18th of June, 1900	Extract from Clinical Research Association's Report (Mr. Targett):—"The growth may be described as a villous carcinoma."
2	8. Mrs. S.	Cancer of body	June 8th, 1893	Private; with Dr. Howse, then of Barking Rd., E., now of Reading London Hospital	7 years. Examined July 19th, 1900, at the London Hospital. No recurrence	Extract from Clinical Research Association's Report (Mr. Targett):—"A soft columnar- celled carcinoma."
3	15. Mrs. J.	Cancer of cervix, large "cauliflower"	Feb. 27th, 1894	Private; with Dr. Howse, then of Barking Rd., E., now of Reading London Hospital	6 years	Extract from Clinical Research Association's Report (Mr. Targett):—"The nature of the growth is undoubtedly squamous-celled epi- thelioma."
4	19. Mrs. S.	Cancer of cervix	Oct. 13th, 1894	Private; sent to me by Dr. Garry Simp- son, of Acton	5½ years (up to March 15th, 1900, when the patient was last examined)	Extract from Clinical Research Association's Report (Mr. Targett):—"The substance of the cervix is infiltrated with branching pro- cesses of squamous-celled epithelioma."
5	23. Mrs. H.	Cancer of cervix, large "cauliflower." Pyometra	June 1st, 1895	Private; sent to me by Dr. Garry Simp- son, of Acton	5 years. Heard from her in letter dated June 11th, 1900, that she was quite well	Extract from Clinical Research Association's Report (Mr. Targett):—"Solid branching columns of squamous-celled epithelioma are shown infiltrating the substance of the cervix uteri."
6	24. Mrs. H.	Cancer of body	Aug. 30th, 1895	London Hospital	4½ years (up to May, 1900, when I heard from her that she was quite well)	Extract from Clinical Research Association's Report (Mr. Targett):—"The growth from the body of the uterus is a soft, glandular- celled carcinoma, which has extensively in- filtrated the muscular substance."

No.	No. in Table I.— a consecutive series of 40 vagi- nal hysterecto- mies or cancer.	Disease.	Date of operation.	Place of operation.	Interval since operation without recurrence.	Microscopic characters of growth.
7	26. Mrs. S.	Cancer of cervix	April 16th, 1896	London Hospital; sent to me by Dr. Dunlop, of Purfleet	Nearly 4 years. I saw her on March 15th, 1900, and she was quite well	Extract from Clinical Research Association's Report (Mr. Targett):—"This portion of the cervix uteri is thickly infiltrated with a soft squamous-celled epithelioma."
8	29. Mrs. W.	Sarcoma of body, "deciduoma malignum"	Feb. 11th, 1897	London Hospital	More than 3 years (up to March 24th, 1900, when I last examined her)	For full report of this case see Lond. 'Obst. Soc. Trans., vol. xxxix, for 1897, p. 246.
9	32. Mrs. B.	Carcinoma of body	Sept. 4th, 1897	Private; with Dr. Pope, of Kingston, Hereford	2 years and 10 months (up to July 7th, 1900, when I last examined her)	Extract from Clinical Research Association's Report (Mr. Targett):—"The tissue from the body of the uterus is extensively in- filtrated with a soft columnar-celled carci- noma."
10	33. Mrs. R.	Cancer of cervix	Sept. 6th, 1897	London Hospital	2 years and 9 months (up to June 21st, 1900, when I last examined her)	Extract from Clinical Research Association's Report (Mr. Targett):—"The substance of the cervix uteri is extensively invaded with thick branching processes of squamous- celled epithelioma."
11	36. Mrs. F.M.	Cancer of cervix	Dec. 2nd, 1897	London Hospital	2 years and nearly 4 months (up to March 22nd, 1900, when I last saw her)	Extract from Clinical Research Association's Report (Mr. Targett):—"The growth in the cervix uteri is a malignant adenoma,—that is, a columnar-celled carcinoma in which the cells preserve their tubular arrangement."
12	39. Mrs. R.	Cancer of cervix	Feb. 16th, 1899	London Hospital	1 year and 5 months (up to July 26th, 1900, when I last examined her)	Extract from Clinical Research Association's Report (Mr. Targett):—"The cervix is very extensively invaded with squamous-celled epithelioma. Some of the epithelial pro- cesses are large, and the central cells have undergone granular degeneration. In others there are small cell-nests."

DISCUSSION.

Professor BRIGGS (Liverpool).—The results of the operation of vaginal hysterectomy produced this evening by the author of the paper would tend to establish the position of the operation much more than, for instance, my own. Since March 24th, 1889, up to the present I have operated on 83 cases of malignant disease of the uterus (4 deaths). Of the 83 growths 3 were sarcomata of the cervix, 65 carcinomata of the cervix, and 15 carcinomata of the body of the uterus. To adopt Dr. Lewers's date limit, up to the end of April, 1899, I have 57 cases to report upon, and amongst them the deaths from operation were 3, including 1 from embolic septic pneumonia on the thirty-fifth day. Of the 57, 10 were cases of corporeal cancer, showing 5 patients apparently cured, and 46 were cases of cervical cancer. Of these my reports at present are incomplete, but in 5 cases complete cures are apparently promised. Adding to these the 1 case of cervical sarcoma operated upon over ten years ago, there are perhaps only 11 out of 57 patients completely cured by operation. Delay in diagnosis has always to be remarked upon, and if patients object to examination use could be made of the ordinary medico-legal tests for blood in the intermenstrual discharges. I have brought with me water-colour drawings, from which you will gather that many of my cases were in a more advanced stage than Dr. Lewers's; but I intend to complete my clinical reports in the near future, and on one or two occasions in the past I have been gratified to find the results better than I had anticipated, owing to broad ligament indurations not having proved malignant. Recently my assistants and I have analysed 100 malignant growths of the uterus in the laboratory of University College, Liverpool; 74 were carcinomata, 6 sarcomata, and 20 malignant adenomata. Amongst the 83 included in my series of operations already referred to, I may mention that the cases of malignant adenoma were 4 of the cervix and 11 of the body of the uterus. The cases of malignant adenoma have proved more amenable to radical treatment. I have adopted similar methods to those of Dr. Lewers in diagnosis and in operating. I desire to congratulate Dr. Lewers on results which at present seem to be better than my own. Amongst even advanced cases of corporeal cancer the proportion of cures is high, but amongst the cervical carcinomata cures have rarely been obtained. It is amongst the latter that Dr. Lewers's results are so strikingly good when compared with my present acquaintance with my own.

Dr. A. ROYTH.—Those who know how difficult it is to follow

up hospital cases will realise how indefatigable the author must have been to obtain the histories up to date. I also have done 41 cases of vaginal hysterectomy, 15 being of the body and 26 of the cervix. That seems a large proportion of cases in which the body was affected, resembling in this respect the author's figures, he having had 11 out of 40. I imagine the reason why one gets a larger proportion of cases of cancer of the body of the uterus is that they come earlier under observation. The history in such cases of cancer of the body is usually that there has been a period of amenorrhœa after the menopause, and that then hæmorrhage comes on which attracts attention, and this leads to an examination being made, whereas the disease at an early stage in the cervix may escape notice for a time. Moreover pain is a much more marked symptom in cancer of the body. This is also probably one reason why recurrence does not take place so often in cancer of the body. My mortality was also three. One, on the sixth day, was a case of cancer of the cervix in an alcoholic patient who died of exhaustion, and two of cancer of the body, both dying suddenly when I thought they were convalescent, one on the sixteenth day and the other on the twenty-fourth day, acute mania having developed on the sixteenth day. One of the cases of cancer of the body was a malignant papilloma, and two malignant adenoma with one sarcoma, removed by the vagino-abdominal method. The cases of cancer of the body did not as a rule recur as early as in cancer of the cervix. One patient is alive two and a quarter years after the operation, one three years and a half, another four years after. Of twenty-six cases of cancer of the cervix with one death, recurrence was less satisfactory. Four or five cases recurred within a year, one two years after, another three years after, and one, a very disappointing one, seven years after. I saw this latter patient every six months, and I thought she had permanently recovered. Recurrence took place in the cicatrix and she soon died. One patient operated on in 1892 and another in 1895 are still free from recurrence. From the figures brought forward to-night it is evident that the opinion expressed by some that this operation is useless cannot be true. I am surprised that the author uses forceps for his broad ligaments. He began with using ligatures and finished with forceps, whereas I began with clamps and turned to ligatures. It seems to me more surgical to use ligatures than forceps, as the latter leaves a large slough to come away, the temperature is higher, and the discharge very offensive. If I have to use forceps I generally leave them on thirty-six hours. Like the author, I never attempt to operate if I find that the uterus has indurations in the broad ligament, or if the utero-sacral folds or bladder are implicated. If in

doubt I get permission to do the operation if possible, and after drawing the uterus down I open Douglas's pouch first, instead of making an anterior incision and separating the bladder, which is usually the first step. I then pass the finger into Douglas's pouch, and am able to accurately estimate whether the disease has spread into surrounding tissues. If it has I prefer not to remove the uterus. How long does the author keep the gauze in? At first I took it out in two or three days, but I find now that I get better results if I leave it in seven or eight days. Of 15 cases of vaginal hysterectomy for cancer of the body two died (sixteenth and twenty-fourth day) suddenly. Of the others two recurred within one year, two within two years, and one in three years. Of those who are still alive without recurrence after two years from the operation, one has lived six and a half years, one four and a quarter, and two two and a half years. Of 26 cases of cancer of the cervix one died on the sixth day (chronic alcoholism). Of the others six recurred within one year, three within two years, and one after six and a quarter years. Of those who have survived more than two years and are now without recurrence, one has lived five years, two over four years, three over three years, and two over two years.

MR. BOWREMAN JESSETT.—I have published 107 cases in the 'British Gynaecological Journal' of hysterectomy for cancer, and have now performed the operation in over 150 cases. My conclusions support those advanced by the author, viz. that a very fair percentage of cases of carcinoma of the uterus, if seen in an early stage, can be what may practically be called cured, i.e. patients are relieved from their sufferings for a period of from three to nine years. The whole gist of the author's paper is early diagnosis, and I am quite sure that this is the practical conclusion of any one who has had much experience of these cases. Patients go about a long time with purulent or sanguineous discharges without consulting a doctor, or the doctor contents himself with prescribing some astringent lotions, etc. Many patients object to being examined, but I contend that it is the duty of the medical man, to himself and to his patient, to insist upon a thorough examination in all cases where there is any abnormality of the uterine secretions, especially in women at or after the menopause. If such an examination were made the disease would be recognised at a much earlier stage than is often done at present, and if treated surgically such patients would stand a very much better chance of obtaining relief. The operation is not of itself dangerous, the mortality being at the outside from 5 to 8 per cent. Some people say vaginal hysterectomy is a very simple and easy operation, but though I have seen some very simple operations where the

uterus could be drawn completely outside the vulva, and the fornices readily opened, I have also seen others of extreme difficulty. The difficulty in many cases of getting the uterus down is very great. I have in nearly every instance used ligatures for securing the vessels, which seems to me a far more surgical procedure than adopting forceps for the same purpose. The only mishap I have ever had was after using forceps. The principal difficulties are that you often get adhesions of the uterus in Douglas's pouch to the rectum, the sigmoid flexure, or small intestine, and these adhesions are very difficult in many cases to separate. In one case in which I used forceps there were very strong adhesions to the upper part of the rectum, and I had some difficulty in catching up the ovarian artery. Everything went well for ten days, when a large slough came away and a recto-vaginal fistula formed. I endeavoured to close this later, and found it quite at the top of the vagina, and it was next to impossible to do so. Another danger is that of tearing into the bladder, which happened to me in one case which appeared on examination to be one of the simplest of cases, but on opening the anterior fornix I found that the disease had extended from the cervical canal into the cellular tissue and had infiltrated the bladder, and in trying to separate it the bladder was torn and a fistula formed which caused much trouble later. The ureters are also much more likely to be injured by forceps than by the ligatures. I would ask the author whether he employs Doyen's plan of splitting the anterior wall of the uterus and drawing it down. In cases of a very large uterus I always employ that procedure, as it undoubtedly facilitates matters materially. The author has not alluded to the malignant degeneration of fibroids, and I should like to know his experience thereof. Such cases are in my experience the most favourable for operation. I cannot agree with Dr. Routh that cancers of the body of the uterus are the earliest met with and the easiest to diagnose. I think these are very difficult to recognise; indeed, I cannot suggest any way of doing so for certain in the early stages, excepting by anæsthetising the patient, dilating the uterus, and curetting and examining the fragments microscopically. Even in some of these cases pathologists have reported that there was no malignancy, yet they have proved to be malignant, a small growth being found in the fundus which might escape the curette or not be included in the scraping submitted to the pathologist for examination, thus showing how difficult these cases are to diagnose in the early stage. Then as to hæmorrhage being one of the early symptoms after the menopause: that is true, but there is often a still earlier symptom, and that is slight hæmorrhage after coitus, which in my experience is not uncommon. In the post-mortem

room I have been struck by the very small number of cases in which the sacral and iliac glands are involved, and this induced me to urge patients to submit to hysterectomy. I would point out that the treatment of the peritoneal flaps is rather an important point. They have a tendency to curl up after division, and if you do not take the trouble to pull these flaps down you may get an accumulation of blood-clot which may give rise to trouble. I make a point of drawing both flaps well down. Personally I do not withdraw the gauze for five, seven, or even ten days.

MR. CHARLES RYALL.—As to the frequency with which recurrence takes place in epithelioma of the cervix in contradistinction with that of the body of the uterus, we all know how easy it is to transplant malignant disease in a patient already suffering therefrom. It has struck me that at the time of the operation we are very likely to infect the wound. In carcinoma of the fundus the disease is surrounded by a certain amount of healthy tissue, which is not the case in the cervix. Pieces of the growth may be broken off and left in the wound, thus leading to recurrence. The author's large percentage of recoveries may be accounted for by the use of the actual cautery, which is a strong antiseptic. This danger of infection was very strongly brought before me not long since in a case of cancer of the fundus, for which *abdominal* hysterectomy was performed. During the operation I tore the uterus across at the internal os, so that the fundus had to be removed first. A year later that patient came with recurrence in the vaginal *cul-de-sac* causing pressure on the rectum. She also had a swelling in the abdominal scar the size of a large walnut. Colotomy was done, and I took advantage of the opportunity to examine the peritoneal scar, and there were no adhesions at all. I removed a piece of the growth in the scar, and it was found to be identical with the original growth in the uterus of a year before. I agree with Mr. Jessett that Doyen's method is of great assistance when the uterus is large. As to palliative cases, the author goes perhaps a little too far. He gives us 27 per cent. of cures, so that the other 71 per cent. can only have been palliative operations. In cases of cancer of the uterus which cannot all be got away, if an operation only rids the woman of the foul discharge for six months, it is only fair to give her the benefit of it, without thinking only of our statistics.

DR. WALTER TATE.—In eleven of Dr. Lewers's cases of cancer of the body of the uterus there was freedom from recurrence in five, or about 50 per cent., and in twenty-eight of cancer of the cervix there was freedom in 25 per cent. That corresponds to the results obtained by most operators. I have myself operated on upwards of seventy cases by vaginal hysterectomy for cancer

of the cervix and body, but I do not agree with the author in looking upon this operation as only suitable for cases in which the disease is definitely localised in the uterus. So far as I have seen, cases in which there is a slight amount of thickening in the broad or utero-sacral ligaments, in which patients suffer from discharges and severe hæmorrhages, may be very greatly relieved by removing the uterus, provided the thickening is slight and not sufficient to interfere with the uterus being brought down near the vulva. Dr. Martin, of Berlin, made the remark some years ago that where vaginal hysterectomy is an easy operation it is a good operation, and where it is difficult it is a bad operation. In some of these cases where there has been thickening of the broad ligament the freedom from recurrence has been much longer than in cases in which *a priori* the prospect was much more favourable; therefore the operation is desirable and justifiable as a palliative measure even when to all appearance it is not likely to be permanently curative. I have lost five cases out of the seventy, but only two out of the last fifty.

Dr. CHAMPNEYS.—The author's statistics quite negative the view that vaginal hysterectomy is an operation which should not be performed, or which is of no use. Of course cancer is altogether a very disappointing disease, and even when the prognosis before operation seems excellent we must expect a great many disappointments. I have not my figures with me to-night, but I have several patients going about apparently quite well several years after operation. The author did not allude to one of the most valuable forms of examination, viz. *per rectum*. This enables us to reach much higher than *per vaginam*, and you can feel the back of the uterus and the utero-sacral ligaments. Like him I began with ligature, and have finally adopted the forceps, but I do not use Wells's forceps, because the serrations seem to me to be in a wrong direction; they are transverse, which is parallel to the direction of the vessels; so I have mine made with diagonal serrations. With regard to the operation in advanced disease I agree with what he has said. If the disease has gone beyond the uterus I do not think the operation should be done. In many cases it is very difficult to make out whether it has or not. There may be some induration or want of elasticity in one or other broad ligament, and one cannot know beforehand whether it is due to inflammation or to malignant infiltration. Very often in such cases anæsthesia will clear up the doubt, for the induration may become plastic, so to speak, and there is an absence of nodules; in such cases I give the patient the benefit of the doubt. The difficulty and danger of the operation is certainly very much enhanced when the broad ligament is involved. I should like to know what

evidence Mr. Jessett has of fibroids undergoing malignant degeneration. That fibroids may coincide with malignant disease merely shows that a woman with fibroids is not protected against the common fate. It must of necessity be very difficult to prove that an innocent growth takes on malignant degeneration, and I have only met with one case in which I suspected that this might have been the case.

Dr. HERBERT SPENCER.—I cannot speak too highly of the truly admirable way in which the author has tabulated and arranged his cases. I am particularly interested in the instance of implantation mentioned by Mr. Ryall. Many examples of this cancer-implantation ("Impfmetastase") are on record. In order to prevent its occurrence I have usually performed high amputation in cancer of the cervix instead of vaginal hysterectomy, the operation being performed of late years entirely by the actual cautery with extremely good results. I have not lost a case in thirteen years. It is an idea I owe to Dr. Byrne, who in some 400 cases has only had one death. It is practically a safe operation. I believe that the improvement in this operation will lie in the fact of using the cautery for the removal of the cervix or uterus. I have used it for the removal of the whole uterus *per vaginam*. With modern improvements in cautery forceps I think we shall be able to remove the uterus with a very small mortality, and that the searing of the tissues prevents the implantation of cancer on the raw surfaces. It is remarkable that Jacobs, of Brussels, with a very large experience, has not had one case survive for two years. Most foreign operators cut up the uterus with knife or scissors, but this, I think, ought not to be done in cancer. It is especially in fibroids that it renders excellent service, but in cancer the less we cut up the better. I have a number of cases remaining well five years or more after operation. I cannot to-night give figures, but I am convinced that the use of the cautery is an extremely valuable method of operating.

Dr. LEWERS.—I could not follow Dr. Briggs's figures, but I hope he will publish his cases in a form we can refer to. I must congratulate Dr. Routh on his low mortality, which is the same as my own. I should think we get cases of malignant disease of the body more frequently in an earlier stage than cases of cancer of the cervix. It is difficult to account for this so far as the examination of the case goes, for a well-marked projecting growth on the vaginal cervix is much more easily recognised by the doctor examining than a malignant growth in the body of the uterus, which can only be recognised by dilating the cervix. My own belief is that in cases of cancer of the cervix and probably of the body there are no symptoms at all in the early stages. It is not a question which is the earlier

symptom, but I believe there is a stage in which the growth can be recognised by local examination at a time when there are no symptoms. I remember the case of a patient from whom I removed a fibroid polypus projecting through the cervix; she was of a nervous temperament, and she came at intervals afterwards to see that she was all right. On one occasion there was a well-marked raised patch on the anterior lip of the cervix, not the spot where the polypus was attached, and I felt convinced that it was malignant. Sir John Williams took the same view, and the case was operated on; the growth was found to be malignant. At that time the patient complained of nothing whatever—no hæmorrhage, either after coitus or at other times, and she had no discharge. Therefore we may take it that in the early stages there may be no symptoms, and that is one reason why we get these cases as a rule so late. A case of recurrence after seven years, such as that mentioned by Dr. Routh, is particularly disappointing. I may repeat what I have said in my paper that these cases do not include all my cases of operation for cancer. Many of my earlier cases were operated upon by high amputation of the cervix, and some of my best cases, as regards the after results, belong to that series; possibly because, being the first cases, they have had longer to run. In December, 1892, I read a paper before this Society comprising a number of these cases; among that series I can say positively that there are two patients, one thirteen and one twelve years after operation, who are now perfectly well. In regard to forceps *versus* ligatures, I confess that I should like to use ligatures. I used them in the first three cases. They are much more comfortable for the patient, and the use of the forceps entails much discomfort; but as a rule I have been well satisfied with forceps. My experience of ligatures is that it is extremely difficult to apply them in such a way as to secure us against hæmorrhage. The uterus is drawn down and the parts are put on the stretch, and I have seen cases where, when the operation was supposed to be over, the ligatures slipped off with profuse hæmorrhage, and it has been necessary to use the forceps after all. I, too, have altered the serration of the forceps, and use a pattern with oblique grooves as mentioned by Dr. Champneys. I generally leave the gauze in for four days,—in fact, until I take out the forceps. With ligatures it is quite a different matter, and it then may be good practice to leave it in longer. You cannot do this if you use forceps, because they cannot be removed easily with the gauze in position. In reference to the formation of a fæcal fistula, it often happens that a piece of intestine is infiltrated, and possibly that might have been so in Mr. Jessett's case. I agree with Dr. Champneys that it is important to preserve the integrity of the specimen, because you

lose the outline of the malignant growth if you cut it up, and may easily leave some of it behind. With regard to cancer of the body of the uterus, by the time the uterus is so large as to offer an obstacle to removal by the vaginal route, it is usually beyond the reach of surgical aid. In such cases it is best to do abdominal hysterectomy, which seems to me better than splitting the uterus. I know nothing at all about fibroids becoming malignant. I have been looking out for such cases for years, and I cannot recall a single instance in which I have satisfied myself that such has been the case; for the present, therefore, I remain sceptical. Some Continental observers assert that the glands are always affected in all the cases of cancer of the cervix that come to us, but my results, I think, show conclusively that this cannot be the case. I approve of the idea of drawing down the peritoneal flaps, and shall bear Mr. Jessett's remarks in mind. I decline to admit that it can be right to perform vaginal hysterectomy on advanced cases of cancer when there is no prospect whatever of permanent relief. Unless we are going to get something better than a few months of comparative comfort I really do not think it is worth while. Recurrence usually takes place in the scar and quite exceptionally in the glands, so that it is rare for the patient to be relieved from the misery of a constant discharge. I myself never do vaginal hysterectomy as a palliative measure. It is true that many of my operations proved to be nothing more, but I did not operate with that object in view. I have never removed the uterus entirely by the cautery, as mentioned by Dr. Spencer, but I shall be prepared to try it.

No. 7. (Table I.)

Cancer of body of uterus. Vaginal Hysterectomy, February 1st, 1893. The patient was seen on June 18th, 1900, and expressed herself as feeling quite well.

A.—Peritoneum. B.—Cut surface of uterine wall. C.—Growth in endometrium (villous carcinoma). D.—Cervical canal.

No. 7. (Table I.)

Fig. A depicts the general characters of this growth under a low power. In the muscular tissue of the uterus there are spaces of various sizes from simple gland-like tubules to cysts visible to the naked eye, and filled with a complicated epithelial new growth. This growth has resulted from proliferation of the columnar epithelium lining the tubules, and has assumed in part the form of intra-cystic villous papillomata. At the periphery of the largest space the malignancy of the growth is shown by extension of solid processes into the adjacent uterine tissue. (Mr. Targett's description for the Clinical Research Association.)

Bale and Danielsson, Ltd.

In **Fig. B** (high power) two of these intracystic papillomata are represented. In the vascularity and delicacy of their structure they closely correspond with the fimbriated papilloma of the urinary bladder. The growth may be described as a villous carcinoma. (Mr. Targett's description for the Clinical Research Association.)

Bate and Danielsson, Ltd.

No. 8. (Table 1.)

Primary carcinoma of the body of the uterus. Vaginal Hysterectomy, June 8th, 1893. The patient was seen and examined at the London Hospital on July 19th, 1900, and found to have no sign of recurrence.

A.—Peritoneum. B.—Cut surface of uterine wall. C.—Growth in endometrium "a soft columnar-celled carcinoma." D.—Cervical canal.

No. 8. (Table I.)

Fig. A (low power) represents a portion of uterine wall composed of interlacing bundles of muscular tissue, and extensively infiltrated by a soft columnar-celled carcinoma. The cells are arranged in tubules and elongated spaces, which have a definite lumen, and the stroma consisting of uterine muscle is very scanty. (Mr. Targett's description for the Clinical Research Association.)

In Fig. B (high power) the columnar character of the cells is shown, and although the tubules and alveoli everywhere exhibit a central cavity or lumen, yet there are abundant signs of epithelial proliferation, and the cells encroach on the central space. In the well-marked tubular structure of the growth, this specimen resembles the so-called malignant adenoma. (Mr. Targett's description for the Clinical Research Association.)

No. 15. (Table I.)

Vaginal Hysterectomy, February 27th, 1894. No return six years after the operation, when the patient died of phthisis.

- A.—Epithelial surface of vaginal portion of cervix.
- B.—Growth "squamous-celled epithelioma."
- C.—Portion of peritoneum on body of uterus.

No. 19. (Table I.)

Fig. A (low power) shows epithelial alveoli of varying size and shape, which are invading the muscular substance of the uterus. The cells are arranged in a single row of columnar epithelium at the periphery, and towards the centre they become much more irregular and compressed. A few cell-nests are seen, and they are numerous throughout the section. (Mr. Targett's description for the Clinical Research Association.)

Fig. B (high power). In this sketch one alveolus is represented made up of typical squamous epithelial cells. To the left a whorl of cells is seen in process of formation of a cell-nest. The vessels in the adjacent uterine substance are dilated. (Mr. Targett's description for the Clinical Research Association.)

No. 23. (Table I.)

Vaginal Hysterectomy, June 1st, 1895. I heard from the patient (who is in Australia) in a letter dated June 11th, 1900, that she was quite well.

- A.—Cavity of body of uterus enormously dilated. It contained foetid pus (pyometra) which escaped during the operation.
- B.—Cut surface of wall of body of uterus.
- C.—Growth in cervix, "squamous-celled epithelioma."
- D.—Internal os.
- E.—Peritoneal surface of uterus.
- F.—External os.

No. 23. (Table I.)

Fig. A (low power). Solid branching columns of squamous-celled epithelioma are shown infiltrating the substance of the cervix uteri. Large cell-nests are present, as well as areas of granular degeneration in the centres of the epithelial processes. (Mr. Targett's description for the Clinical Research Association.)

Fig. B (high power). The squamous character of the epithelial cells is here represented, and in the centre keratoid and other degenerative changes have taken place. The cells have separated from the stroma by shrinkage. (Mr. Targett's description for the Clinical Research Association.)

No. 24. (Table I.)

In Fig. A (low power) is represented a soft glandular-celled carcinoma invading the body of the uterus. The cells are arranged in elongated alveoli, and a strand of muscle tissue crosses the section. At the highest point in the field the growth has undergone granular degeneration and necrosis. (Mr. Targett's description for the Clinical Research Association.)

Bale and Danielsson, Ltd.

In Fig. B (high power) the cells are seen to be of moderate size and fairly uniform shape. There is no indication of a tubular arrangement, and the small round nuclei indicate leucocytes invading the degenerate areas. (Mr. Targett's description for the Clinical Research Association.)

No. 26. (Table I.)

Fig. A represents a low power view from the edge of the growth. In the cervix uteri, at the top, is a portion of vaginal mucous membrane, and beneath this comes a broad strand of submucous tissue and bundles of unstripped muscle. The vessels are distended, and many foci of small round inflammatory cells are seen in this layer. At the bottom of the field are depicted the solid elongated processes of the squamous-celled epithelioma. (Mr. Targett's description for the Clinical Research Association.)

Bale and Danielsson, Ltd.

In Fig. B a portion of the growth under a higher power is represented. The cells are large, but their outlines are very indistinct, so that merely their large round or oval nuclei with nucleoli are recognisable. No cell-nests are present, but a few larger cells may be seen which exhibit the so-called "cell-inclusions." One of these is represented in the drawing. The stroma consists of a few strands of cellular tissue in which the vessels run. (Mr. Targett's description for the Clinical Research Association.)