employment of a daily performance test makes available a considerable amount of information on the manner of development of disability, as well as on its causes and duration.

The respective parts played by skin burns and systemic poisoning in the development of disability has already been pointed out, and it has been suggested that with high dosages subjects protected only by respirators would rapidly become disabled by a combination of systemic intoxication, generalized burns, and surgical shock. With moderate dosages, however, skin burns play the major part in the production of disability in such men, and the most important areas of skin in this respect are the genitalia and the axillae. Owing to the delay in the full development of burns in these areas there was in this series a corresponding lag in the development of disability, the overall group disability not becoming maximal until the eleventh day after exposure. Because of this delay, the ability to formulate an early and accurate prognosis becomes of considerable importance. It has been pointed out that in unprotected men it seems to be possible to make such a prognosis from the skin condition in the first 48 hours after exposure. A rough prognosis may also be based, irrespective of the skin protection afforded, on the occurrence of nausea and vomiting within 24 hours of exposure.

The possible prevention or reduction of disability is naturally a most important aspect of the subject. With moderate dosages it seems unnecessary to protect the whole body, and the prime object of any defensive measure designed to prevent disability should be the safeguarding of the genitalia and axillae. The provision of effective protection for the genital area and the buttocks would itself reduce the incidence of disability to a very large extent. It has also been remarked that the treatment of scrotal lesions with suspensory bandages has a considerable effect in reducing the incidence of disability due to burns in this region.

Summary

An account is given of the factors responsible for the production of disability in a series of 179 men exposed to the vapour of mustard gas in the Tropics.

The influence of dosage on the type of disability produced is discussed, and the importance of protecting the genitalia is stressed.

It seems to be possible to formulate a reasonably accurate prognosis on the basis of the skin condition within 48 hours of exposure. The occurrence of nausea and vomiting within 24 hours is also of value in this respect.

The kindness of the Australian Chemical Defence Board in permitting publication of the work carried out under its direction is gratefully acknowledged. Permission to publish has also been received from the Chief Scientist, Ministry of Supply. The work of assessing and recording the disability which is the subject of this paper was done by a team, to the other members of which my grateful thanks are due, in particular to Dr. F. S. Gorrill, Dr. A. H. Ennor, and Mr. J. W. Legge.

REFERENCES

- Sinclair, D. C. (1948). *British Medical Journal*, 2, 290.
 — (1949a). *Ibid.*, 1, 476.
 — (1949b). *Brit. J. Derm. Syph.*, 61, 113.

A new unit, the Radiological Health Branch, has been formed in the United States Public Health Service to investigate and advise on risks to health due to the increased use of radioactive materials. It is under the direction of Dr. Edwin G. Williams. The unit will develop a training programme for public health workers and collect information on possible dangers to health from radioactive isotopes, x-ray apparatus, cyclotrons, beta-trons, and other atomic-particle accelerators.

POSITION OF THE APEX BEAT AS A CLINICAL SIGN

BY

D. H. ISAAC, M.B., B.S.

Late Captain, R.A.M.C.

AND

A. H. LEVY, M.B., B.S.

Late Captain, R.A.M.C.

Previous Standards

Palpation of the apex beat is one of the commonest physical signs to be elicited, yet reference to various authorities reveals considerable divergence of opinion regarding its normal limits. There is no agreement even upon the three most important criteria concerning its definition and position—namely, (1) the point of cardiac pulsation to be regarded as the apex beat; (2) whether the limit of normal should be taken in relation to an anatomical landmark; and (3) the distance from the midline to be taken as the limit of normal where direct measurement is used.

Price (1927) gives one of the most useful definitions, "as that part of the area of pulsation, distinctly palpable, which is lowest and farthest to the left." This is supported by Lewis (1946), who describes an area 1 in. (2.5 cm.) in diameter, the outermost limit of which "is the best clinical guide to the heart's size," provided that there is no displacement. Bramwell and King (1942), although regarding this as the theoretical limit of the left border of the heart, for practical purposes prefer to take the centre of the area of maximal impulse. A third view is expressed by Carlisle (1934), who states that "the apex beat is not the point of maximum impulse, but is at the point farthest down and farthest out that the beat of the left ventricle is felt."

There is similar lack of unanimity regarding the anatomical point within which the apex beat should lie. Bramwell and King, since they consider that the size of the chest is one of the chief causes of variation in position of the normal apex beat, bound it within the mid-clavicular line. Parkinson (1936), however, holds that the mid-clavicular line is difficult to apply, especially in women. The nipple line is strongly advocated in men by various authorities, Parkinson regarding it as fairly constant and an "indispensable guide"; Noble Chamberlain (1947) states that the normal cardiac thrust is "almost invariably within the nipple line." These views are supported by Lovatt Evans (1945), Price (1927), and Carlisle (1934). However, Grassmann (1921) firmly denies the value of the mammillary line in determination of the size of the heart.

A fixed anatomical line is not upheld as more than a rough guide by Campbell and Conybeare (1942), who state that the apex beat should always be given as a distance from the midline. It must be emphasized, however, that there is a wide variation in the outermost distance from the mid-sternal line which is regarded as normal. Thus Gray's *Anatomy* (1946) states that as a general rule it will be found $3\frac{1}{2}$ in. (8.9 cm.) from the midline; Noble Chamberlain gives it as up to 4 in. (10.1 cm.), while pointing out that where an abnormal apex beat is found the chest structure must be examined for scoliosis, etc., to exclude cardiac displacement. Lewis put 3–4 in. (7.6–10.1 cm.) as the usual limits of normal, and goes on to say that "a well-defined maximal impulse, the outer border of which is situated $4\frac{1}{2}$ in.

(11.4 cm.) or more to the left of the mid-sternal line . . . in an adult male is to be taken as a definite sign of the heart's enlargement in the absence of cardiac displacement." Conybeare (1942), however, includes $4\frac{1}{2}$ in. as within the normal range.

Some attention is also paid to the intercostal space in which the apex beat is found. Thus Noble Chamberlain maintains that where the impulse lies in the sixth or seventh space left ventricular enlargement is indicated. Other writers, on the contrary, regard an apex beat in the sixth space, at least, as a normal variation, especially in the aged.

"To locate accurately the apex beat is the most important step in the entire examination . . . in palpation of the heart" (Carlisle, 1934). Despite this importance it is evident, with such wide disagreement on the definition and standards of normality of a basic physical sign, that considerable confusion is likely to arise in the clinical estimation of the heart's size.

Procedure

A series of young men who presented themselves for routine Army medical examinations were examined for the position of the apex beat. These men had all been examined on at least two previous occasions and pronounced fit for military service; over 95% were in category A1, and those in a lower grade were so classified for reasons other than cardiovascular or respiratory disease.

Our aim was to determine accurately the limits within which the apex beat of a normal heart may lie, and an attempt was made to correlate this position with the structure of the thorax. Those cases in which the apex beat was over 4 in. to the left of the midline were compared with the remainder of the series.

The apex beat was defined as follows: (1) The point farthest out and lowest down at which the finger was distinctly raised by the cardiac impulse. (2) In those cases in which the finger was not raised and/or in which there was much diffusion of the impulse, that point was taken farthest out and down at which a distinctly localized maximal impulse was felt. The apex beat was regarded as impalpable and not measured in those cases in which it was too diffuse and no localized maximal pulsation was found.

The cardiovascular and respiratory systems were examined clinically and were in all cases normal. X-ray examination was not employed as a routine.

In assessing the structure of the chest, the girth was measured in mid-expiration at the level of the nipples. Any scoliosis and/or kyphosis were noted; as an additional aid in detecting spinal curvature which was not obvious from inspection of the vertebral spines a comparison was made of the height of the shoulders.

Results

The present series consists of 500 unselected subjects: each of us, independently, examined 250 cases. The age distribution of the men is shown in Table I. The average age was 21.5 years and the maximal variation was from

TABLE I.—Age Distribution of Total Series

Age	No.	%	Age	No.	%
17	1	0.2	30	2	0.4
18	8	1.6	31	0	0
19	89	17.8	32	2	0.4
20	213	42.6	33	5	1.0
21	59	11.8	34	1	0.2
22	26	5.2	35	3	0.6
23	20	4.0	36	2	0.4
24	20	4.0	37	3	0.6
25	9	1.8	38	0	0
26	19	3.8	39	4	0.8
27	3	0.6	40	2	0.4
28	6	1.2	41	0	0
29	1	0.2	42	2	0.4

17 to 42 years. In 105 (21%) of the 500 cases the apex beat was not felt or was not sufficiently localized according to the definition. Of the 395 palpable cases the apex beat was found to be over 4 in. from the midline in 45 (11.4% of the total palpable). The ages of the 45 are shown in Table II.

TABLE II.—Age Distribution of Those with Apex Beat Over 4 in. from Midline

Age	No.	%	Age	No.	%
19	8	17.8	23	1	2.2
20	19	42.2	24	4	8.9
21	5	11.1	25	2	4.4
22	5	11.1	26	1	2.2

The apex beat in each subject was measured to the nearest $\frac{1}{4}$ in. (0.6 cm.) from the midline, with the patient recumbent. It was found that the variation was from 2 in. (5 cm.) (1 case) to $5\frac{1}{2}$ in. (14 cm.) (1 case) to the left of the midline, the maximal incidence being $3\frac{1}{2}$ in. (8.9 cm.) (134 cases). The accompanying Graph gives the distribution of all the apex beats which were palpable.

Graph showing distribution of all apex beats that were palpable.

The average girth of the chest in the 500 cases was 34.6 in. (87.8 cm.); and in those in whom the apex beat was more than 4 in. from the midline it was 35.1 in. (89.1 cm.). Scoliosis as visible in the vertebral spines was found in 113 (22.6%) of the total series; and in 12 (26.6%) of the over-4-in. group. One or other shoulder was found to be significantly higher in 130 (26%) of the total, as compared with 16 (35.6%) of those with an apex beat over 4 in. These findings are shown in Table III.

TABLE III

	Total Series	Apex Beat Over 4 in. from Midline
No. of cases	500	45
Scoliosis	113 (22.6%)	12 (26.6%)
Scoliosis to left in thoracic region	67 (13.4%)	10 (22.2%)
Scoliosis to right in thoracic region	30 (6.0%)	2 (4.4%)
Shoulder higher	130 (26.0%)	16 (35.6%)
Average chest girth	34.6 in.	35.1 in.

The apex beat was found almost equally in the fourth and fifth intercostal spaces, and only in a very few in the third (2) and sixth (10) intercostal spaces. Occasional radiographs were taken in some of the cases above 4 in., and in all cases the apex beat as found clinically corresponded with the apex of the heart radiologically, and in no case was any cardiac enlargement detected.

Discussion

Examination of the results obtained shows that in no less than 11.4% of palpable cases the apex beat was over 4 in. to the left of the midline. According to previous standards many of these cases might have been regarded as due to cardiac enlargement. This is, however, extremely improbable in view of the subjects examined, who were all healthy young soldiers showing no clinical evidence of heart disease. Hartley (1948) has shown that in a large series of recruits examined radiologically 1% manifested apparent cardiac enlargement; out of 18 such cases followed up for some years, only eight had pathological enlargement.

Our impression was that anatomical landmarks, particularly the mid-clavicular line, were not easy to define and apply. From Table III it will be seen that there was little difference (less than 2%) in the average girth of the total series and of those with apex beat over 4 in. from the midline. Thus there would seem to be little to bear out the assumption of a direct relationship between the girth of the chest and the position of the apex beat.

Price (1927), Parkinson (1936), Lewis (1946), and Evans (1948) state that displacement of the heart may be caused by scoliosis. At first sight (Table III) this would not seem obvious; when, however, the direction of the thoracic scoliosis is analysed a marked preponderance of scoliosis to the left, as compared with that to the right, is apparent in those cases over 4 in. The ratio of thoracic scoliosis to the left compared with the right was 2:1 in the total series, and 5:1 in those with an apex beat over 4 in. It would therefore seem that scoliosis to the left in the thoracic region is a cause of cardiac displacement in a number of cases. Some, however, had no scoliosis, and in these another cause must be sought. Bramwell (1932), King, and Parkinson regard the height of the diaphragm as important in this respect; but this point could not be investigated in a purely clinical series. Foti (1937) points out that in a narrow thorax the apex beat may be carried to the left by the lateral thrust of the heart. Sternal depression has been emphasized by Evans (1946) as a cause of displacement.

In recent years much prominence has been given to the correct assessment of patients with abnormal cardiac signs. The "innocent" murmurs have been classified (Evans, 1947) and their recognition has prevented much unwarranted cardiac invalidism. R. K. Price (1949) showed that of 200 children referred to a cardiac clinic no fewer than 39 normal children had been restricted in their activities. In a similar way the finding of a displaced apex beat on medical examination may lead to the diagnosis of cardiac enlargement, with perhaps unnecessary invalidism and the production of a cardiac neurosis. Thus the finding of an apex beat outside the range previously accepted as normal cannot be regarded as pathological unless confirmed by other signs of heart disease or by cardioscopy. The cause may be sought in the structure of the chest, particularly with regard to scoliosis, or in the relation of the heart to the diaphragm. The recognition of this was emphasized by Sir John Parkinson, who said: "In the past it has often happened that this deceptive apex beat has been attributed to real enlargement indicating a cardiac lesion, and the stigma has followed a healthy subject, perhaps for a lifetime."

Summary

A series of 500 cases examined for the position of the apex beat is presented.

The normal distribution of the position of the apex beat is shown.

The apex beat was found to be more than 4 in. from the midline in 11.4% of those palpable.

Thoracic scoliosis to the left was considered to be a cause in some cases.

The necessity for differentiation between displacement due to structural causes and true cardiac enlargement is pointed out.

We wish to thank Colonel G. Moulson, A.D.M.S., Salisbury Plain District, for permission to publish this paper.

REFERENCES

- Bramwell, C. (1932). *Heart Disease*, p. 116. London.
— and King, J. T. (1942). *Principles and Practice of Cardiology*, p. 23. London.
Carlisle, G. L. (1934). *Practical Talks on Heart Disease*, p. 14. London.

- Chamberlain, E. Noble (1947). *Symptoms and Signs in Clinical Medicine*, p. 115. Bristol.
Conybeare, J. (1942). *Textbook of Medicine*, p. 494.
Evans, C. Lovatt (1945). *Starling's Physiology*, p. 535. London.
Evans, W. (1946). *Brit. Heart J.*, **8**, 162.
— (1947). *Ibid.*, **9**, 1.
— (1948). *Cardiology*, p. 42. London.
Foti, A. (1937). *Med. Rec., N.Y.*, **146**, 438.
Grassmann, K. (1921). *Münch. med. Wschr.*, **68**, 1420.
Gray's *Anatomy* (1946). 29th ed., p. 1520. London.
Hartley, R. (1948). *British Medical Journal*, **2**, 673.
Lewis, T. (1946). *Diseases of the Heart*, p. 118. London.
Parkinson, J. (1936). Lumleian Lecture. London.
Price, F. W. (1927). *Diseases of the Heart*, p. 2. London.
Price, R. K. (1949). *British Medical Journal*, **1**, 515.

A SIMPLIFIED METHOD OF BLOOD-SUGAR ESTIMATION*

BY

D. A. HALL, B.Sc., Ph.D.

Assistant Lecturer in Medical Biochemistry, Department of Medicine, University of Leeds

The majority of the standard methods for blood-sugar determination are neither simple enough nor rapid enough for general use outside a laboratory. Furthermore, many of the existing methods measure, in addition to glucose, other reducing components of the blood. In recent years new or modified tests have been devised in an attempt to provide the general practitioner with a reliable method for use at the bedside, or at least in the normally equipped surgery. With the latter end in view some of the recent modifications have been examined and a modification of one of them (Haslewood and Strookman, 1937) is proposed.

Recent Modifications

Hagedorn, Halstrom, and Jensen (1935).—This simplification of the earlier method of Hagedorn and Jensen uses a shorter period of heating, carried out over a naked flame, thus decreasing the amount of equipment required. An accuracy to within 3% of the older method is claimed. The combination of zinc hydroxide precipitation for protein removal with ferricyanide oxidation of the glucose gives results which approximate closely to true sugar values. The final operation, however, consists in the titration of liberated iodine, a procedure not readily performed outside the laboratory. Further, as the method is based on the estimation of surplus oxidizing agent, it is necessary for a blank determination to be carried out with each estimation—an unfortunate complication to add to any method designed for rapid use.

Lauber and Mattice (1944).—These authors describe an adaptation of the original method of Folin and Wu (1920), employing zinc hydroxide to precipitate the protein, but the precipitate has to be removed by centrifuging, necessitating the use of apparatus available only in the laboratory. Further, owing to the fact that some of the red cells may be laked before precipitation is completed, the values obtained are greater than the true glucose value.

Leech and Woodford (1948).—The use in this method of an oxidizing agent other than the conventional alkaline copper or ferricyanide solution permits a far simpler procedure to be planned. The reduction product of the oxidizing agent (3:5 dinitrosalicylic acid) has a sufficiently strong colour to enable it to be compared visually with

*A paper the substance of which was read in the Section of Medicine at the Annual Meeting of the British Medical Association, Harrogate, 1949.