delivery, compared with 26% after a delivery under general anaesthesia.

Post-operative Complications

Headache was the only early complication noted; 19 (3.8%) women complained of a moderate or severe degree of headache. In Table II the incidence of headache after spontaneous deliveries is compared with that following deliveries under spinal analgesia.

TABLE II.—Incidence of Headache

	Mild	Moderate	Severe
Deliveries under spinal analgesia	10.5% .. 2.2% .. 1.6%		
Spontaneous deliveries	9.7% .. 1% .. 0.3%		

There were no sequelae affecting the chest. No remote complications of spinal analgesia were brought to our notice, although a large percentage of the patients attend the post-natal clinics and many have been seen during a subsequent pregnancy.

Over the same period 150 caesarean sections have been carried out under heavy cinchocaine analgesia, with no maternal mortality. In the six-year period 1943–9—i.e., including Resnick's series—there have been 287 consecutive caesarean sections under spinal analgesia without a maternal death.

It is important to stress the fact that the safety of the method depends on using the minimal dosage required to produce adequate analgesia, and that for a vaginal delivery it is rarely necessary to use more than 0.8 ml. of heavy cinchocaine. In these quantities the method is very safe.

Discussion

None of the theoretical evils of spinal analgesia has been encountered. The advantages of this form of analgesia to the mother, the baby, and the obstetrician have been adequately summarized by Thomas, Resnick, Anderson, and others. Their conclusions have been confirmed in this series. One cannot fail to be impressed by the well-being of the mother and the absence of shock after a difficult vaginal delivery under spinal block.

I was brought up in the "anti-spinal school" and was thus well primed in all the weighty arguments which are levelled against its use in obstetric practice. I then found myself in the dilemma of working in a department where spinal analgesia was an established practice and where the results had been extremely satisfactory. It was not without qualms that the decision was made to follow the established tradition rather than upset a machine which was running smoothly.

Summary

A series of 511 operative vaginal deliveries under heavy cinchocaine spinal analgesia is presented and analysed. The technique is simple and safe in the dosage employed. The analgesia has been excellent and no ill-effects to the child have been noted. The only factor unfavourable to the mother was headache.

My thanks are due to Mr. George Brown, medical superintendent of this hospital, for permission to publish the results of these cases. I am especially grateful to my predecessors, and to the resident obstetrical officers (Drs. Donovan and Ridehalgh) who have performed a large percentage of the operative deliveries, and to the midwifery staff for their excellent co-operation and high standard of nursing care.

REFERENCES

- Anderson, A. F. (1946). *J. Obstet. Gynaec. Brit. Emp.*, **53**, 347.
 Andros, G. J., Dieckmann, W. J., Ouda, P., Priddle, H. D., Smither, R. C., and Bryan, W. M. (1948). *Amer. J. Obstet. Gynec.*, **55**, 806.
 Cosgrove, S. A. (1927). *Ibid.*, **14**, 751.
 Hugill, J. T., and Gillespie, C. (1948). *Canad. med. Ass. J.*, **58**, 146.
 Resnick, L. (1945). *British Medical Journal*, **2**, 722.
 Thomas, R. C. (1942). *J. Obstet. Gynaec. Brit. Emp.*, **49**, 247.
 — (1944). *Ibid.*, **51**, 324.

CAUDAL ANALGESIA

BY

T. C. THORNE, M.C., M.R.C.S., L.R.C.P., D.A.

Anaesthetic Registrar, Rochford General Hospital;
 Clinical Assistant, Southend General Hospital

Professor Hingson's recent visit to this country and his lectures and publication (Hingson, 1949) on continuous caudal analgesia will undoubtedly give a fresh stimulus to the interest aroused on this subject by his earlier publications.

It is unlikely that continuous caudal analgesia will be used extensively in this country in the immediate future for the relief of pain in normal labour and delivery, because of the lack of trained personnel to induce, supervise, and maintain analgesia, though Galley (1949) has published a series of such cases. Nevertheless, it is clear that the continuous method can play an extremely useful part in the treatment of angiospastic states, and, in the words of a recent *B.M.J.* Editorial (1949), "can confer benefits to mother and baby which they should not be denied, and it is to be hoped that obstetricians in this country will not be satisfied until they can obtain it at least for those mothers for whom it is specially indicated."

The attention which has now been focused on continuous caudal analgesia should not be allowed to push further into the background the single-injection technique of producing caudal analgesia in surgery and obstetrics, a method which has been neglected in this country. Even in the U.S.A. there seems to be a body of opinion which prefers the single-injection technique (which has been widely used in surgery) rather later in labour, employing long-acting drugs and repeating the needle insertion and injection if necessary (Cullen, 1946; Brown, 1948; Crawford and Chester, 1949) because of the possible dangers of the continuous technique using a needle and the increased difficulty of insertion using a catheter. Moreover, in the average hospital in this country it is unlikely that enough continuous caudal blocks will be required to allow many to become really proficient in the art of inserting a needle into the caudal canal.

It is my object to draw attention to the usefulness of the single-injection technique in inducing analgesia in surgery and obstetrics and at the same time in providing training in needle insertion, to describe some cases in which continuous caudal analgesia has been used, and to mention some of the difficulties and complications associated with the insertion of a needle into the caudal canal.

Single-injection Caudal Analgesia

Up to 1943 very little had been published on the subject in this country. Wilson (1931), Poole (1941), and Anderson (1949) reported on its use in labour; Mitchell (1933) recommended it in prostatectomy; and Hewer (1943) described its use but preferred low spinal analgesia. Molesworth (1946), Lee (1947), Goldman (1948), Minnitt and Gillies (1948), and Dawkins (1949) describe and generally recommend it, mainly in vaginal and rectal operations and for forceps delivery. Hewer (1948) still prefers low spinal analgesia. Corlette (1948) regards it as a dangerous method apt to produce shock and with a high failure rate. At the 1948 Annual Meeting of the B.M.A. Steel (1948) recommended its use, but in the discussion which followed several speakers objected to it because of the risk of collapse and because of the unpredictability of analgesia. Kenny (1947) introduced "proctocaine" by the caudal

route for the treatment of pain due to inoperable pelvic carcinoma.

In my series it has been used in the following cases:

	Successes	Failures
Prostatectomy and cystoscopy	33	5
Haemorrhoidectomy	26	1
Vaginal repairs	109	14
Forceps delivery	10	0
Herniotomy	3	0
Relief of pain	2	0
	183	20

In addition there have been three cases in which the sacro-coccygeal membrane could not be located with the needle and two cases which were rejected at pre-operative examination because of inability to palpate the sacral hiatus.

Prostatectomy.—In this operation it has been used to reduce stimulation from digital enucleation under moderately deep general anaesthesia and to secure haemostasis (Rowbotham, 1947), though for the latter purpose with no noticeable success, possibly because the block extended above the sacral segments with the dose used—20 ml.

Haemorrhoidectomy, Including All Anal Operations.—Caudal block is considered to be the ideal method for these cases, as doses of 15 to 20 ml. provide complete relaxation and analgesia to be equalled only by the low spinal method with its risk of headache.

Vaginal Repair, Including Vaginal Hysterectomy.—The main advantage is reduction in haemorrhage. For Fothergill's repair and vaginal hysterectomy 35 to 40 ml. is required.

Forceps Delivery.—For the handicapped mother or foetus low spinal analgesia has been used with good results in over 40 cases, but with severe post-operative headache in seven. Caudal block is now used in preference to spinal analgesia when time is not a factor; dose—35 to 40 ml.

Herniotomy.—Excellent operating conditions are provided without complete muscular flaccidity by doses of 40 ml. with analgesia at about T 10.

Pain Relief.—A 43-year-old woman with fourth-stage carcinoma of the cervix required morphine for intractable pain in the legs. Proctocaine, 35 ml., gave immediate relief. There was paralysis of sphincters for 24 hours. On discharge three weeks later she was still pain-free and ambulant. She died three weeks after discharge.

Continuous Caudal Analgesia

1. With Hingson's caudal needle (4 cases)

Case 1.—A 26-year-old primipara at the 37th week of pregnancy, with eclampsia (one fit), had induction performed after 36 hours; 12 hours later she was in labour. As she had oliguria—36 oz. (1 litre) in 48 hours—and was very restless though heavily sedated, caudal block was established, being maintained for 6½ hours; delivery was by low forceps and episiotomy. The baby weighed 4 lb. 2 oz. (1.87 kg.). During block, 9 oz. (256 ml.) of urine was passed, and diuresis continued.

Case 2.—A 23-year-old primipara at term, in labour with a large vulval haematoma increasing in size, painful, and extremely tender. Two injections were given before delivery by low forceps and episiotomy.

Case 3.—A 19-year-old primipara, at the 32nd week of pregnancy, had bilateral ovarian cystectomy performed, and three days later went into premature labour. Towards the end of the first stage she felt as if the abdomen were bursting, and became very distressed. Two injections were given before delivery, by low forceps and episiotomy, of a 4½-lb. (2-kg.) baby.

Case 4.—A 22-year-old primipara, at term with very flat pelvis, received six injections for forceps delivery, which pro-

vided good analgesia for 12 hours; delivery was by high forceps and episiotomy.

The babies in the above cases cried immediately, and the third stages were normal, with minimal bleeding.

2. With polythene tubing (size 1) passed through a 15-gauge Barker spinal needle (4 cases)

Case A.—Thrombophlebitis with typical white leg occurred in a woman of 29 three weeks after caesarean section. The right leg was very painful and tender, with 3-in. (7.6-cm.) swelling; temperature 101° F. (38.3° C.). Caudal analgesia provided complete relief from pain, so that she was able to move the leg and carry out exercises. The temperature fell rapidly. In 3½ days she received 14 injections to produce intermittent analgesia as recommended by Lull and Hingson (1948). The bed was fixed in the 15-degree Trendelenburg position and she was seen by the physiotherapist. The polythene tube was removed eight days after insertion; she was then allowed up. The small skin wound healed cleanly. On discharge 23 days after admission there was a ¼-in. (1.9-cm.) swelling and slight ache on prolonged standing. Three months later there was no oedema, aching, or disability, and the swelling was reduced to ½ in. (1.27 cm.).

Case B.—A 25-year-old primipara, with severe pre-eclamptic toxæmia and a very small baby. Two hours after caudal block was instituted it was found that the polythene tube had broken at the skin surface. The proximal piece was recovered with some difficulty, after incision under local analgesia. The tube was acutely angulated as it left the skin, and this, combined with the supine position, was the probable cause of rupture. Fourteen days later, the wound having healed by first intention, the block was re-established and maintained for 3½ days, though after the first day without much effect on the blood pressure; it was therefore abandoned and labour induced, with delivery of a 20-oz. (567-g.) baby, which lived a few hours. As the result of experience with this case, it is believed that to maintain a reduction of blood pressure the tube should be inserted well inside the canal, as high as the lumbar region if possible.

In addition, polythene tubing has been used in two prostatectomies without incident.

Technique and Complications

1. Invariably it has been found that when the needle is correctly placed injection is extremely easy, and that analgesia of S 4 and 5 is present within five minutes.

2. Lumbar puncture and Hingson's caudal needles have been most often employed, but in thin patients No. 1 intravenous or No. 17 hypodermic needle is quite satisfactory.

3. When boiled, polythene tubing tends to swell and may no longer pass through a 15-gauge needle; therefore the alternative method of sterilization by soaking in 1% "cetavlon" has been used.

4. After the first 30 cases the test dose was not employed except in those requiring continuous caudal analgesia. Dural tap has occurred once, and was immediately obvious.

5. Bloody tap has occurred 32 times; after moving the needle, aspiration is usually negative and injection can then be made safely.

6. Toxic reactions occurred in six cases, including convulsions in three, all in elderly patients; some of these reactions may have been partially intravenous, but it must be remembered that symptoms may be produced by too rapid injection, particularly in the elderly. Except in the eclampsia cases the blood pressure has not fallen more than 25 mm. in the whole series.

7. In five of the 20 failures injections were not completed, because of bloody or dural tap. Of the 15 cases in which injections were completed, in three the injections were unilateral and in four probably partially intravenous;

the remaining eight occurred early and were probably dorsal to the sacrum.

Summary

A series of 203 cases of single-injection analgesia and 8 cases of continuous caudal analgesia have been described. The single-injection technique is shown to be an extremely useful method of producing analgesia in vaginal repairs and forceps delivery, and an ideal one for anal operations. In addition it provides training in needle insertion. Some of the difficulties and complications are mentioned.

I wish to thank Dr. J. A. Lee for interesting me in caudal analgesia and for continuous help and encouragement; Miss E. M. Whapham, Mrs. F. Bridge, and Mr. B. J. Sanger for their co-operation and permission to publish case reports; and the theatre and labour-ward staffs for their assistance.

REFERENCES

- Anderson, A. F. (1949). *British Medical Journal*, **1**, 233.
British Medical Journal (1949). **2**, 801.
 Brown, W. E. (1948). *Amer. J. Obstet. Gynec.*, **55**, 717.
 Corlette, C. E. (1948). *A Surgeon's Guide to Local Anaesthesia*. Bristol.
 Crawford, O. B., and Chester, R. V. (1949). *Anesthesiology*, **10**, 473.
 Cullen, S. C. (1946). *Anesthesia in General Practice*. Chicago.
 Dawkins, C. J. M. (1949). In *Modern Practice in Anaesthesia*, edited by Frankis T. Evans. London.
 Galley, A. H. (1949). *Anaesthesia*, **4**, 154.
 Goldman, V. (1948). *Aids to Anaesthesia*, 2nd ed. London.
 Hewer, C. L. (1943). *Recent Advances in Anaesthesia*, 4th ed. London.
 — (1948). *Ibid.*, 6th ed. London.
 Hingson, R. A. (1949). *British Medical Journal*, **2**, 777.
 Kenny, M. (1947). *Ibid.*, **2**, 862.
 Lee, J. A. (1947). *A Synopsis of Anaesthesia*. Bristol.
 Lull, C. B., and Hingson, R. A. (1948). *Control of Pain in Child-birth*. Philadelphia.
 Minnitt, R. J., and Gillies, J. (1948). *Textbook of Anaesthetics*, 7th ed. Edinburgh.
 Mitchell, W. E. M. (1933). *Lancet*, **2**, 348.
 Molesworth, H. W. L. (1946). *Regional Anaesthesia*, 2nd ed. London.
 Poole, W. H. (1941). *J. Obstet. Gynaec. Brit. Emp.*, **48**, 84.
 Rowbotham, S. (1947). *Anaesthesia*, **2**, 70.
 Steel, G. C. (1948). *British Medical Journal*, **2**, 92.
 Wilson, J. St. G. (1931). *Lpool med.-chir. J.*, n.s. **5**, 138.

All patients were subjected at different times to various methods of control. Generally speaking, alkaloid treatment, if still in progress, was alternated with diparcol, dummy tablets containing lactose and also capsules of "benadryl" being substituted for the latter. Three patients (two post-encephalitics and one senile vascular case) showed significant improvement. In one of these it was dramatic. From being absolutely helpless and bedridden the patient was at the end of two months able to walk unaided, dress herself, knit, and generally help the nursing staff in ward duties. She was discharged home after maintaining this sensational improvement for several months. In another case the improvement was pronounced. The third patient, who was previously rigid and helplessly bedridden, was rendered free to move about in bed and feed herself.

Experience with these cases revealed that, whereas the subjective and objective signs of illness were not appreciably altered, the degree of motor control regained was such that the patients were enabled to lead a purposeful life—knitting, sewing, or doing useful domestic work. This improvement has been maintained.

Six patients showed no significant response to diparcol treatment, but one of these was found post mortem to have a cerebral glioma, so he could justifiably be excluded from the series.

No blood dyscrasia or other serious complication was observed during treatment. The only side-effect of diparcol requiring comment is the mental depression which occurs in some patients; amphetamine should be given in an attempt to control this symptom.

In general, the results of this investigation revealed that in certain instances diparcol may appreciably benefit sufferers from Parkinson's syndrome. It seems likely that only an adequate therapeutic test will reveal the type of case likely to show the greatest degree of response.

I wish to express my thanks to Dr. W. R. Thrower, consulting physician to May and Baker, Ltd., for advice and assistance during the investigation and for the supply of diparcol tablets for clinical trial.

R. M. GRAY, M.B., B.S.,
 Medical Officer, British Railways,
 Newcastle-upon-Tyne.

REFERENCE

- Sigwald, J., et al. (1947). *Rev. Neurol.*, **79**, 683.

Medical Memoranda

Experiences with "Diparcol"

Many methods have from time to time been introduced for treating Parkinson's syndrome and later discarded after apparent early successes. The evaluation of any remedy under controlled conditions is complicated by the varied aetiological factors and clinical features of the sufferers. Nevertheless any new therapeutic measure which is basically sound justifies critical examination. The claims made by Sigwald *et al.* (1947), using diethazine hydrochloride, or "diparcol," were such that a confirmatory investigation was made for a period of six months during 1948 at the Grove Road Hospital, Richmond, Surrey.

CLINICAL INVESTIGATION

Nine unselected patients with Parkinson's syndrome were treated with diparcol. Four cases were probably post-encephalitic; three were of the senile vascular type; one was of uncertain aetiology, possibly the result of head injury; and one was ultimately found to be suffering from a small glioma in the right putamen. All had been under continuous observation for several years and had been given solanaceous alkaloids, with no more than temporary benefit. The average daily dose of diparcol, slowly attained, was 1 g. in three equal portions, the maximum daily dose being 1.5 g.

Profuse Haemorrhage from Ruptured Corpus Luteum Simulating an Ectopic Pregnancy

Rupture of a corpus luteum or Graafian follicle may occasionally give rise to symptoms suggestive of an extra-uterine gestation or appendicitis. For this reason it is well known that at operation for ectopic pregnancy blood may be found in the pelvis although no ectopic pregnancy is discovered. What is, perhaps, less well known is the extent of the bleeding which may follow such follicular rupture.

Weil (1939) reported 30 cases in which bleeding was found at operation to be due to rupture of a corpus luteum or Graafian follicle, and five others in which the diagnosis was made and conservative treatment carried out. In 11 cases the amount of blood found was about a pint (570 ml.). Weil states that any amount from 10 ml. to a litre may be discovered. This is illustrated in the case here described, in which 2 pints (1.1 litres) of blood were removed from the abdomen at operation.