Twenty-five active and 4 inactive cases of pulmonary tuberculosis were disclosed.

Twenty-seven further cases of disease were revealed. They included tuberculous effusion (6), bronchiectasis (11), new growth (4), lung abscess (1), spontaneous pneumothorax (1), cardiac (3), and cough fracture (1).

It has been argued that radiography should be widely applied to the symptom-complex of "pneumonia."

REFERENCES

- Baum, O. S., and Amberson, J. B. (1942). *Amer. Rev. Tuberc.*, **45**, 243.
 — and Baum, L. F. (1949). *Ibid.*, **59**, 68.
 Cohen, R. C. (1949). *British Medical Journal*, **1**, 133.
 Davies, D. T., Graham-Hodgson, H., and Whitby, L. E. H. (1935). *Lancet*, **1**, 191.
 Douglas, A., Pinner, M., and Wolepor, B. (1929). *Amer. Rev. Tuberc.*, **19**, 153.
 Eberle, A. J. F. (1949). *Tubercle*, **30**, 146.
 Morgan, T. N., and Wylie-Smith, R. (1943). *Lancet*, **2**, 731.
 Nairn, R. C. (1949). *Thorax*, **4**, 101.
 Paulley, J. W., Lees, D. H., and Pearson, A. C. (1949). *British Medical Journal*, **1**, 135.
 Rist, E. (1929). *Canad. med. Ass. J.*, **21**, 143.
 — and Ameuille, P. (1922). *Paris méd.*, **33**, 14.
 Toussaint, C. H. C., and Pritchard, E. K. (1944). *Post.-Grad. med. J.*, **20**, 143.
 Williams, L. R., and Hill, A. M. (1930). *New Engl. J. Med.*, **203**, 418.

ADRENAL CORTICAL TUMOURS OF UNUSUALLY LARGE SIZE

BY

GEORGE LUMB, M.D.

Honorary Consultant, Westminster Hospital; Senior Lecturer in Pathology, Westminster Medical School

Tumours arising from the adrenal glands are rare, especially those primarily in the cortex. It is well known that cortical hyperplasia of either a focal or a generalized type occurs during middle life and is often associated with feminine pseudo-hermaphroditism (Glynn, 1912). Where the hypertrophy is generalized, a gland may enlarge and even approximate in size to the kidney. All gradations occur between simple hypertrophy and true adenoma formation, and as the two conditions appear during the same age period it is sometimes very difficult to differentiate one process from the other.

Primary malignant tumours arising from the glandular elements of the cortex are usually referred to as adenocarcinomata—a term which has largely replaced that of hypernephroma. As it was common to misapply the older term to renal tumours and non-cortical adrenal tumours it is obviously an advantage to abandon it, but since the adrenal cortex is developed from mesoderm it is clearly not entirely accurate to employ a title which indicates an epithelial origin. However, most pathologists agree that the histological appearances are of "adenocarcinomatous" type, and I shall use the term here.

In the same way as difficulty is encountered in drawing a sharp line between simple hyperplasia and true adenoma formation, so it is not always easy to differentiate histologically between benign and malignant conditions. Thus when metastases can be demonstrated the diagnosis is clear despite the fact that the primary tumour may be made up of cells apparently identical with those of a normal adrenal gland. Often, however, spread occurs late, and large tumours are encountered which undergo necrosis and extensive haemorrhage yet possess thick well-defined capsules.

Adrenal cortical tumours may be further subdivided into those which are associated with hormonal disturbances and those which are not. Examples of cases occurring without hormonal disturbances are well recognized but rare. The majority occur in adults over the age of 30 and have been described in both sexes. Cahill *et al.* (1942) found five such cases in a series of 13 adrenal cortical tumours. The histological appearances vary from those that are well differentiated to those showing an extreme degree of anaplasia.

The following two cases fall into the group of tumours of adrenal cortical origin without hormonal disturbances. In each case the clinical picture was that of a large abdominal mass causing minimal discomfort. In both cases the tumours were discovered on routine examination, when the patient had sought advice for other symptoms. Both were easily removable at operation.

Case 1

A man aged 69, under the care of Sir Stanford Cade, was first seen at Westminster Hospital on May 28, 1949, complaining of difficulty in micturition. Five months previously he had noticed a painless swelling in the right side of the abdomen.

Examination revealed a heavily built man with dyspnoea on exertion and a blood pressure of 240/120, but without clinical evidence of Cushing's syndrome. Prostatic enlargement was thought to account for his urinary symptoms. A very large, tense, immobile mass was present in the right side of his abdomen extending up to the xiphisternum and across the midline towards the left. The possibility of a large hydro-nephrosis was excluded because the excretion pyelograms showed normal kidney patterns.

At laparotomy on June 7 a large retroperitoneal tumour lying anterior to and above the right kidney was discovered. No adrenal gland could be found on that side. The tumour was encapsulated and stripped off the kidney with ease, leaving it normal, with no suggestion of its having any association with the mass. After operation the blood pressure fell to 120/80, and the patient seemed weak and collapsed. He developed a basal bronchopneumonia and died on June 13, six days after operation. Post-mortem examination was not allowed.

Case 2

A woman aged 57, under the care of Mr. G. T. Mullally, was first seen at Westminster Hospital on August 2, 1946, complaining of a painless swelling in the right upper abdomen which had been present for 3½ years. She had no symptoms apart from occasional abdominal colicky pains and a mild degree of constipation.

Examination showed that she was well nourished and of slight build, with a blood pressure of 130/80, and no clinical suggestion of any hormonal abnormality or of Cushing's syndrome. A hard rounded mass was palpable in the right upper quadrant of the abdomen. It was not attached to the skin, and moved down on respiration. A barium meal revealed displacement but no abnormality of the bowel. Excretion pyelograms showed normal renal shadows, and the blood urea was 34 mg. per 100 ml.

Laparotomy, performed by Mr. Mullally on August 18, revealed a large encapsulated retroperitoneal tumour lying slightly behind and to the right of the right kidney, but in no way associated with it. No adrenal gland could be found on the right side. The tumour was easily removable, but after operation the patient was collapsed, with depressed blood pressure of 80/40. Plasma and blood transfusion produced no improvement. On the fourth day a basal bronchopneumonia developed, and the patient died on the sixth day after operation. Post-mortem examination was not allowed.

Morbidity and Histology

The morbid anatomical and histological appearances of the tumours were so similar that they may be described together.

In each case roughly spherical completely encapsulated tumours were found, with the following dimensions and weight :

	Diameter	Weight
Case 1 ..	22.5 cm. (8½ in.)	3,750 g. (8¼ lb.) (see Fig. 1)
Case 2 ..	23.5 cm. (9¼ in.)	5,350 g. (11½ lb.) (see Fig. 2)

with lipid globules. Fuchsinophil staining by Broster and Vines's (1933) method gave for the most part negative results, although occasional granules scattered through the cells of the tumours took up the red stain feebly.

A marked feature of the tumours was their extreme vascularity, and large numbers of small vessels had cells of cortical

FIG. 1.—The cut surface of the tumour in Case 1.

There was no evidence of a pedicle or of any association with the kidney, and in neither case was an adrenal gland found on the same side as the tumour. The capsules were quite thin but extremely tough. The cut surface of the tumour showed considerable haemorrhage and necrosis, so that most of the substance consisted of pinky-yellow friable material, leaving only small areas of yellowish fleshy tumour substance lying just beneath the capsule. Spaces containing blood clot were a noticeable feature of both cases.

Microscopically the central part of the tumours was found to be composed mostly of necrotic debris with occasional groups of cells remaining. Only in those areas lying immediately under the capsule was it possible to determine the histological pattern. In each case the picture was similar, and consisted of groups of polyhedral or round cells with deeply

FIG. 3.—Typical microscopic picture at periphery of tumour (Case 1). (× 65.)

suprarenal type either closely attached to or actually forming their walls (see Fig. 4). Many areas of both tumours showed evidence of thrombosis in vessels.

In the necrotic areas there were small granular zones of calcification, not large enough to cause noticeable "grittiness" on cutting, but demonstrable microscopically by special staining methods. This feature is considered to be evidence that the

FIG. 4.—Photomicrograph showing close association of tumour cells with a small vessel (Case 2). (× 65.)

FIG. 2.—The cut surface of the tumour in Case 2.

staining clear-cut nuclei and foamy pale-staining cytoplasm, bounded by a well-marked cell membrane. These cells were arranged principally in columns, but often enclosed spaces. The general appearances were similar to those found in normal adrenal cortex (see Fig. 3).

Frozen sections of the tissue stained by methods to demonstrate fat showed the cytoplasm of the cells to be packed

breakdown of the tumour was not a very recent occurrence. The capsule was thin and in each case made up of fibrous tissue with intermingled smooth-muscle fibres.

Careful search of both tumours failed to reveal any of the cytological features usually associated with malignancy, such as giant-cell forms or hyperchromatism of the nuclei.

Discussion

These cases present interesting features both clinically and pathologically. Laparotomy was essential in order to make a diagnosis, and easily removable tumours were found, but in both cases blood pressure fell sharply afterwards, suggestive of suprarenal shock, and it seems probable that this feature was in part, if not entirely, the cause of death. It is therefore important to consider the possibility of a cortical suprarenal neoplasm when a very large encapsulated retroperitoneal tumour is found which is causing no symptoms referable to the organ of origin; for it seems likely that if the diagnosis were made early enough hormonal replacement therapy might prevent post-operative collapse. It is therefore particularly valuable to note that the naked-eye appearances in my two cases and in the case described by Anderson *et al.* (1943) were exactly similar.

Walters and Kepler (1938) pointed out that probably the most important single feature in the surgical treatment of cortical adrenal tumours was the control of post-operative shock. Cecil (1933) stated that 39% of patients operated on died of shock. It may be added in this connexion that Cahill *et al.* considered that post-operative shock occurred only in those cases in which, as a result of the tumour, a syndrome of Cushing's type was present and hormonal excess was affecting metabolic functions such as salt-and-water balance and capillary permeability. They further state that, in their series, cases without clinical hormonal upsets were never associated with post-operative collapse. This view cannot be confirmed from the present cases.

From the pathological point of view the gross and the microscopical structure of these tumours provokes discussion whether they are benign or malignant. Features in favour of their benignity are the presence of the well-formed capsule and the absence of evidence of metastases, although it must be admitted that it was not possible to perform necropsies. The absence of demonstrable mitoses and the close resemblance of the tumour cells to those of the normal adrenal seem to point to a non-malignant origin, although it may be repeated that numbers of metastasizing tumours with complete differentiation of cell type have been described.

It is somehow difficult to reconcile such very large masses showing so much necrosis and haemorrhage with benign growths. Another feature suggestive of malignancy is the association of tumour cells with the vessel walls, and in many cases the presence of strands of cells lying free in blood spaces. The marked tendency towards thrombosis in the vessels is more commonly seen in malignant than in benign tumours.

Examples in the literature of large cortical adrenal tumours are rare. Among the largest is the case recorded by Holmes (1925) in which a tumour weighing 1,025 g. (2 lb. 4 oz.) and measuring 17 by 14 by 9.5 cm. was found to be of benign type, and that of Anderson *et al.* (1943), who described a malignant cortical tumour weighing 1,780 g. (3 lb. 14 oz.) in which a solitary metastasis was found in the liver. The latter case is of interest, as the appearance of the tumour was identical with those described above, although a relatively anaplastic histological pattern was recorded.

It may again be stressed that histologically benign tumours merge imperceptibly into the malignant varieties, and in the absence of demonstrable metastases opinions might well differ about their correct interpretation. It might be argued that haemorrhage into a simple adenoma is the more correct view in the above cases, but it is my

opinion, based on the reasons already given, that these are examples of slowly advancing malignant tumours.

Summary

Two cases of very large cortical adrenal tumours are presented.

The complete absence of localizing clinical signs is noted.

The importance of their recognition in order to institute post-operative treatment is stressed.

The question of their benign or malignant origin is discussed.

My grateful thanks are due to Mr. G. T. Mullally and Sir Stanford Cade for permission to make use of the case records.

REFERENCES

- Anderson, A. F., Hain, A. M., and Patterson, J. (1943). *J. Path. Bact.*, **55**, 341.
 Broster, L. R., and Vines, H. W. C. (1933). *The Adrenal Cortex*. London.
 Cahill, G. F., Melicow, M. M., and Darby, H. H. (1942). *Surg. Gynec. Obstet.*, **74**, 281.
 Cecil, H. L. (1933). *J. Amer. med. Ass.*, **100**, 463.
 Glynn, E. E. (1912). *Quart. J. Med.*, **5**, 157.
 Holmes, G. (1925). *Ibid.*, **18**, 143.

A SECOND EXAMPLE OF THE "DUFFY" ANTIBODY

BY

ELIZABETH W. IKIN, B.Sc.

A. E. MOURANT, D.Phil., D.M.

AND

GERTRUDE PLAUT, B.A., M.D.

(Blood Group Reference Laboratory, Ministry of Health and North London Blood Supply Depot, National Blood Transfusion Service)

Cutbush, Mollison, and Parkin (1950) have described an antibody which defines a new blood-group system. The antigen (*Fy^a*) distinguished by this antibody was found to be present in 66% of the English population, and appeared to be independent of the *ABO* and *Rh* groups of the blood tested. The antibody, anti-*Fy^a*, was found after a transfusion reaction in a haemophilic patient who had received previous transfusions. It was accompanied by the *Rh* antibody anti-*D*, which could, however, be removed by absorption. The reactions of the new antibody could be detected with certainty only by means of the Coombs test.

A second example of the same antibody has been detected in the serum of Mr. J. S. In or about the year 1943 he had twice been transfused for haemorrhage from a peptic ulcer. In August, 1949, he was again admitted to hospital for the same condition. His blood was found to be group *AB*, and two bottles of group *B* blood were directly matched against his serum at the hospital. He had a severe rigor after receiving about 20 ml. of blood from one bottle. The transfusion was interrupted and begun again from the other bottle with a similar result. It unfortunately proved impossible to trace the donors of these bottles of blood.

A full investigation of the patient's blood was then carried out. He was found to be group *A₂B*, *R₁R₁*, (*CDe/CDe*), and his serum, though containing no saline agglutinins, was found by means of the Coombs test to contain an antibody which reacted with more than half of the red-cell samples tested. It was almost immediately