FIG. 4.—Case 4. Thorax after removal of aberrant intrathoracic toxic goitre.

lifted out of the thorax and removed after two venous channels below had been divided. These appeared to pierce the pericardium to empty direct into the auricle. The patient was given a blood transfusion at the conclusion of the operation and, as might be expected, had a severe thyrotoxic reaction during the days following. Her convalescence was complicated by collapse

of the lower lobe of the left lung, but she eventually made a good recovery with disappearance of all signs of thyrotoxicosis. She has been well since that time. Fig. 4 shows the condition of the thorax nearly three years after operation, and Fig. 5 depicts the specimen removed.

FIG. 5.—Case 4. The specimen removed, showing vascular pedicle. Weight, 354.5 g.

place in the surgery of goitre. I have also described a form of intrathoracic goitre in the posterior mediastinum which has seldom been recognized or described. Lastly, I have shared with you the alarming experience of operating on a true, very large, aberrant intrathoracic goitre with severe thyrotoxicosis, an association which, so far as I know, has never been described.

[The lecture was illustrated with a coloured film of the operation of mediastinotomy for removal of the thymus gland.]

REFERENCES

- Dunhill, Sir T. P. (1922-3). *Brit. J. Surg.*, **10**, 4.
Joll, C. A. (1932). *Diseases of the Thyroid Gland*. Heinemann, London.
Mora, J. M., et al. (1944). *Surg. Gynec. Obstet.*, **79**, 314.
Rives, J. D. (1947). *Trans. Amer. surg. Ass.*, **65**, 417.

ORAL HORMONAL THERAPY FOR MENSTRUAL DISORDERS

BY

G. I. M. SWYER, M.A., D.M., D.Phil., M.R.C.P.
Endocrinologist, University College Hospital Medical School

Disorders of menstruation comprise a large and important group of conditions for which many women at some time or other seek medical advice. While it is true that they are essentially the outward symptomatic manifestation of underlying disturbances, in such a large proportion is the condition dysfunctional that it is permissible to consider them as disease entities in their own right and to prescribe treatment on that basis. Clearly the adoption of this attitude presupposes the elimination by the appropriate methods of systemic or local disease which might itself give rise to the menstrual disorder in any particular case, though it must be admitted that this is often easier in theory than in practice. In the account which follows it is to be understood that, except where specifically indicated, only disorders of function are under consideration and that where underlying organic disease exists the correct treatment is that directed towards the cure of the disease itself.

During recent years the ready availability of hormones—particularly the steroid hormones and their synthetic analogues—has led to their increasing use in the treatment of menstrual disorders, and the fact that oestrogens, androgens, progestogens, and gonadotrophins have all had their enthusiastic advocates would suggest strongly that empiricism has played an altogether excessive part in determining such therapy. It is not, however, the purpose of this communication to discuss the relative merits of the various techniques that have been proposed; on the contrary, attention will be directed solely towards a therapeutic regime based upon that originally described by Hamblen (1945) and involving the oral administration of oestrogens and progestogen. This method has a rational basis (which may, of course, be founded upon partially false premises) and, in addition, has the great practical advantage of avoiding parenteral therapy, inconvenient to doctor and patient alike.

The results of treatment of 52 patients complaining of menstrual disorders form the basis of the following report. These cases, treated in the Endocrine and Fertility Clinics at University College Hospital, have been selected solely on the grounds of the treatment having extended over a sufficient length of time and of adequate follow-up. They are conveniently grouped into cases of prolonged and excessive bleeding with irregular cycles (19) and with regular cycles (7); cases of irregular (chiefly scanty and infrequent) and anovular but otherwise regular cycles (13); and cases of secondary amenorrhoea (13). The plan followed consists, first, of a detailed description, along somewhat dogmatic lines, of the methods of treatment that have been adopted, and, secondly, of a critical evaluation of the results achieved. It is my intention to present only what I have myself found to be the case, for which reason no specific references to the results of other workers will be made. In addition to the inherent justification for such a course there is the further fact that, so far as is known, no report of the use of the precise regime considered here has hitherto appeared.

Control of Excessive Uterine Bleeding

Prolonged and excessive uterine bleeding is the most urgent menstrual disorder for which patients seek treatment.

It is convenient, therefore, to consider this aspect of therapy first, and in so doing emphasis may be laid at the outset upon the necessity for establishing the *dysfunctional* nature of the condition. Particularly in the case of older women must steps be taken to exclude neoplastic causes; however, in most cases admission to hospital for the necessary curettage is seldom possible without more or less considerable delay, during which the patient naturally expects relief. Unless, therefore, the patient's condition is so extreme as to warrant immediate admission, it is clearly permissible to employ such therapy as may be expected to control dysfunctional uterine bleeding, when, should the treatment fail, the case automatically becomes one for urgent admission, the suspicion of neoplasm being rendered greater by the failure of hormonal treatment.

For the control of excessive uterine bleeding oestrogens are generally effective and reliable, statements to the contrary notwithstanding. The action is undoubtedly initially that of a direct haemostatic, and, in general, adequate dosage secures complete arrest or obvious amelioration of the bleeding within 48 hours of the beginning of treatment. Stilboestrol in a dose of 2-5 mg. daily is suitable for this purpose, but it has the drawback of producing toxic symptoms in a small but important proportion of patients (important because, as will appear, interruption of therapy is not permissible). Dienoestrol and hexoestrol, in my experience, have proved far less effective than stilboestrol, even allowing for the differences in potency, and it is to be hoped that ethinyl oestradiol may solve the problem posed by the tendency of stilboestrol to be poorly tolerated. However, limited experience with this new and highly potent synthetic oestrogen is not too encouraging at present.

In the event of the initial dose proving insufficient for the arrest of haemorrhage, there should be no hesitation in doubling it; the patient, in fact, is told to do this if after 48 hours there is no sign of improvement. Whatever be the appropriate dose for the arrest of haemorrhage, it is clearly imperative that this dose must be taken daily after the bleeding has stopped, since to reduce it or to stop its administration would provoke an oestrogen-withdrawal bleeding a few days later. Daily administration should therefore continue for a total of 20 days. After an interval which may vary from three to 10 days, an oestrogen-withdrawal bleeding begins and oestrogen administration should be recommenced so as to start the process of cycle regulation (see below). Occasionally, bleeding recommences while the course of oestrogen is being taken; in such an event the patient should be told to stop the treatment, wait for the fifth day of the bleeding episode, and then start treatment again. If the initial dose has been 5 mg. of stilboestrol daily, it may usually be reduced to 2 mg. daily in subsequent courses, but it seems that a smaller dose is seldom advisable in these early stages of treatment.

There are other hormonal methods for the control of excessive uterine bleeding, such as the use of testosterone and progesterone (preferably in combination), but since they involve parenteral administration no further attention will be given to them here. In severe cases the efficacy of complete curettage deserves careful consideration, and this measure is rendered more effective if oestrogen administration is begun at the same time.

Where anaemia of more than minimal degree exists treatment with haematinics should be given *pari passu* with other therapy.

Cycle Regulation

The immediate control of uterine bleeding having been achieved, it now becomes necessary to attempt regulation

of the cycles so that normal regular bleeding episodes may ensue. Mere cessation of treatment will be followed in most cases by a relapse to the pre-treatment condition; ignorance of this fact has been the basis for many opinions that endocrine treatment has little value in dysfunctional uterine bleeding. Since the great majority of cases of irregular excessive uterine bleeding are associated with failure of ovulation, treatment cannot claim to have been fully curative unless ovulation has been re-established. Where there is a complaint of infertility this is clearly of paramount importance, but its desirability in other cases is also worthy of emphasis, since relapse is decidedly less probable if ovulation has been re-established than if the succeeding cycles are anovular. In cases of climacteric bleeding the position is plainly different, and attempts at inducing ovulation are doomed to failure; certain modifications in treatment are therefore necessary, and will be considered later.

On the fifth day of the oestrogen-withdrawal bleeding resulting from the controlling course of treatment, a new course should be started: 2 mg. daily of stilboestrol is usually a suitable dose and should continue for 20 days. The third course starts on the fifth day of the resultant "period" and is similar in all respects to the preceding course. The effects of the oestrogen therapy are fivefold: there is direct uterine haemostasis; the pituitary is inhibited and as a result ovarian inhibition follows; the pituitary-ovarian relationship which had previously resulted in a more or less constant oestrogen output by the ovary is brought to an end; and finally, when the oestrogen administration is stopped, there is a temporary release of pituitary activity. It is probable that the regular releases of pituitary activity occasioned by the regular cessation of oestrogen therapy play an important part in determining the resumption of cyclic pituitary function, and at this point the opportunity may be taken of stressing the fundamental importance of absolute regularity of treatment if regularity of results is to be obtained. In this connexion it has been found useful to employ a duplicated form providing the necessary instructions to the patient, and in which the appropriate data (number of tablets, number of days, etc.) are written. I use the following:

Start taking.....small tablets daily on the...../.....th day of the cycle for.....days.

Start taking.....large tablets daily on the...../.....th day of the cycle for.....days.

Wait for the period to commence, and on the...../.....th day of the cycle start taking.....small tablets daily for.....days.

("Small tablets" refers to oestrogens, "large tablets" to ethisterone; the form allows either the date for starting the course of tablets or the day of the cycle to be specified, unnecessary words being deleted as appropriate.)

The fourth, fifth, and sixth cycles are conditioned by combined treatment with oestrogen and progestogen. The following appears to be a satisfactory regime: 2 mg. of stilboestrol is given daily for 20 days, starting on the fifth day of the cycle, and from the 15th day of the cycle ethisterone (anhydrohydroxyprogesterone) in a dose of 20 mg. daily is given for 10 days. During the last 10 days of the treatment course both hormones are thus taken simultaneously. For the fifth course the dose of stilboestrol is reduced to 1 mg. daily and that of ethisterone increased to 30 mg. daily; for the sixth course the daily doses are 0.5 mg.

and 40 mg. respectively, other details of administration being the same as in the fourth cycle. With the combined treatment the interval between cessation of treatment and onset of uterine bleeding is less than that with oestrogens alone, usually being two or three days as compared with four to six days. This regime results in ovulation in a useful proportion of cases, as will be shown later. With the low dose of stilboestrol employed in the last treatment cycle, ovulation may occur during that cycle (as may be concluded from the occurrence of conception during such a cycle in patient No. 35—see below). There seems, in fact, to be good reason to conclude that the administration of oestrogen and progestogen has an action on the pituitary gland, causing the latter to function in the manner necessary to determine ovulation. Regular ovular cycles follow cessation of treatment in successful cases, regular anovular cycles or the occurrence of irregular (but normal) periods at long intervals in less successful cases.

In some cases difficulty may be experienced in the initial production of regular oestrogen-withdrawal bleedings; it seems wise in these circumstances to persist with oestrogens alone until regularity has been achieved before starting the combined courses of treatment. As mentioned previously, the induction of ovulation is not attempted in patients with climacteric bleeding, and, in these, several regular courses of oestrogen therapy, the dose per course being reduced gradually, are usually adequate to secure the occurrence of moderate bleeding episodes leading to a calm menopause. For these patients individualization of therapy is essential, and attempts at following a rigid system are unlikely to yield satisfactory results.

Regular Menorrhagia

Excessive uterine bleeding of an entirely different type is that which occurs as regular periods in which the loss of blood is very heavy, though not necessarily prolonged. In many cases premenstrual endometrial biopsies show secretory changes, so that the occurrence of ovulation may be inferred.⁹ There seems to be little doubt that the cause of the excessive loss is different from that in the irregular type previously considered. Investigation may reveal the presence of fibroids, hypothyroidism, anaemia, or various other local or systemic conditions, and thereby indicate the correct line of treatment. Careful questioning may indicate an emotional basis, the correction of which, though often difficult, may effect a cure. There remains, however, a residue of cases in which no obvious underlying disturbances can be discovered; in these, treatment with hormones is seldom effective and is very likely to make matters worse by disturbing the menstrual rhythm. In such cases good results can be achieved by correcting the accompanying anaemia that is nearly always present and by the administration of a non-saponifiable liver extract (antimenorrhagic factor "glanules," Armour). It is probable that the active principle of this extract, whatever it may be, has a direct action on the blood vessels. The chief drawback, in my experience, is the necessity for uninterrupted treatment, at a dosage of two capsules thrice daily, in order to prevent relapses.

Irregular (Chiefly Scanty and Infrequent) and Anovular Cycles

Patients whose complaints are referable to menstrual disturbances comprised under the above heading are of two types: those whose primary complaint is of infertility, and those who believe the menstrual irregularities are themselves indicative of ill-health. For the latter, after systemic diseases have been excluded, reassurance is theoretically

sufficient if child-bearing is not contemplated, though in practice this may be impossible to achieve, so deep-rooted is the conviction that unless a woman has a "good" period at reasonably regular intervals something must be "wrong." If present sterility and future child-bearing are pertinent considerations, the production of ovulation, and preferably regular ovulation, is the obvious object of treatment.

Where the cycles are irregular, regulation as well as induction of ovulation is necessary, and, for reasons already given, treatment is best started with oestrogens alone, to be followed later with oestrogens and progestogens. The regime described under "Cycle Regulation," using stilboestrol and ethisterone, is suitable for this purpose. Where the cycles are regular but anovular, cycle regulation is unnecessary, and the combined treatment alone may be expected to produce ovulation in a useful proportion of patients so treated.

Secondary Amenorrhoea

Where no spontaneous period has occurred in the preceding year or more, treatment along the lines described usually produces withdrawal bleedings, but, as will be seen, the resumption of regular spontaneous cycles after the completion of treatment is exceptional.

Results of Treatment

It is now proposed to offer evidence for the foregoing statements, based on experiences with the group of patients already described. This will be mainly of summary type, but reports of more outstanding cases are given in some detail. The overall results of treatment are given in Table I.

TABLE I.—Overall Results of Oral Treatment Summarized

Group	No. of Patients in Group	Results of Treatment*				
		Excellent	Satisfactory	Failure	Ovulation After Treatment	Pregnancies
1. Prolonged and excessive bleeding; irregular cycles	19	7	9	2	8	3
2. Excessive bleeding; regular cycles	7	5 (A.M.F. 4 myomectomy 1)	—	2	—	—
3. Irregular (chiefly scanty and infrequent) periods	9	4	2	3	3	1
Regular anovular cycles	13	7	3	3	7	3
4. Secondary amenorrhoea	4	3	1	—	4	2
Totals	52	20	13	16	15	6

* These results do not include two spontaneous cures (Groups 1 and 4) and one pregnancy following gonadotrophin therapy after failure of oral treatment (Group 4).

A.M.F. = Antimenorrhagic factor "glanules" (Armour).

1. Prolonged and Excessive Uterine Bleeding (Irregular Cycles)

The ages of the 19 patients in this group ranged from 13 to 45 years, and the duration of symptoms before treatment from four months to 15 years. There were two patients (Nos. 2 and 3) in whom psychogenic factors loomed large; good initial control was secured in one of these, though relapse soon occurred in spite of continuing treatment, but removal of the cause—the patient was very unhappy as a student nurse at another hospital and very happy on giving up this activity—produced a cure. In the other case the treatment was never taken properly and so was a complete failure; this patient was referred for psychotherapy. One patient (No. 4), with symptoms lasting six months, underwent spontaneous cure, a secretory premenstrual biopsy

FIG. 1.—Chart of Case 6

having demonstrated the re-establishment of ovulation. In the remaining patients the immediate control of bleeding was good in 14 instances, fair in one, and poor in one; this last patient (No. 15) underwent a second course of treatment, good immediate control being obtained on this occasion.

In the earlier part of this study the combined treatment courses were of constant dosage—stilboestrol, 2 mg. daily for 20 days, and ethisterone, 20 mg. daily for the last 10 of these days, in each of the three combined treatment cycles; these are indicated by the term “C.D.” (constant dosage). Where treatment has followed the plan previously described, the term “D.S.—I.E.” will be used as symbolical of the diminishing stilboestrol and increasing ethisterone dosages. Endometrial biopsies* were taken before treatment in nine cases; they showed a proliferative type of endometrium in seven cases and cystic glandular hyperplasia in two. In no case, therefore, was evidence of ovulation obtained. Biopsies were taken after treatment in eight instances (excluding the case of a spontaneous cure) and showed secretory changes in seven of them. The single instance of a post-treatment proliferative biopsy (No. 9) followed a “C.D.” course of treatment, and relapse occurred three

months afterwards; a “D.S.—I.E.” course of treatment was then used, and three months after the end of this second course the endometrium was found to be secretory. Three pregnancies have occurred 2 months, 5½ months, and 9 months respectively after the end of treatment. As post-treatment secretory biopsies had been obtained in two of the pregnant patients, evidence for the re-establishment of ovulation is therefore available in eight of the treated cases.

Results have been excellent in seven of the 18 treated cases, regular ovular cycles having been established or conception having occurred; they have been satisfactory in nine cases, the resulting periods being irregular (but otherwise normal) or anovular, or subsequent relapse having occurred; and there have been complete failures in the two psychogenic cases. Further courses of treatment have been necessary in six of the nine “satisfactory” cases. Five patients received the “C.D.” course of treatment; an “excellent” result followed in one of these only, further treatment being necessary in the remainder.

Some representative case reports are given below.

Case 1.—A schoolgirl aged 13. Menarche 11 years 9 months. Periods were regular for nine months, lasting five or six days and recurring about every 28 days. Since then there has been irregular prolonged bleeding. No flooding. Examination under anaesthesia (August 28, 1947) revealed an anteverted slightly enlarged uterus; curettage produced polypoidal proliferative endometrium. Heavy, irregular bleedings followed. Treatment,

*In all the instances recorded the endometrial biopsy has been taken during a bleeding episode or in the pre-bleeding phase, as evidenced by the occurrence of the period within a few days of the biopsy.

FIG. 2.—Chart of Case 14.

begun December 1 with stilboestrol 3 mg. daily, brought prompt arrest of bleeding. In the 12 months following cessation of treatment there have been only three periods, all of normal type. Further treatment is contemplated.

Case 4.—An unmarried packer, aged 22. Menarche 12½. Periods regular, lasting three days (rather scanty) and recurring every 28 days. In December, 1947, she moved house and changed her job; a period started December 8 and lasted for three months. Following this there was irregular bleeding until she was admitted on April 14, 1948; examination revealed no abnormalities, and the bleeding stopped the next day. Since then regular periods have occurred, although no treatment had been given, and a biopsy taken at the start of the second period showed a normal secretory endometrium.

Case 6.—A housewife aged 23. Menarche 15. Periods were irregular, lasting five or six days and recurring at intervals of one to three months. The irregularity persisted, passing into amenorrhoea, which lasted for three years. During the year preceding admission prolonged irregular heavy bleedings had been the rule in spite of treatment with various hormones, including oestrogens, progesterone, and gonadotrophins. At the time of admission bleeding had been continuous for four weeks. Examination revealed no gross abnormalities, and bleeding was controlled with four daily intramuscular injections each of 25 mg. testosterone propionate and 10 mg. progesterone. On the fifth day of the resultant progesterone-withdrawal bleeding, treatment with stilboestrol, 2 mg. daily, was started. The case is illustrated in Fig. 1, which shows the occurrence of regular cycles during and after the end of treatment. Pregnancy was diagnosed nine months after the completion of treatment.

Case 9.—A housewife aged 27. Menarche 12. Periods were irregular for the first year or so, lasting six or seven days and recurring every 14–50 days. They were heavy. Regular periods followed until six months before she was first seen. At that time irregular intermenstrual spotting began, gradually developing into prolonged irregular and excessive bleeding. She had had one child three years before. Examination revealed no abnormalities. Bleeding was easily controlled with stilboestrol, 5 mg. daily, and regular cycles followed the "C.D." course of combined treatment. A biopsy taken just before the second spontaneous cycle showed a proliferative endometrium. Shortly afterwards a relapse ensued, and dilatation and curettage was performed because of the severity of the bleeding. Subsequent treatment with a "D.S.-I.E." course led to the re-establishment of normal ovular cycles.

Case 14.—A housewife aged 35. Menarche 12. Periods were regular, lasting three days and recurring every 28 days. She had had two pregnancies, the last eight years previously. Five months before she was first seen continuous heavy bleeding began (October, 1947). In January, 1948, dilatation and curettage revealed cystic glandular hyperplasia, and bleeding ceased for three weeks. Moderate bleeding then followed for two weeks, followed by heavy bleeding for a further two weeks. Stilboestrol, in a dose of 3 mg. daily, reduced the bleeding considerably, and after one week the dose was increased to 5 mg. Slight oozing persisted, becoming a withdrawal bleeding of the usual type on cessation of oestrogen administration. Thereafter fairly satisfactory control was obtained, a secretory endometrial biopsy being found in the third cycle after cessation of treatment. Four months later, however, prolonged

FIG. 3.—Chart of Case 31

bleeding began again, and further treatment is now being given (Fig. 2).

2. Regular Menorrhagia

The ages of the seven patients in this group were between 16 and 39 years, and the duration of symptoms at the time when the patients were first seen varied between one and 16 years. Secretory endometria were found in three patients before treatment, the endometrium being proliferative on three occasions (not necessarily premenstrual) in another patient, on whom myomectomy was eventually performed, after the failure of several forms of therapy, with complete cure. Poor results with oestrogens were obtained in two cases and with antimenorrhagic factor in two cases, one of which had the myomectomy described above. The other failure with antimenorrhagic factor is considered to be a case of emotional menorrhagia, and is described briefly below.

Case 21.—A housewife aged 25. Menarche 11; except for amenorrhoea for one year at 17, periods were regular, lasting three or four days and recurring every 28 days. Some time ago the patient came under the influence of a fortune-teller, who predicted that she would lose her first child; in 1944 she gave birth to a stillborn foetus with spina bifida. There followed two very heavy periods; then she conceived again, a normal infant being born in 1945. Since then she has worried over the possibility of losing her surviving child, and the periods, not quite so regular as before, have been very heavy, lasting eight or nine days. Examination revealed a large cervical erosion but no other abnormalities; curettage produced secretory endometrium. Haemoglobin 96%. Treatment with anti-

menorrhagic factor had no appreciable effect. She has been advised to have another pregnancy.

Good results with antimenorrhagic factor were obtained in four cases; in one of these incapacitating dysmenorrhoea has also been relieved; in two others retroversions were present, and correction with Hodge pessaries had had no effect on the menorrhagia.

3. Irregular (Chiefly Scanty and Infrequent) and Anovular Cycles

The ages of the 13 patients in this group ranged from 17 to 42 years and the duration of symptoms from seven months to 19 years in the cases of oligomenorrhoea. Biopsies taken before treatment in five cases showed proliferative endometria; after treatment four secretory biopsies were obtained. Evidence of ovulation occurring after treatment (secretory endometrium, conception, ovular type of basal temperature chart) was obtained in six cases. In three cases of oligomenorrhoea regular withdrawal bleedings occurred during treatment, but no spontaneous cycles followed its cessation; these, then, were complete failures. In two other cases regular spontaneous cycles were produced, but these were most probably anovular; in the remainder, ovulation was either definitely or possibly produced in one or more cycles following cessation of treatment. One patient conceived (see below, Case 35). There were four cases of regular anovular cycles; after treatment evidence of ovulation was obtained in all four and two conceived (Cases 32 and 34). Three case reports are given below.

FIG. 4.—Chart of Case 45

Case 31.—A married part-time secretarial worker aged 26. Menarche 15; periods always irregular, lasting six or seven days and recurring at two-monthly to three-monthly intervals. On examination moderately obese; external genitalia normal; uterus of normal dimensions (sound passed $3\frac{1}{2}$ in.—8.75 cm.); vaginal smear of castrate type; B.M.R. +3. Treatment with 1,000-calorie diet, thyroid sicc. gr. 1 (65 mg.) daily, and dextro-amphetamine sulphate, 5 mg. thrice daily, produced moderate weight reduction (15 lb. (6.8 kg.) over a period of seven months). The doses of steroid hormones given are shown in Fig. 3, and it can be seen that regular periods have followed cessation of treatment. The basal temperatures are suggestive of ovulation, but the endometrial biopsy produced only minute fragments of endometrium in spite of repeated scraping, and it seems reasonable to infer that some degree of unresponsiveness on the part of the endometrium must exist.

Case 34.—A housewife aged 28. Menarche 14. Periods always regular, lasting three days and recurring every 28 days. Patient investigated for infertility; basal temperatures suggested anovular cycles, confirmed by premenstrual endometrial biopsy. After a "D.S.—I.E." course of treatment the premenstrual biopsy was found to be secretory, and basal temperatures many months later were of ovular type. She conceived, but aborted at 13 weeks.

Case 35.—A married part-time secretarial worker aged 32. Menarche 13; periods always irregular, lasting generally three to four days and recurring at intervals of three months to two years. She had had various treatments, including thyroid, stilboestrol, and gonadotrophins. Examination revealed no abnormalities. A biopsy taken on the first day of a presumably spontaneous period showed bleeding from a proliferative endometrium. Treatment started with stilboestrol, 5 mg. daily, and progressed to the usual "D.S.—I.E." course. No withdrawal bleeding followed the last course of treatment, and two months

after the last period the Friedman test was found to be positive. It is reasonable to conclude that ovulation occurred during the last course of treatment. The pregnancy is progressing normally.

4. Secondary Amenorrhoea

The ages of the 13 patients in this group ranged from 17 to 37 years and the duration of symptoms from one to seven years. In one case (No. 41) there was spontaneous resumption of regular ovular cycles (as indicated by the basal temperatures) after 16 months' amenorrhoea without treatment. In another case (No. 45), with two-and-a-half years' amenorrhoea, a castrate type of vaginal smear and a B.M.R. of -16.5 were found; treatment with thyroid substance and later with dienoestrol and ethisterone along lines previously described produced regular spontaneous cycles, which, however, appeared to be anovular as judged by the basal temperature chart (Fig. 4). Spontaneous periods were resumed nine weeks after the cessation of treatment in a third case (No. 42), in which there had been seven years' amenorrhoea. There was a further case (No. 48) in which, the second pregnancy having produced an anencephalic monster, two scanty periods followed, to be succeeded by amenorrhoea lasting one year; withdrawal bleedings were produced by the usual oral treatment, but there was no spontaneous resumption of menstruation. Conception, however, occurred during gonadotrophin treatment (pregnant mares' serum, 500 i.u. intramuscularly daily for eight days, followed immediately by chorionic gonadotrophin, 500 i.u. intramuscularly, daily also for eight days), but was followed by abortion at approximately seven weeks. Within two months conception again occurred, as proved

by a positive Friedman and male-toad test; and as a prophylactic measure six pellets of progesterone, 25 mg. each, were implanted subfascially. The pregnancy continues. The remainder of the cases of this group may be summarized as complete failures, in so far as no spontaneous periods followed the cessation of treatment.

Intolerance to Stilboestrol

Treatment with stilboestrol was attempted in 46 of the cases reported here; intolerance was shown by seven (15%), necessitating change of treatment. Dienoestrol was used in six cases; it gave good results in two of these (Nos. 37 and 44) and poor results in the remainder.

Discussion

The foregoing results appear to justify certain definite conclusions: (1) oral stilboestrol is an effective and reliable agent for controlling prolonged and excessive uterine bleeding; (2) given in the manner described, and subsequently combined with ethisterone, it can produce excellent end-results in about 40% of patients having irregular prolonged and excessive uterine bleeding, and satisfactory results in a further 50% of these patients; (3) it has no value in the treatment of regular menorrhagia and little or none in the treatment of secondary amenorrhoea of more than a year's duration; (4) the combined treatment seems to be of considerable value in producing ovulation in cases of regular anovular cycles and of less definite value in the treatment of oligomenorrhoea.

The relapse rate in apparently cured cases of irregular prolonged and excessive uterine bleeding is comparatively high; relapse occurred in five patients, three of whom had received a combined course in which the dosages of stilboestrol and ethisterone remained constant in successive months. In only one of the cases wherein relapse had occurred was evidence of ovulation after treatment obtained, which provided the justification for the remark made previously that the re-establishment of ovulation renders the liability to relapse considerably less. The fact, however, that relapse may occur even when ovulation has been restarted is not altogether surprising, and is a reflection of the instability of the menstrual rhythm which is a fundamental defect in all cases of menstrual irregularity.

The question of dosage in the controlling course of stilboestrol deserves further consideration. But for the liability to produce toxic symptoms, and the obvious disadvantages of this, there would be much to be said for erring on the side of excessive rather than inadequate dosage. There seems little doubt, however, that many patients who have experienced toxic symptoms from stilboestrol will continue to have them, even on much smaller doses, so that avoidance of such side-effects at the outset is much to be desired. The dosage levels in common use, such as "5 mg. t.d.s.," are for most purposes grossly excessive and largely responsible for the prevalent opinions regarding the high toxicity of this excellent, potent, and cheap oestrogen. A dose which is sufficient to have a haemostatic action may in fact prove insufficient to prevent the recurrence of bleeding while it is still being taken: it is probable that this occurs most often when the endogenous oestrogen production is high (as where cystic glandular hyperplasia exists). A reasonable explanation would be that the oral oestrogen, by inhibiting the pituitary, causes a reduction of the endogenous oestrogen output, so that the combined stimulus to the endometrium becomes reduced, the overall effect being similar to an oestrogen-withdrawal bleeding. Had the dose been higher the combined oestrogen level would have fallen

FIG. 5.—Diagrammatic representation of the effects of moderate and high exogenous oestrogen doses in cases with low and high endogenous oestrogen levels.

relatively less and so no withdrawal bleeding would have resulted (Fig. 5). This recurrence of bleeding need not be considered as of serious import if the advice given—namely, to stop oestrogen administration and to start again five days later—is followed. The pituitary inhibition will not have passed off completely within this time and the endogenous oestrogen production will not have reached its pre-treatment level, so that the second oestrogen course will be far less liable to interruption through the inopportune return of bleeding. One final point regarding oestrogen dosage may be made: in general it is safe to assume that the therapeutic dosages of oestrogen and the tolerance towards them bear a direct relation to the level of endogenous oestrogen production. In practice, this means that dosage must be carefully limited at either end of the reproductive period and can be almost unlimited during pregnancy.

Great interest attaches to the not infrequent occurrence of ovulation after treatment in patients who were presumably or definitely anovular before treatment. In the series of patients reported there were, among the cases of irregular excessive bleeding, oligomenorrhoea, and regular anovular cycles, 14 instances in which definite evidence of non-occurrence of ovulation was obtainable and a further 17 in which it was presumptive. Evidence of ovulation after treatment was obtained in 15 of these cases, representing nearly 50% success. (In one patient with menopausal menorrhagia, induction of ovulation was not attempted.) In a further six patients the opportunity of deciding whether or no the post-treatment cycles are ovular has not arisen, so that the actual success rate may be even higher. To be set on the debit side of the account are the two instances of spontaneous resumption of regular ovular cycles—in one case following prolonged irregular bleeding, and in the other 16 months' amenorrhoea, both without any form of treatment. The superiority of the "D.S.-I.E." type of

combined course over that represented by "C.D." for the induction of ovulation would appear to be demonstrated by the fact that ovulation was shown to have occurred after the "C.D." course in only two out of the nine cases in which it was used, whereas the "D.S.-I.E." course succeeded in one case where previously the "C.D." course had failed. Final conviction would seem to be provided by the occurrence of six pregnancies within 0 to 9 months of the completion of treatment, failure of ovulation having been the case prior to treatment in all of these patients.

A sobering note must, however, be struck by the conspicuous failure to produce ovulation in cases of long-standing secondary amenorrhoea as well as most cases of marked oligomenorrhoea.

It is tempting to speculate upon the mechanism whereby treatment with oestrogens and progestogen can lead to ovulation, but in the present lack of factual knowledge it would be unwise to proceed too far. That a direct action on the ovary could be implicated seems scarcely possible, so that one must invoke the anterior pituitary lobe as an intermediary, causing ovulation by the secretion of gonadotrophin. Unfortunately, although oestrogens can undoubtedly inhibit the pituitary, it is much less certain that they can have a stimulant action, and such an action is indeed strenuously denied by some. Be this as it may, it seems that the regular administration of oestrogens and progestogen, in the proper dosage, can so condition the pituitary gland as to secure its production of gonadotrophin in the manner necessary to cause ovulation.

Summary

A regime involving the oral administration of oestrogen and ethisterone for the treatment of menstrual irregularities has been described. Encouraging results have been obtained in cases of irregular prolonged and excessive uterine bleeding, using stilboestrol for the immediate control of bleeding and proceeding to treatment designed to regulate the cycles and re-establish ovulation. Moderate success was achieved in cases of oligomenorrhoea and regular anovular cycles; the treatment has had little value in cases of regular menorrhagia (for which the antimenorrhagic factor ("glanules," Armour) proved useful) and of long-standing secondary amenorrhoea.

The treatment appeared to be responsible for the re-establishment of ovulation in at least 15 out of 30 patients (not including cases of secondary amenorrhoea and menopausal menorrhagia) in whom failure of ovulation had been presumed or proved. Pregnancy occurred in six patients.

Intolerance to stilboestrol was encountered in only seven out of 46 patients to whom it was given.

[Restriction of space precludes publication of the details of all the cases in this study. However, for those interested, Tables II-V, listing these details in summary form, will be provided on application to the author.]

I wish to express gratitude to my colleagues in the Obstetric Hospital, University College Hospital, for their kind co-operation in this study, and to Messrs. Armour and Co., Ltd., for supplying some of the antimenorrhagic factor "glanules" used.

REFERENCE

Hamblen, E. C. (1945). *Endocrinology of Woman*. Thomas, Springfield, Ill.

The National Smoke Abatement Society has published as a booklet the proceedings of its conference held at Harrogate last autumn. The booklet includes the presidential address by Lord Simon of Wythenshawe on "A Survey of Domestic Smoke Abatement," and an address by Mr. Arthur Blenkinsop, M.P., Parliamentary Secretary to the Ministry of Health. It is obtainable for 4s. 6d. from the society at Chandos House, Buckingham Gate, London, S.W.1.

DEFICIENCY PATTERNS IN METALLIC POISONING

BY

WILLIAM HUGHES, M.D., M.R.C.P.

Resident Physician, Belmont Road Hospital, Liverpool

The account of the investigation which led finally to the discovery of 2:3-dimercaptopropanol (dimercaprol; British anti-lewisite—"BAL") (Carleton, Peters, *et al.*, 1946) makes one of the most fascinating chapters in the history of medical research. Peters (1936) gives as his starting-point the observation that the convulsions in thiamin deficiency could be traced to a biochemical lesion in which the pyruvate-oxidase system broke down through lack of thiamin. The enzyme was shown to consist of thiamin pyrophosphate linked with a sulphur-containing protein. The protein moiety of the system could be poisoned by iodoacetic acid, and thus the biochemical lesion could occur even in the presence of adequate thiamin. Iodoacetic acid is a mild vesicant—vesication is the outstanding effect produced on epithelial surfaces by lewisite. It was demonstrated that lewisite poisoned the pyruvate-oxidase system, and that its action was due to its avidity for the SH groups of proteins in the enzyme. The next step was to find a sulphur-containing compound which would compete with the protein for the available arsenic, and the dithiol, 2:3-dimercaptopropanol, was found to do this most effectively. Subsequent investigations have shown that dimercaprol is effective in other forms of metallic poisoning, including mercury (Longcope *et al.*, 1946) and gold (Burnett, 1947; Cohen *et al.*, 1947; Davison, 1947; Slot and McDonald, 1947).

While Peters's hypothesis hangs on the biochemical lesion of the pyruvate-oxidase system, it must not be forgotten that many other enzyme systems occur in which the coenzymes contain members of the B₂ complex, and that the specific proteins to which these are attached may also contain SH groups susceptible to poisoning by heavy metals. There is also the possibility that vitamins and factors outside the range of the B complex may form similar linkages with SH-containing proteins. Wherever such systems occur there is the possibility of interference with the protein by a heavy metal. We should expect to find, therefore, as a result of the "biochemical lesions" of metallic poisoning, many anatomical lesions, patterns of lesions, and syndromes which we are accustomed to associate with specific vitamin deficiencies. That these do occur is quite apparent from the accounts of cases in the literature of the subject. There are good examples in the comprehensive review by Carleton, Peters, *et al.* (1946), and in the paper by Longcope *et al.* (1946). In the following case the pattern of lesions was that which might have been expected in vitamin-K deficiency.

Case 1

A man aged 61 was admitted on September 18, 1948, for rheumatoid arthritis. The history went back 20 years, when the condition started in the small joints of the hands.

On admission there was severe involvement of all the peripheral joints of the limbs and also some involvement of the spine, leading to pronounced kyphosis. Psychologically he was difficult to treat and did not co-operate in the physical therapy prescribed. He was therefore put on a course of sodium aurothiomalate ("myocrisin"), starting with a dose of 10 mg. weekly, subsequently increased to 50 mg. weekly. By January 19, 1949, he had received a total of 0.68 g., and on