REFRESHER COURSE FOR GENERAL PRACTITIONERS

CAUSES AND TREATMENT OF ECZEMA

BY

R. T. BRAIN, M.D., F.R.C.P.

Physician-in-charge, Skin Department, Royal Free Hospital and the Hospital for Sick Children; Physician, St. John's Hospital for Diseases of the Skin

Eczema may be defined as the eruption peculiar to the hypersensitive skin, the hypersensitivity being inherited or acquired. Although the histological features of eczema are essentially inflammatory in nature, the clinical picture of the early lesions, consisting of grouped, discrete, minute papulo-vesicles tending to crust or exude, is characteristic enough to distinguish it from a dermatitis due to an obvious irritant. It is evident, however, that an irritant which will produce a dermatitis characterized by erythema, oedema, heat in the skin, irritation, and disturbance of the skin metabolism is also likely, particularly in predisposed subjects, to produce a secondary eczematous response.

In eczema two groups of factors are important aetiologically. First, constitutional factors have to be considered, and these are particularly prominent in infantile eczema, which provides an interesting introduction to the inherited or constitutional factors that may be more difficult to understand when occurring later. Secondly, eczema may have its cause entirely in some primary or secondary irritant, often called an allergen, which may be the result of simple contact or be due to a pathological process remotely affecting the skin by damaging some other organ of the body. The simplest and most satisfactory solution of the problem of eczema is that occasionally discovered by a careful history, when the eruption is found always to follow or to be aggravated by contact with a specific allergen.

Confirmation of the suspected causative agent may be obtained by what is called the patch test. This consists in applying a very small quantity of the suspected irritant, or if it is a plant or fabric a $\frac{1}{4}$ sq. in. (0.4 cm.²) portion set in $\frac{1}{2}$ sq. in. (1.6 cm.²) of fine mesh bandage affixed to 1 sq. in. (6.5 cm.²) of adhesive plaster. A patch test with enough controls should be applied to the flexor aspect of the forearms or near the edge of the eczematous eruption, because the whole skin surface may not be sensitized to the irritant. If the patch produces an erythematous area, with or without small papulo-vesicles set in it, within 24 to 48 hours it indicates a positive specific reaction provided the plaster and the control-test substances produce no corresponding effect. Prepared plaster strips are now available and provide a convenient means of carrying out patch-testing. Many cases of eczema of obscure origin have been elucidated by this simple measure.

For a long time it was thought that allergens were necessarily of organic origin and were usually proteins, but it has now been established that many simple chemicals can act as allergens. One point of difficulty to the inexperienced in dermatology is the fact that one individual may be exposed for a long time to a mild irritant before developing a reaction to it, whereas another will produce an eczematous eruption after the first contact with the irritant. If a reaction occurs after the first contact it must be regarded as an idiosyncrasy, but since it has been established beyond doubt that sensitization can occur *in utero* it will rarely be possible to distinguish a true idiosyncrasy from an acquired hypersensitivity.

It may be difficult to recognize an external irritant as being the cause of eczema, but it is even more difficult when the eczematous eruption is secondary to some toxin produced by bacterial or mycotic infection. These secondary eruptions are sometimes called "ides," the best-known example occurring in epidermophytosis of the feet, when the epidermophytide affects the hands as a deep vesicular eruption often indistinguishable from the idiopathic eczematous eruption cheiropompholyx. The association of eczematous eruptions with epidermophyton infections is well established, and as the infection responsible may be limited to one or two toe clefts it is very likely that other foci of infection, such as in the teeth, tonsils, or sinuses, and all varieties of skin infection or injury may provide allergens causing secondary eczema, which is usually symmetrical and mostly affects the hands, arms or legs, before involving the face, neck, or trunk. Thus it is common for an ulcerative condition affecting the legs, or even a localized dermatitis such as occurs from the overtreatment of a simple injury, to be complicated by the sudden appearance of a symmetrical itching eruption presenting the grouped papulo-vesicles characteristic of simple eczema. When this happens and is followed by increased irritation of the skin, increased anxiety, and disturbance of sleep it is easy to understand how these factors, plus the damage caused by uncontrollable scratching, are likely to extend the eczema. An occupational dermatitis due to simple irritants such as lime, cement, or corrosive acids may in this manner be converted into an eczematous eruption which may run a chronic course and show no striking inclination to clear up even when the patient is away from work. Unless this mechanism of producing secondary eczema is understood and the importance of an anxiety state in maintaining the eruption is appreciated, the results of treatment are likely to be very disappointing.

Treatment: General Measures

The success of any scheme of treatment will depend on the recognition of the irritating factors with which the individual has been in contact, his reaction to the disease and to the various forms of treatment that have already been given, and on the choice of suitable local applications. It is essential to have the confidence of the patient, and this is not increased by disclaiming any knowledge of dermatology. It is often helpful to explain to the patient that his eczema is but the reaction of a tissue that has been offended and which has therefore become jumpy or hypersensitive and in consequence has to be humoured and protected from further insult. Doctor and patient should understand that the reaction in the skin produces substances (autolytic) which circulate in the blood or lymph channels and which are likely to provoke further eruptions of a similar nature. It is therefore better to abandon the idea of some specific cure and to concentrate upon soothing and protecting the skin, and when this is done a period of wholesome neglect may be beneficial. Acres of eczema have been cultivated by overtreatment.

When due thought has been given to the patient's general health, particularly to focal sepsis, constipation, dyspepsia, and insomnia, the relief of his irritation and anxiety often calls for a mild sedative such as small doses of phenobarbitone. Amphetamine or dextroamphetamine, 5 mg. once or twice in the early morning, will often relieve the depressed patient, and, provided it does not keep him awake, is undoubtedly useful. The obese seborrhoeic subject may tolerate more frequent doses, but the reaction to this drug must be carefully watched.

On the presumption that dilatation of the vessels in eczema is, as in the case of urticaria, due to histamine, the antihistamine drugs have been hopefully employed, and, while occasionally in acute phases they seem to be of considerable value, they are often profoundly disappointing. Even so, "benadryl," 50 mg. two or three times a day, may prove more helpful as a sedative than a barbiturate. The mechanism of itching is thought to be compounded of the sense of pain and that of touch, so that

FIG. 1.—The early lesions of eczema. Showing the grouped minute papulo-vesicles set in non-inflamed skin.

analgesics may be more useful than hypnotics, and acetylsalicylic acid or compound tablet of codeine should be tried. The more recently introduced analgesics have been used with varying success. It must be admitted, however, that no drug is always successful in controlling irritation.

Ephedrine hydrochloride in $\frac{1}{4}$ - or $\frac{1}{2}$ -gr. (32- or 16-mg.) doses can be well combined with phenobarbitone, and the combination is particularly useful in those cases of flexural eczema (Besnier's prurigo) which are not uncommonly associated with asthma. The synergic effect of a combination of atropine and barbiturate is suitable for the reactive individual with hyperhidrosis, and the addition of small doses of ergotamine may increase its sedative effect upon the cutaneous vessels. Seborrhoeic subjects of either sex are sometimes helped by small doses of stilboestrol, and the torpid patient with a dry skin who has eczema may derive benefit from small doses of thyroid, which is obviously contraindicated in the nervous and excitable patient. Autohaemotherapy or protein shock is occasionally used to provoke a response in a chronic phase.

FIG. 2.—Lichenized eczema of the flexures.

Local Treatment

Although many doctors and some dermatologists advise their patients not to use water on the skin, most patients with eczema will tolerate and even enjoy bathing in isotonic saline or 0.5–1% solution of sodium sulphate. To this bath may be added 2 to 3 oz. (57 to 85 ml.) of solution of coal tar, and 1 to 2 oz. (28 to 57 ml.) of the new emulsifying ointment of the *British Pharmacopoeia* may be creamed in a jug of warm water and added to the bath to give it emollient qualities, which are particularly beneficial in the generalized eczema of infants. Vegetable oils and liquid paraffin are useful for the occasional cleansing of an irritable skin, but their constant use raises the skin temperature by preventing evaporation of sweat and greatly enhances the risk of secondary infection, which is often a complicating factor in eczema. Further local treatment of eczema must be varied according to the phase.

Weeping Phase

In this phase lotions and powders are indicated. Calamine lotion, "siccolam," or liquid magnesia are bland and on drying leave a fine powdery film which tends to dry up exudate and at the same time to form a protective layer upon the surface. Calamine lotion is made more "drying" by the addition of 5% of titanium dioxide and is further improved if 2% of "bentonite" is added. These lotions may be modified by the addition of

FIG. 3.—Eczematous dermatitis in an elderly seborrhoeic subject. The left arm shows a confluent scaly dermatitis with lichenification about the wrist somewhat obscuring the eczematous pattern (which is clearly visible on the right forearm).

2 or 3% of solution of coal tar or 1% of strong solution of lead subacetate. If a clear lotion is required for use as a wet compress the tar and lead solutions may be used together, or the tannic-acid lotion of the *National Formulary*, which tends to coagulate the exudate and to harden the surface of the skin, may be employed. The latter lotion contains mercuric chloride and is therefore mildly antiseptic. As the normal pH of the skin is on the acid side (pH 5-6) an acid lotion may be regarded as somewhat more physiological. It may account for the success which follows the use of simple boric lotion, though boric acid is somewhat discredited as an antiseptic. Solution of proflavine (N.F.) or 1% of gentian violet may be used when pyogenic infection is present. Gentian violet has the disadvantage of obscuring the lesion and making observation of its progress difficult, if not impossible. Sometimes 1% of strong solution of lead subacetate and 2 or 3% of tar solution in boiled milk seem to satisfy the very irritable skin, and bland talcum powders are valuable in drying the lesions and in preventing friction between irritable skin surfaces.

Drying Phase

The skin may become uncomfortably tight in the drying phase, particularly on the face and flexures, and then a bland cream such as the oily calamine lotion or a zinc cream is indicated, but it should be applied very lightly and the treated area left uncovered. Eczema rarely tolerates cotton-wool and bandages, and the more lightly eczematous lesions are covered the less they irritate as a rule. Bland creams can be made with the emulsifying ointment bases now available in the B.P., but occasionally these agents act as irritants and must immediately be discontinued. In fact it is impossible to predict the reaction of the eczematous area to any application, however bland it may have proved in the prescriber's experience, and when the application is followed by redness, increased irritation, or weeping it should be discontinued. Subsequently the skin may tolerate the preparation, but to ensure success most careful observation is essential with every variation of treatment.

An oil-in-water cream may be prepared without the use of an emulsifying agent, with 1.25% of methyl cellulose and 10% of liquid paraffin in water. To this and to the other creams may be added 2 or 3% of solution of coal tar, but though tar is one of the most commonly used antipruritic substances in dermatology it occasionally acts as an irritant. For the erythematous and oedematous skin 2% of ichthyol in an oily cream is useful. Sometimes considerable benefit results from the alternation of a lotion or a cream with warm or cold saline compresses applied for a few hours. The latter soften and remove exudate. If pyogenic infection is present penicillin solution may be sprinkled on the surface of the compress.

The old starch compress is still worthy of a place in the treatment of crusted eczema, and penicillin solution may be applied to its surface, or proflavine solution may be painted on the infected area before the poultice is applied. Dermatologists rarely use the sulphonamides locally, because they are so apt to provoke sensitization of the skin, either to themselves or to sunlight; sometimes penicillin is equally irritating. Creams containing the antihistamine substances are occasionally of value, and they may be tried when other attempts at relieving the irritation of localized lesions fail. Creams and ointments containing local analgesics—for example, ung. benzocain. co. B.P.C.—are often temporarily effective, but their continued use usually does more harm than good.

Dry Phase

In this phase pastes are most useful, particularly when thickening or lichenification is present. Plain zinc paste or Lassar's paste, which contains 2% salicylic acid, is most commonly used, and 2 or 3% crude coal tar or 5% solution of coal tar may be incorporated to relieve irritation. Tar paste is the classical application for the lichenized flexural lesion which is the common feature of Besnier's prurigo, and sometimes the occlusion of an area of eczema under a thick film of paste, left undisturbed for four or five days, proves very beneficial.

Ointments are less often used by dermatologists, because they are apt to increase the heat and irritation of the skin. With the new emulsifying bases this criticism may be less justified, but ointments in this category really correspond in their properties to the creams which are mixtures of oil and water. Ointments of salicylic acid, tar, or zinc oxide are helpful on the very dry skin.

The ointment bases of the *British Pharmacopoeia* and the indications for their use in dermatology were described recently (Hadgraft and Brain, 1949). Mention was made of a formula for a pourable cream having a pH value of 5, with bland antiseptic, antipruritic, and cooling properties. It is often useful in the dry or drying phases of eczema. The formula is:

Emulsifying ointment	..	..	..	80 g.
Wool fat	..	..	..	50 g.
Liquefied phenol	..	..	..	20 ml.
Buffer solution pH5*	..	..	..	7 ml.
Water	..	..	..	to 1,000 ml.

*Prepared, according to the B.P. Appendix III, from potassium hydrogen phthalate and sodium hydroxide.

Physiotherapy

This has a limited field in the treatment of eczema. Undoubtedly the most useful agent is x rays in doses of 50-100 r, preferably with a kilovoltage of 60-70, but not exceeding 90. Doses of this order may be given every one, two, or three weeks on three occasions, and sometimes the relief from irritation is almost immediate. In the adult it might be suspected that there is a psychological element of suggestion in this response, but similar results have been obtained with infants and young children. In the latter, doses of 30-50 r are rarely exceeded, and it is quite certain that no ill-effects can follow such small doses, though it is important not to irradiate the genital organs. The chronic phases of eczema, particularly those associated with lichenification, are most likely to be improved by x-ray therapy. Equally good results have been obtained with low-voltage rays, including Grenz rays, and also by painting the lichenized areas with thorium X varnish containing 1,000 electrostatic units per ml.

Ultra-violet light, being an irritant in erythema doses, almost always aggravates acute eczema and may cause a relapse in a chronic case; sometimes an eczematous dermatitis affecting the face, hands, and exposed parts of the forearms may be entirely actinic. However, the exposure of a hypersensitive skin to cooling currents of air and to sky-shine exercises the normal mechanism of adaptation of the skin and its complicated vascular system to environment, and if cautiously practised is one of the most beneficial techniques in treatment. Many cases of infantile eczema have cleared up completely by being gradually acclimatized by exposure.

In addition to controlling irritation and restraining scratching by sedatives and antipruritic medicaments some method of mechanical restraint is often indicated. In small children loose cardboard cylinders may be applied to the

arms, or canvas bands may be fastened to a strip of canvas across the bed to immobilize the subject. Adults may wear thin cotton gloves and keep their nails very short, or their wrists may be tethered to the pyjama girdle. It is obviously impossible to clear up a lesion of the skin if it is actively scratched. The intelligent patient who can refrain from scratching while fully awake may be found to scratch violently during sleep, and will often admit to finding the lesions bleeding and excoriated on waking in the morning.

Occlusive dressings over tar or zinc paste are effective partly because of their barrier to scratch injury, and the old technique of painting the irritable area with crude coal tar was effective for the same reason. A 10% solution of crude coal tar in acetone is somewhat more elegant than plain tar. Of the proprietary preparations "tar derma-ment," which contains a varnish basis, is sometimes very useful. When the tarry film is tacky it may be reinforced by the application of a talcum powder, and "Fissan ichthyl powder" is usually well tolerated by the eczematous skin. "Dermatol" (bismuth subgallate) has a reputation for its astringent and sedative properties. Another protective preparation is 10% ichthammol in flexile collodion.

Psychological Considerations

No discussion of modern treatment is complete without a consideration of the patient's psychological reaction to his illness and of measures to diminish his anxiety and fore-sight in preparing him for his return to useful work.

Young patients should be provided with a variety of toys, and in severe cases of eczema they are undoubtedly better if nursed in a ward where they have playmates. Children unhappy and fretful with restraint in their cots, doing their best to scratch or rub, are often seen to play happily without obvious discomfort from their eczema when allowed to be up.

Occupational therapy cannot be introduced too early in the treatment of adults with eczema, and if interesting handicraft can be provided it is often possible to dispense with sedatives during the daytime. In milder cases of eczema it is advantageous for the patient to continue at work, provided that further contact with irritants is avoided, and the co-operation of the employer should, if possible, be obtained to secure this. If alternative clean light work cannot be found it would be better for the subject to seek some other job. Many patients with intractable chronic eczema get a fixed anxiety neurosis, and the patient's sole interest is his skin. In this phase the prognosis is almost hopeless, and it is believed that the tragedy and economic wastage can usually be averted by giving as much thought to the maintenance of the subject's morale and confidence in his or her ability to return to full employment as to the details of local treatment.

REFERENCE

Hadgraft, J. W., and Brain, R. T. (1949). *Lancet*, 2, 78.

A six-months course for training men and women who wish to take posts of responsibility in Old People's Homes will be held in September by the National Old People's Welfare Committee and the National Corporation for the Care of Old People. Students will be given lectures on institutional management, first aid, and home nursing, and practical training in a hospital geriatric unit and in an old people's home. Inquiries about the course should be addressed to the secretary, National Old People's Welfare Committee, 26, Bedford Square, London, W.C.1.

THE DEVELOPMENT OF THE ELECTRO-CARDIOGRAPH IN GREAT BRITAIN

BY

S. L. BARRON

Electrocardiographic research may be said to have begun in the latter end of last century with the work of A. D. Waller (1890), of Bayliss and Starling, and of Gotch and Burch, who, employing a capillary electrometer, established that a form of electrical potential curve was associated with the contraction of the human heart.

Einthoven's Researches

Willem Einthoven, of Leiden University, was prompted by these researches to make similar investigations, and published a paper in 1895 describing work also using the capillary electrometer. Realizing the limitation of this method, he devised a more simple, sensitive, and accurate means of recording these fluctuating electrical potentials. The instrument we know as the "string" galvanometer had been known in theory for some years, and had been applied and described by Ader in 1897 for investigations in cable telegraphy. It consisted of a single metal fibre stretched in a magnetic field, and was intended as a quick-period system rather than a sensitive one. Ewing in 1892 described a bifilar galvanometer containing two conductive fibres bridged across the centre by a small mirror and the whole inserted in a strong magnetic field. This was used for tracing magnetization curves and escaped notice as a reflecting mirror galvanometer. As both these papers were out of Einthoven's field he was unaware of them when he began to develop the string galvanometer.

The problem that confronted Einthoven was to make an instrument of high internal resistance, dead beat, with quick period, highly sensitive, and stable for accurate calibration. As sensitivity is normally obtained at a sacrifice of period and stability it was at that time a great problem. He experimented first with a highly sensitive D'Arsonval moving-coil galvanometer and found that the smaller the numbers of windings on the coil the greater was the normal sensitivity. This led first to a single-turn coil and then to stretching a single wire in a strong magnetic field. In order to get a high electric resistance in the wire or "string" as it became termed, he developed a method of chemically plating with silver fine quartz fibres made by the classic method devised by C. V. Boys (1887). These fibres were electrically conductive, of high resistance, and owing to their small mass had negligible moment of inertia; on them the whole of the development of electrocardiography depended, and they are still in common use to-day.

The fibres were tautly suspended between the poles of a powerful electromagnet, and the shadow of the "string" was projected by a large arc lamp through a specially designed optical system which photographed the movement of the string shadow on a moving photographic plate. The electromagnet was large and unwieldy, and was cooled by a water-jacket (see Fig. 1).

In 1903 Einthoven described his equipment in a historic paper, "Ein neues Galvanometer," quickly followed by a paper (Einthoven, 1904) which compared the earlier work of Waller, using a capillary electrometer, with the new string galvanometer, and established the latter as highly accurate and suitable for electrocardiographic work.

Einthoven (1908) followed by a long and painstaking research in which he established the forms of connexion to