of yesterday, and may be the discarded fallacy of to-morrow. Methods, like men, must establish their own grading.

Above all things it is important that the relation of confidence between the doctor and his patient should not be disturbed. The patient should look on his doctor as an individual whom he has chosen to help his own individual needs, as a man of power who can invoke State assistance and command State resources, never as an official at the end of an office string. For this relationship to be maintained it is essential that private practice should be encouraged rather than driven into the hands of the small group of charlatans who would still exist to prey on the stragglers of a society in which all reputable doctors were perforce full-time employees of the State. In matters of decisive concern to the individual—his health, his house, his legal affairs, his money—it is essential that each man should be free, if he so desires, to seek advice from an expert of acknowledged position and integrity, chosen by himself. Only when such a body of experts is available, as a standard for comparison and as a Court of Appeal in doubt or difficulty, can a Government service be wholly efficient or command complete confidence. It is clearly in the interests of a State medical service that the men who work within it, and the men whose wisdom or whose contributions to science have earned them a position in which they are sought as second opinions, should be the same.

At the Cross-roads

American medicine is at the cross-roads, and American doctors would be wise to face that fact. They can see, if they will look, other nations proceeding along, or being dragged along, some of the many paths that lead from the crossing. They are still free to choose their own. They can pick the good and reject the bad from other systems if they will study them, and from these pickings they have to-day an unparalleled opportunity, one that will never come again, to produce their own system and make it work. They can then go to their Government and say, "Here is a plan that we, the doctors of America, have devised and tested. We claim that it will keep our scientific research in the forefront of world progress, that it will allow every man who enters medicine to find that work to which he is most suited by inclination and ability, and that it will enable every citizen of our great country to get the treatment that his disability demands. We can prove that the cost of this system to the country is less than that of the National Health Service in Great Britain, that the benefits it confers are greater, and that it is more readily adaptable to changing needs. We ask you to take it over as a going concern. We will run it for you."

The organization of medical services must come, all over the world, and the best people to initiate and develop such a public service are the doctors themselves. They alone have a clear and consistent grasp of the true relation of those services to the people, which is essentially that of servant and master—meaning that the true centre of interest and action is not in the authority but in the patient. And that patient is not a number in an office, but a unique person, a living soul. It is because doctors, and especially that most responsible of doctors the general practitioner, can never forget that all-important fact, so easily mislaid in offices and ignored by Governments, that I hope the American profession itself will take the first steps in its own development on a modern nation-wide scale. Will it do this? Not if it goes on worrying over Granny.

NEW ORIENTATIONS IN EPILEPSY*

BY

DENIS WILLIAMS, M.D., D.Sc., F.R.C.P.

*Neurologist, St. George's Hospital; Assistant Physician,
National Hospital, Queen Square*

Epilepsy has always aroused the interest of physicians, and it continues to do so probably because of our ignorance of the mechanisms responsible for its maintenance. There are other reasons for the present interest in this disorder. First, its manifestations are dramatic and its origins are certainly obscure—it is arresting and has always been mysterious. Secondly, it presents a social and economic problem of some size, for its incidence is surprisingly high. Statistical studies in America, Switzerland, and Holland (Ledeboer, 1949) suggest that at least 1 in 200 people in each of these countries suffers from epilepsy in one form or another. If the figure for the British Isles is similar, there must be 200,000 subjects of epilepsy in our midst, and rough estimations have ranged from 100,000 upwards. Lennox (1947a) has estimated that there are about 10,000,000 epileptics in the world, although no more than about 300,000 receive treatment. Thirdly, as Hughlings Jackson's work showed, the disturbances of function occurring in the epileptic fit have helped towards the understanding of the localization and integration of cerebral activity. Lastly, in the past 15 years the electroencephalogram (E.E.G.) has shown more of what is taking place in the cerebral cortex during epileptic fits and between them. It has shown differences between epileptics and normals, and it has revealed new facts about the causal lesions. In doing this it has not only given us an instrument through which we can see more, but it has provided a sharp stimulus to the work and interest in epilepsy. Observations made with the E.E.G. have confirmed many of the deductions of Jackson and Gowers, but they have also given us new knowledge and so have modified our view of the clinical state of epilepsy. It may be profitable, therefore, to take stock, and to consider what reorientations have been brought about by these recent advances in knowledge.

The electroencephalograph itself is an impressive piece of apparatus; the electroencephalograms are tangible and convincing; the jargon which surrounds the subject, derived as it is from general physics, electronics, and physiology, creates an atmosphere in which the clinical worker feels bewildered and ignorantly respectful. He feels that this is something beyond the range of his education and experience, and he has not quite realized that most of the advances it has made possible have been in clinical knowledge; for the physiology of the E.E.G. is almost unknown, and the records, the result of activity in millions of cells, cannot yet be interpreted through the medium of axonal activity. The electroencephalographic changes found in epilepsy had to be related to careful clinical observation, and most of the interpretations of the E.E.G. made so far have been clinical.

But there are other reasons, mainly clinical, for a changed approach to epilepsy. One is the discovery of new anti-convulsants, such as the hydantoins, which are effective in one form of epilepsy and not in another, and so call for a differentiation of types. Another is the advance of neurological surgery, which requires detailed knowledge of the lesion causing epilepsy. Penfield's work has been largely responsible for this particular advance. A further reason

*An Open Lecture delivered in Edinburgh University on October 25, 1949.

is the realization that much abnormal behaviour—some psychotic, some criminal, and some only eccentric or anti-social—has a physical cause which is closely related to that of epilepsy. Interesting observations of this sort on people with abnormal behaviour, and particularly on psychopaths, have been made by Lindsley (1944) and Hill (1944).

The Electroencephalogram and Classifications

The E.E.G. has influenced knowledge in the field of epilepsy in four main ways: (1) types of electrical discharges have been related to clinically differing seizures; (2) discharges identical with those seen in an attack but of shorter duration have been seen without visible changes in behaviour (larval attacks); (3) focal origins of seizures have been demonstrated and studied; and (4) differences have been found between the E.E.G.s of epileptics and normals irrespective of the occurrence of a fit.

On the basis of these changes, which need not be described here, new classifications of the epilepsies have been attempted. These have been of two kinds: those based upon the form of electrical pattern occurring in the fits (Gibbs, Gibbs, and Lennox, 1943); and those based upon the site of origin as well as the kind of electrical discharges (Jasper and Kershman, 1941). Both the classifications have of course depended on clinical as well as electroencephalographic information and have involved the use of clinical nomenclature, so that some confusion has arisen. Without entering into any of the controversy that has centred around these classifications, their main implications may now be considered.

On the Basis of Electrical Patterns

Three main groups of epilepsies having this basis can be recognized: (1) grand mal, associated with runs of very fast spikes in the E.E.G.; (2) petit mal, associated with the three-a-second wave-and-spike discharge; (3) psychomotor epilepsy, which is associated with various wave patterns, which can, however, be grouped into those in which waves are linked to give a rhythmic complex wave form, those which Gibbs calls a slow or modified wave and spike, and those in which no characteristic linkage is evident.

Much has already been written about each of these types of epileptic discharges, but some general conclusions and also some tentative hypotheses can be based upon the new knowledge from which the classification has arisen.

Grand Mal

The first type of discharge with the very fast, spikes (usually about 22 a second) may be accompanied by motor or sensory disturbances. When the epileptic attack is motor it has as its characteristic result the simultaneous contraction of muscle groups. This contraction may be continuous (tonic) or intermittent (clonic); it also may be generalized and symmetrical, or focal and asymmetrical. But wherever the contraction occurs it has the same form. If, for instance, it is in the hand, both the muscles causing flexion and extension of the digits and also the synergic muscles contract simultaneously, so that the position the digit assumes during the contraction is determined by the relative strength of the opposing muscles.

There has been a simultaneous discharge of cortical neurones which has caused an unphysiological response with no regard for the relationship of agonistic, antagonistic, and synergic muscles, resulting in a movement that has no resemblance to the function of the part. It is an unorganized contraction. While this mass contraction is taking place the cortex is discharging high-voltage spikes,

which are of course simply fast waves. These fast waves are often generalized and symmetrical, but they may be limited to one area of the cortex. When the movements and discharge are unilateral the patient is usually conscious, but when the movements become bilateral consciousness is nearly always lost. If the attack is sensory its manifestations are again primitive and unorganized, the disturbance being described as tingling or pins-and-needles. There is nothing about its form to suggest integration in the cortex.

Psychomotor Epilepsy

The third type of discharge will be considered next because it has some points of affinity with the first. It is characterized in the E.E.G. by a rhythm made up of waves of one, two, three, or more frequencies linked together. This linkage is most often harmonic, but the patterns of waves which result are diverse not only because of the many relationships between different frequencies that might occur, but also because a wave of a given frequency does not necessarily repeat at its own frequency. Thus a wave with a period of one-sixth of a second may occur twice a second, linked with a wave having a period of half a second. Because of these linkages such patterns as "square waves," "sawtooth waves," and "waves and humps" have been described, but these obvious patterns are really the result of this kind of organized linkage of sine waves of different frequencies (Dawson and Walter, 1944). The rhythms making up these wave complexes are mostly in the band between 3 and 10 a second.

From clinical observation it seems to me that the motor expression of these disturbances is composed of an organized contraction of muscles. Agonists contract, antagonists relax, and synergists contribute to the movement. Thus, again taking the hand as an example, whereas in the tonic-clonic attack the finger may be in tonic spasm, in the "psychomotor" attack it flexes in a natural way. When other digits move in a similar manner the movement, which may be unconscious and is certainly not purposive, appears to be purposive, so that it may be described as "beckoning," or "rolling a piece of paper," or "clenching his fist," but the observer is really ascribing purpose to an unconscious purposeless act. The contractions, then, in this form of epilepsy are co-ordinated and physiological. Just as the first form of epilepsy may be motor as well as sensory, this form may be sensory also.

Whereas the counterpart of the unorganized contraction of muscles seen in tonic-clonic epilepsy is an unorganized sensory experience felt as tingling of part of the body with perhaps loss of sensation also, in the "psychomotor" attack sensory changes are organized and are much more like those felt by the normal person. He experiences fear, or anger, or a sense of pleasure, or a compulsive urge, or a mixture of the lot. The sensation he experiences need not be that of an affective change, but if it is a somatic sensation it relates to the whole body, in relation either to itself or to its environment. On the one hand, things seem too near or too far, minute, crowding in, or distorted, or the position of the body in space may be misrepresented; while, on the other hand, the sensation of the body itself has changed—it is too big, or the patient feels he is projected outside it.

Whatever the disturbance of feeling, whether it is of affect, proprioception, or exteroception, it is at least highly enough organized to be felt as a caricature of normal feeling. One of the characteristics of this sensation is that it is compounded of many feelings, and so cannot easily be described. Once again it is as false to attribute psychological content to this feeling as it is to ascribe purpose to its

accompanying movement. Although location will be discussed later it is important to mention here that these "psychomotor" disturbances seem to arise in the temporal lobe (Gibbs, Fuster, and Gibbs, 1948) in contrast to the tonic-clonic disturbances, which are usually frontal or parietal in their origin. The importance of the temporal site of origin in relation to the memory patterns which emerge in the attacks (Penfield, 1949) and the abnormal behaviour associated with the attacks will be described later.

Petit Mal

The other of the three groups is that of petit mal, and it is in this group that it has been most difficult to reconcile clinical and electroencephalographic classifications. Petit mal used to mean just what it indicated, a small attack, but the classical three-a-second wave-and-spike rhythm is seen in only a fraction of patients having brief epileptic attacks. Lennox (1947b) has found that the wave-and-spike rhythm is associated with three kinds of attacks: (1) true petit mal—brief and frequent periods of unconsciousness, seen most often in children; (2) akinetic attacks in which there is sudden falling, with muscular hypotonia; and (3) myoclonic twitches.

The first of these disorders was that which was originally observed by Gibbs, Davis, and Lennox (1935) to accompany the wave-and-spike discharge, and which led them to apply the term "petit mal" to this rhythm. But the difficulty soon arose that brief periods of interruption of consciousness were occasionally found to be accompanied by bursts of spikes (grand mal) or by complex wave forms (psychomotor), so that the wave-and-spike complex was not peculiar to petit mal in the clinical sense. Petit mal had in fact become split up into "true" petit mal, small grand mal, small psychomotor attacks, and combinations of the three. Another difficulty was that the pattern of the larval attack seen in the E.E.G. did not necessarily reflect the patient's clinical disability, so that, for example, a patient with rare major convulsions might show only larval petit mal (wave and spike) in the E.E.G. We know now that the pattern of the discharge seen in the E.E.G. between fits does not necessarily indicate the kind of clinical epilepsy from which the patient suffers. It merely shows he has epilepsy, but it may also show that the epilepsy is of a mixed form. This distinction of the different forms of petit mal has led to a therapeutic advance, for the clinician may now recognize the nature of the patient's attack, however brief, and then plan specific anticonvulsant treatment accordingly.

There is no difference at all between pyknolepsy and true petit mal—clinical, electrophysiological, or therapeutic—and the use of the term should cease. Lennox (1947b) has tried to overcome the confusion of terms by using the word pykno-epilepsy for frequent stereotyped brief interruptions of consciousness, usually occurring in childhood and associated with a three-a-second wave-and-spike discharge in the E.E.G. Although the experienced neurologist can usually distinguish this condition from others by careful case-taking, the term "petit mal" will continue in use for brief attacks of any sort; and indeed it should, for that was its original meaning. It seems to me, therefore, that it would be less confusing to retain the phrase "petit mal" and to refer to this special well-recognized subgroup as "juvenile petit mal."

Juvenile Petit Mal

The three-a-second wave-and-spike phenomenon associated with juvenile petit mal has many features by which it differs from other forms of epileptic discharges. Its distribution and spread over the hemispheres is unusual, for often this is not only generalized and symmetrical, but each

phase of the wave complex appears synchronized in all positions—a fact difficult to explain on the basis of known mechanisms. There is also a most interesting variability in the rate and mode of spread of the rhythm over the scalp, and study of this spread, which has not yet been made, might be most profitable. Because of these features workers have felt that the rhythm arises in the basal ganglia, and recent experimental work on animals, particularly that of Lewy and Gammon (1940) and Jasper and Droogleever-Fortuyn (1946), supports the view that the optic thalamus is closely concerned with the cortex in the initiation of the disturbance.

With Dr. Parsons-Smith I have been making simultaneous records from the thalamus, from the overlying white matter, from the cerebral cortex, and from different points on the scalp. The results, which are not yet published, show plainly that a thalamo-cortical mechanism is used in man for the maintenance of this peculiar rhythm of electrical potential change. This intimate role of the thalamus in petit mal naturally leads to speculation, for this is the only occasion in which structures other than the cortex have been observed to be responsible for the actual discharge of epilepsy. This fact is probably the basis of all the fundamental ways in which juvenile petit mal differs from other forms of epilepsy. May it not be, for instance, a disturbance in the thalamus—the head ganglion of the afferent system—that leads to the inactivity, hypotonia, and amnesia of petit mal? If during the attack the hemispheres are deprived of the normal flow of afferent impulses all responses to environmental change will cease, spontaneous voluntary movement will not occur, and amnesia must result. Petit mal therefore is epilepsy disturbing the afferent (receptive) side of the cerebral mechanism. If this is simply speculation it is certain that petit mal differs entirely from other forms of epilepsy in its clinical manifestations, course, and response to treatment, as well as in its E.E.G. patterns.

A final attribute of the three-a-second wave-and-spike outburst of petit mal that has been revealed by the E.E.G. is that it does not arise as a result of acquired lesions. This must, however, be qualified by stating that it is impossible to say whether the wave-and-spike discharge can be induced by cerebral disorder occurring during development *in utero*, at birth, or very shortly after. In a personal series of more than 20,000 E.E.G.s, wave-and-spike outbursts have not been seen to follow brain damage caused after the first few months of life. In only three instances have they been seen in patients not known to have epileptic attacks, and one of those three was the identical twin of an epileptic. In consequence they are taken as certain evidence of constitutional or idiopathic epilepsy, for no real distinction can exist between epilepsy caused by unrecognized brain disorder arising during development and that arising without any evident acquired cause.

On the Basis of the Origin of Epileptic Discharges

Jasper and Kershman (1941) first emphasized the importance of epileptic foci, although these had been recognized in the E.E.G. in 1936 by Walter; and indeed Hughlings Jackson discussed their clinical evidences half a century ago. Penfield has recently presented the evidences of many operations in which epileptic foci have been located by direct cortical recording; he has then reproduced the features of the seizure by electrical stimulation of the site, and has then abolished the focus and the fits by cortical excision. When this process of investigation has involved simple focal fits, whether sensory or motor, it has served to confirm and amplify clinical opinion, but when the epileptic disorder has included disturbances of affect, abnormal and apparently

purposive behaviour, or disorders of thought, the investigations have obviously added greatly to our knowledge of higher cerebral function. It is not too much to say that an era of cerebral surgery controlled by electrical recording and stimulation is beginning, and that elaborations of this technique will replace the cruder "psychosurgery" of to-day. Of great interest in this respect is the observation of Gibbs, Fuster, and Gibbs (1948) that the focus of origin of the discharges characteristic of "psychomotor" epilepsy is usually in the anterior part of the temporal lobe, particularly on the left, for foci of abnormality are also found in this position in patients with behaviour disorders.

When due attention has been paid to the focal onset of epileptic discharges, and in particular to the remarkable experimental surgery of the Montreal Institute, it is still the widespread distribution of the epileptic activity in the hemispheres which seems all-important. A study of the results of head injuries in the second world war (Williams, 1941a, 1941b) allows some general statements upon the development of epileptic changes as a result of brain injury. It is of course true that many local brain injuries give rise to a focus of epileptic activity the position of which coincides with the site of injury. This follows clinical experience, but what seems to be of greater importance is that more often a severe focal brain injury gives rise to episodic changes in the E.E.G. which are typical of epilepsy, and which are seen equally well in both hemispheres at points remote from the site of injury. For instance, a penetrating injury in the posterior part of one hemisphere may be followed by bursts of high-voltage 22- to 28-a-second waves arising in both frontal poles, although at the site of injury the kind of electrical disturbance may simply be that of organic brain damage apparently uncomplicated by secondary epileptic changes. Something must have happened to the whole brain in most of these cases, and in many no focal initiation could be found at the site of injury. A primary change at the site of injury has been followed by secondary changes occurring symmetrically in the two hemispheres. A further important fact is that an interval of time must elapse before these secondary changes occur.

In serial records made at intervals after an injury, during which we observed the development of secondary episodic discharges, and also when studying the records made in many cases at varying intervals after injury, we did not find this interval to be less than eight weeks, but it was more usually three to six months. Cases were collected in which these disturbances preceded the first fit; in many others in which they were found no fits have yet occurred; and in others, again, traumatic epilepsy began without any discharges of this sort. Indeed, the E.E.G. may be normal in at least 25% of traumatic epileptics.

We were forced to conclude from this evidence that after the primary local disturbances had occurred secondary changes developed which were delayed and which were often symmetrical or generalized in the hemispheres. It seemed that some change had taken place with the passage of time which had created an epileptic brain, rather than that the whole disturbance was due to a local epileptogenic scar. In support of this view there is no doubt that an inherited factor is potent in the development of traumatic epilepsy. Although removal of the damaged brain at the time of injury greatly reduced the likelihood of secondary changes occurring in the E.E.G., subsequent removal of the scar was not so effective in relieving established epilepsy as seen in the E.E.G. It seemed, indeed, that in some cases once the secondary episodic disturbances had transcended the area of injury they were independent of removal of that area. Clearly, as Penfield and his colleagues have

shown, the local scar causes epilepsy, but general factors are at work in making both the injured area epileptogenic and the whole brain epileptic.

The Epileptic Fit

Form

The features of the clinical attacks associated with the three types of epileptic outbursts seen in the E.E.G. have already been contrasted, but it would also be well to consider the relationship of the fit itself to the persistent disorder which renders the patient epileptic. Before the E.E.G. was used, except in those cases in which a causal lesion gave a continuous disturbance of function, there was little in the patient's story or behaviour to link one fit with the next. Epilepsy was a series of isolated episodes. In some cases this is still so, but in many the E.E.G. shows changes in cerebral electrical behaviour which may be continuous. There is some relationship between the kind of continuous disorder and the kind of fit, so that the sort of E.E.G. records shown by a patient with grand mal is usually quite different from that of a patient who has only petit mal. Except that petit mal is a manifestation of epilepsy beginning in childhood—perhaps related to incomplete myelination—and so tends to appear whenever the cause of epilepsy arises early in development, the kind of epileptic fit in any subject depends upon qualities in that subject as well as upon the kind of local cause.

This was well shown during the war, when we were engaged in research into the oxygen convulsions occurring in deep divers. In a pressure tank one of a team of volunteers breathed pure oxygen under high pressure (three to four atmospheres) until a convulsion occurred. The E.E.G.s and other recordings of body function were made continuously. Although the usual result was a generalized tonic-clonic fit with typical bursts of high-voltage spikes arising in both parietal and frontal regions and increasing in voltage until a convulsion occurred, in some subjects a different pattern of disturbance took place. Rhythmic wave complexes made up of frequencies in the range of three to seven a second began, and a clinical attack with restless, apparently purposive, and even aggressive behaviour followed. The subject had had a "psychomotor" attack in circumstances that produced grand mal in others, but, as is the case in epilepsy arising from other causes, grand mal was more often seen than other variants of epilepsy. Here it may be stated that all fits are epileptic, whether they are symptomatic of brain damage, whether they are associated with the toxæmias of pregnancy, with uraemia, with anaesthetics, or whether they are produced therapeutically. The actual physiology of the fits themselves is similar in all conditions.

Frequency

Many factors are known to influence the rate of occurrence of fits—the kind of attack, the natural rhythm with which fits occur, the age of the patient, his emotional state, the special precipitants, and so on—but these are so well recognized that they need not be re-examined here. There is little relationship between the number of actual clinical attacks and "larval" attacks which are seen as discharges in the E.E.G. but which have no corresponding clinical counterpart.

One patient will show larval attacks every few minutes, and have one or two clinical fits a year, whilst another will never show any disturbance in the E.E.G. except during clinical convulsions. It seems as if in the first instance in which many epileptic disturbances begin but few reach maturity there may be a high resistance to the discharges,

compared with a low resistance in the instance of the patient who has a major fit on the few occasions in which an epileptic disturbance arises in the brain. The first may well be considered to have severe epilepsy with a high resistance, the second mild epilepsy with none. This suggests a dual mechanism in the determination of epileptic attacks, consisting of epileptic disturbances which invade the brain and a natural antagonism to these disturbances. Such a conception might help to explain some of the anomalies in the genetics of epilepsy and in the onset of symptomatic epilepsy.

Site of Origin

The original clinical observations upon the site of epileptogenic lesions were concerned with the primary centres of cortical representation of movement and sensation, for lesions of these areas produced obvious evidences of stimulation or inhibition. Thus the Jacksonian motor fit was first described and was related to the structure of the motor cortex, and understanding of other special areas of cortical representation soon followed. The frontal and temporal lobes offered the least constant and most obscure patterns of epileptic activity. The electroencephalogram has shown that epileptic foci occur more commonly in the frontal lobes—and especially in their poles—than in any other area, and through it localization has become possible where the fits have presented no localizing clinical features.

Complex disturbances of affect can be related to foci in the temporal lobe, and disturbances of visceral activity to the lower part of the frontal lobe. This sort of activity was shown in a boy aged 11 who had had stereotyped attacks for four years, consisting of a painful gripping feeling in the epigastrium with a hungry sensation, followed by very active borborygmi and the passage of flatus. This chain of events was sometimes followed by clonic twitching of the right arm. The E.E.G. showed a well-defined focus of epileptic activity in the left fronto-temporal region, and a ventriculogram a porencephalic cyst overlying the epileptic focus.

Complex disturbances of affect were described by a graduate of 22 who had an epileptic focus above the left ear, over the middle of the temporal lobe. He had attacks heralded by a jumble of recognized voices giving him commands which he could not remember afterwards. He was bilingual, and they were first heard in the language of his home country, but later when he came to England they spoke in English. The voices grew louder and more insistent, and he then had a most unpleasant feeling that was compounded of an unnatural fear, a sense of anger and of frustration, "and a mixture of feelings that I can't describe."

Disturbances of this kind can be accurately located by the E.E.G. Penfield's experimental studies upon the stimulation of the temporal region have evoked sensations and memories in the epileptic subject which were identical with those experienced in the epileptic attack. They could also be evoked in identical form by repeated stimulation. The importance of work of this sort, in which the patient can accurately describe the effects of local stimulation, needs no comment.

Spread

The grand-mal attack and petit mal may spread so quickly over the hemispheres that the discharge in the E.E.G. may appear synchronous in different areas, and the clinical manifestations occur simultaneously throughout the body. Yet other epileptic disturbances spread through the hemispheres very slowly indeed, and others do not spread at all. Twitching may continue for hours

at a corner of the mouth or in a digit without any spread (epilepsia partialis continua), and a Jacksonian fit may slowly spread from finger to finger and so to the thumb, hand, wrist, and up the arm, several minutes being taken in the process. Such a slow spread of the accompanying electrical activity in an attack can often be seen in the E.E.G.

It seems to me that this limitation of the spread of an epileptic discharge is of importance in the understanding of epilepsy. In one case the high-voltage discharges of epilepsy spread over the whole brain unimpeded, in others they are localized or are so inhibited that they spread very slowly over a limited area. In yet another group of patients generalized epileptic discharges are associated only with local movements. Once again there must surely be two factors at work—the epileptic disturbance and a factor opposing it, perhaps the same factor that prevents larval attacks from disturbing consciousness or behaviour.

The Constitutional Factor in Epilepsy

Conrad (1936) found that when idiopathic epilepsy occurred in a twin it affected the other twin in 86% if they were uniovular, and in only 4% if binovular. Lennox, Gibbs, and Gibbs (1942) found that 87% of uniovular twins had identical E.E.G.s, and later that if epilepsy was present in one twin that twin had evidence of an acquired cerebral lesion and the non-epileptic had an abnormal E.E.G. Lennox, Gibbs, and Gibbs (1940) found that 54% of the first-degree relatives of epileptics had abnormal E.E.G.s compared with 6% of a normal control group. They also found that in 94% of a group of epileptics one or more of the parents had abnormalities in the E.E.G. In contrast to this figure is the clinical fact that in only a relatively small proportion of epileptics is there a family history of epilepsy. The figures for this range from 20 to 33%, depending on the group studied. Considering the heredity of epilepsy as seen in the occurrence of fits and in the appearance of abnormality in the E.E.G., it seems that the likelihood of the predisposing cause of epilepsy being inherited is very great; that actual epilepsy appearing in these predisposed persons is not nearly so likely (about 1 in 4); and that a hereditary constitutional factor predisposes subjects to symptomatic epilepsy.

This last point is further supported by observations on eclampsia (Rosenbaum and Maltby, 1943) and on anaesthetic convulsions (Williams and Sweet, 1944). These authors concluded that the occurrence of fits in the toxæmias of pregnancy and during anaesthesia was to a large extent determined by constitutional factors.

In the present search for epileptic foci and in the physiological study of epileptic processes occurring in the brain there is a great tendency to forget the importance of the constitutional factor. This factor does not only predispose the person to epilepsy but helps to determine the form the epilepsy will take. It is potent in predisposing to symptomatic epilepsy whether due to injury, infection, or tumour, as well as in initiating idiopathic epilepsy.

A convulsion is the end-result, in certain circumstances, of cerebral irritation. Its form is not determined by the nature of the irritant, and the likelihood of its occurrence is modified by constitutional factors. It has always been assumed that these factors have simply consisted in a tendency to epileptic attacks, but from time to time in this paper scattered evidence has been presented which raises the possibility that another factor—the failure to prevent the spread of the epileptic state—is also present. A dual inheritance of the state of epilepsy (the epileptic factor)

and the limiting or anti-epileptic factor would offer a convenient explanation of some of the phenomena which have been described.

Abnormal Behaviour in Epilepsy

Abnormal behaviour may occur in epilepsy as a direct result of the fit or irrespective of it. There are indeed four factors which will give rise to the associated abnormalities of thought or behaviour: (1) the fit; (2) the causal disease; (3) the treatment; and (4) a related neurosis.

There is now much evidence, and some acceptance of it, that "the epileptic temperament" does not exist. Alienists dealing with psychotic or deteriorated epileptics in institutions described aberrations in thought and personality which seemed peculiar to the epileptics among the patients under their care. But most of the epileptics were under care because they were otherwise abnormal, while the environment in which they had in consequence to live was very different from that of the normal population. Observation of epileptics who are living in the normal population, earning their living, bringing up their families, and enjoying normal leisure makes it plain that the range of their temperaments is as wide as that of others.

Meyers and Brecher (1941), after a critical psychometric analysis, concluded that there was no significant difference in personality between a group of epileptics and a group of normal people of similar background and age. The same conclusion, based on the result of the Rorschach test, was made on epileptic children by Kogan (1947). Furthermore, Lennox found that the average intelligence quotient in 300 of his own epileptic patients was 104. Epileptic patients would benefit greatly if there was a general realization of the fallacy of applying information from the examination of groups of epileptics in institutions to the great majority of epileptics who are living in their homes and trying to live a normal life.

With the confident statement that there is no disorder of personality or intellect characteristic of epilepsy *per se*, some of the disturbances of thought and behaviour associated with epilepsy will now be considered.

1. Behaviour Associated With the Fit

Abnormal behaviour may be ictal, post-ictal, or replacing. This division has been recognized clinically for a long time, but with the E.E.G. the exact limits of the ictal disturbance can be observed, and it now seems possible to make the distinction clearer on clinical grounds.

The post-ictal behaviour arises from the confusion of a clouded consciousness, and so is fundamentally negative in its character. The patient carries out simple acts in a routine way. He responds to stimuli in a primitive manner, and he shows a gross defect of judgment, understanding, and initiative when faced with more complicated situations. The ictal activity, on the other hand, is the positive result of abnormal cerebral activity. It consequently tends to be aggressive, apparently purposive, and often bizarre and completely illogical. It bears little relation to environmental change and none to events which immediately preceded the attack. Disturbances of thought, affect, and sensation also have this positive quality, so that the patient may be able to describe experiences which are quite outside the range of his normal experience when the attack is over. He will not be able to describe similar feelings if the event is post-ictal, since his amnesia will be complete or nearly complete.

Disturbances which replace the attack may be seen during treatment. They are uncommon, but always seem to consist of an affective disorder in which depression predominates. With the depression the accompanying irritability may lead to aggression and truculence, presumably due to a lowering in the power of judgment.

It is relevant here to mention the temporary change in intellect and personality which may accompany frequent petit-mal attacks. It is quite common to find a sudden and striking improvement in intellectual performance as judged by school work in a child whose petit-mal attacks have successfully been controlled by "tridione." In such circumstances a girl of 12 who had descended from top place in her form to the bottom reached the top third of the form within six months of first taking tridione. The improvement was parallel with the reduction in attacks. As in this case, petit mal may be so frequent that the rapidly repeated interruptions in consciousness will make sustained attention or concentration impossible. The state then simulates mental defect.

Of less common occurrence is the periodic change in behaviour and mood that may accompany an episode of very frequent petit-mal attacks. A parent may complain that an otherwise well-behaved reasonable child will have "bad days." At those times he may be surly and stupid, unreasonable and negativistic. Careful observation may show brief periods of unconsciousness of his surroundings followed by confused, irrational, and uncooperative behaviour.

Another form of episodic abnormal behaviour is that seen in those cases in which disturbances of affect are epileptic in origin. Here the affective disorder is of sudden onset, is brief, and is stereotyped. It has no emotional precipitant, but it comes out of a clear sky. Examples of these disturbances were seen in the following cases.

A nurse had sudden unexplained momentary feelings of compulsion, in which she performed a single aggressive act. She was psychiatrically normal between the episodes, which were common before the menses, and which ceased completely on taking phenytoin sodium alone.

A schoolboy aged 13, known to have convulsions, had rare periods of intense, unreasonable, and as he said "unnatural" fear without cause. They had a sudden onset and ending, and lasted for an hour or so. In one he wrote a suicide letter for which he had amnesia. The episodes stopped dramatically on taking phenytoin sodium and returned when the drug was temporarily discontinued.

2. Causal Disease

The view is widely held that epilepsy leads to intellectual deterioration, and it used to be said that petit mal most rapidly caused this deterioration. There is abundant evidence that neither of these views is true. A number of psychometric studies, of which that of Sands and Price (1947) may be taken as an example, show that the intelligence of ambulant epileptics is the same as that of corresponding groups of the non-epileptic population provided there is no gross organic cause for the epilepsy. The intelligence of patients with symptomatic epilepsy, however, tends to be lower than that in the normal population, the lowering of level depending upon the kind of lesion and not on the epilepsy. The fallacy that petit mal causes deterioration more rapidly than grand mal has probably arisen because petit mal can be symptomatic only if the lesion is developmental or if it occurs very early in life. The fact is that a group of children having congenital cerebral lesions of necessity contains a large number of mental defectives, many of whom also have epilepsy, particularly petit mal. The position may be expressed in the statement that when intellectual defect is present in the presence of epilepsy it is usually the result of the cerebral disorder which has caused the epilepsy. Petit mal alone will not cause deterioration, and grand mal does not usually do so, but histological study suggests that if major convulsions are numerous enough they may lead to cumulative brain damage and progressive deterioration. The occasional fits seen in the socially adapted epileptic do not seem to cause significant deterioration. Such extensive histological studies have been made of epileptic brains that, had there been a constant basic pathological process resulting from epilepsy, discussion would not still be centring on such problems as the significance of the degenerative changes in the cornu ammonis. There seems to be a spreading view that many of the histological changes attributed in the past to the epileptic process were in fact evidences of the lesions giving rise to the epilepsy.

3. Treatment

There are three ways in which anticonvulsant therapy may cause psychiatric changes. Bromism is now uncommon, because of the replacement of bromides by other anticonvulsants, and because of the realization of the effects of prolonged administration of bromides. It is not so widely realized how many days may elapse after beginning its use before an effective level of blood bromide is reached. Giving 30 gr. (2 g.) daily, an adequate level may not be reached for 10–14 days, and the excretion is much slower, so that in a personal case it took five weeks for the bromide level to fall from 180 to 85 mg. per 100 ml.

Intellectual deterioration and retardation, which is always temporary, follows the administration of barbiturates, including phenobarbitone in amounts which are too high for the individual, and there are surprising variations in individual tolerance. A patient and his relatives will describe marked drowsiness and retardation on $\frac{1}{2}$ gr. (32 mg.) twice a day, while another patient of the same age and weight is active and alert on 3–4 gr. (0.2–0.26 g.) a day. Recovery rapidly follows withdrawal.

The hydantoins may cause abnormal behaviour of an impulsive and aggressive kind, and should always be suspect if a sudden marked change in personality has been noticed, particularly in children. The child may become hostile to its parents, resenting discipline, having temper tantrums, and becoming personally aggressive. It may be necessary on rare occasions to withdraw the drug, but more often simultaneous administration of phenobarbitone will relieve the condition.

4. Related Neurosis

Lastly, a neurosis may result from the onset of epilepsy with its consequent limitation of social activity, introspective anxiety, and the retarding effect of therapy. The development of an anxiety state may in turn aggravate the occurrence of attacks, which will increase the neurosis, and such a change in temperament may occur that an organic cause may be suspected. The state should easily be recognized by anyone aware of the difficulties faced by the epileptic patient.

Borderline between Epilepsy and Psychopathy

There is a similarity between the aggressive asocial behaviour of the psychopath and the disturbances seen in psychomotor epilepsy. The epileptic disturbances have three characteristics which will normally distinguish them from the psychopathic—they are episodic, they are associated with amnesia or clouding of consciousness, and they have no apparent relation to preceding events. There are, however, circumstances in which the differentiation of aggressive psychopathic behaviour from epilepsy is difficult. The number of cases in which this is so can be greatly reduced by attention to the details of a careful history.

Because of the similarity between psychomotor epilepsy and aggressive psychopathy there has been a most unfortunate tendency to confuse the two conditions, and through indecision or loose thinking abnormal behaviour has sometimes been called "epileptoid." This tendency has been fostered by ignorance of the constitutional basis of the disorder or by sheer force of circumstance. Impulsive crimes or acts of violence committed without adequate reason have for the sake of the culprit to be explained, if possible in a way that will circumvent the M'Naghten rules. Thus the suspicion of epilepsy is raised, and if there is little else to support that suspicion the opinion grows that, although the crime was not committed during an epileptic state, it was rather like it—epileptoid. There are so few cases in which a sharp distinction cannot be made between epilepsy and other causes of abnormal behaviour that such a compromise is inexact and undesirable.

In making a distinction of this kind on clinical grounds the misconception must not arise that other forms of abnormal behaviour apart from that caused by epilepsy have

not a physical or constitutional basis. Hill has shown that a high proportion of aggressive psychopaths have widespread and severe abnormalities in the E.E.G., and he has drawn attention to the similarity of some of these abnormalities to those of epilepsy. Furthermore, Stafford-Clark and Taylor (1949) have found a relationship between the occurrence of abnormal E.E.G.s and the lack of reasonable purpose in major crimes, particularly murder. Lastly, the association of temporal foci of abnormality in both psychomotor epilepsy and in subjects with disturbances of temperament has already been mentioned in this paper. As more becomes known about the physiology of abnormal behaviour the borderline of epilepsy may appear more and more indistinct, but we must be careful not to confuse the identity of causes with that of the results. In discussing the significance of an abnormal E.E.G. in the absence of a gross lesion (Williams, 1941c) the statement was made that it "indicates an inborn constitutional defect which may find expression in epilepsy or a behaviour disorder such as a neurosis, psychopathy, or psychosis, in the subject himself or his offspring." This appears to be the case, for identical changes in the E.E.G. are seen in diverse states of abnormal behaviour or feeling.

Therapy

The advance in treatment from the use of anticonvulsants which are also sedatives—first the bromides and then the barbiturates (phenobarbitone)—to the hydantoins and trimethyloxazolidine has presented new problems. The hydantoins (e.g., sodium diphenylhydantoinate ("dilanitin"), methyl phenyl-ethyl hydantoin, "mesontoin") have a pronounced effect upon tonic-clonic attacks, a much less beneficial effect on the psychomotor attacks, and are of no benefit in petit mal. Tridione, on the other hand, is specific for those forms of epilepsy accompanied by wave-and-spike discharges in the E.E.G.—petit mal. Because of this specificity it is now essential to make an exact diagnosis of the type of epileptic disturbance experienced by the patient and seen in the E.E.G.; for a "small grand mal" with fast waves may be made worse by tridione, whereas juvenile petit mal with wave-and-spike discharge should be greatly improved.

The mode of action of the hydantoins in grand mal is of interest. They have no sedative or depressant effect; they are most effective as anticonvulsants in a dose only slightly below the toxic level; and, although they are most effective in preventing convulsions, there is a tendency, should an attack occur, for it to be a severe one, and even for it to continue into status epilepticus. There is also a tendency in some children towards hyperkinetic aggressive behaviour. Because of these effects the hydantoins are usually administered with phenobarbitone. In all these respects their action differs from that of the bromides or phenobarbitone. The difference may be summed up by saying that the sedative anticonvulsants protect the brain from the epileptic disturbance, while the hydantoins directly inhibit the epileptic process without depressing cerebral activity. Thus when a convulsion occurs it does so in an unprotected cerebrum which responds maximally. This difference in the pharmacology of these groups of substances supports the view already expressed that a dual mechanism exists in epilepsy—the epileptic state, and the degree of resistance of the brain to that state.

Tridione, which inhibits petit mal, is a convulsant in animals, and has no sedative action. There is no explanation for its dramatic effect in petit mal, but like some of the other drugs effective in petit mal—for example,

amphetamine and caffeine—its action is not sedative but rather the reverse.

The two important advances in medical therapy, therefore, are that it is no longer necessary to procure sedation to achieve control of epilepsy, and that specificity of response is being achieved in the control of the different forms of epilepsy.

Conclusion

In this review I have not meant to record facts about epilepsy, old or new, but have tried to show where new knowledge seems to be leading us. Lest I seem to have integrated clinical and electroencephalographic evidence without details of the supporting data, I would urge that the E.E.G. has presented further signs, which have been used in this paper with all the other evidence to reach conclusions through the clinical method of deduction. This clinical method uses the history, the physical examination, and the results of special investigations, of which the E.E.G. is simply one. This is not belittling the E.E.G., for it has presented us with an objective method of studying cerebral activity which will continue of the first importance to the study of epilepsy.

REFERENCES

- Conrad, K. (1936). *Dtschr. Z. Nervenheilk.*, **139**, 76.
 Dawson, G. D., and Walter, W. G. (1944). *J. Neurol. Neurosurg. Psychiat.*, **7**, 119.
 Gibbs, E. L., Fuster, B., and Gibbs, F. A. (1948). *Arch. Neurol. Psychiat., Chicago*, **60**, 95.
 Gibbs, F. A., Davis, H., and Lennox, W. G. (1935). *Ibid.*, **34**, 1133.
 — Gibbs, E. L., and Lennox, W. G. (1943). *Ibid.*, **50**, 111.
 Hill, D. (1944). *Proc. R. Soc. Med.*, **37**, 317.
 Jasper, H. H., and Droogleever-Fortuyn, J. (1947). "Epilepsy." *Ass. Res. nerv. ment. Dis., Proc.*, **26**, 272.
 — and Kershman, J. (1941). *Arch. Neurol. Psychiat., Chicago*, **45**, 903.
 Kogan, K. L. (1947). "Epilepsy." *Ass. Res. nerv. ment. Dis., Proc.*, **26**, 616.
 Ledebor, B. C. (1949). *Folia Psychiat. Neurol. Neurochir. Neerland.*, **52**, 1.
 Lennox, W. G. (1947a). "Epilepsy." *Ass. Res. nerv. ment. Dis., Proc.*, **26**, xvii.
 — (1947b). *J. Amer. med. Ass.*, **134**, 138.
 — Gibbs, E. L., and Gibbs, F. A. (1940). *Arch. Neurol. Psychiat., Chicago*, **44**, 1155.
 — (1942). *Ibid.*, **47**, 702.
 Lewy, F. H., and Gammon, G. D. (1940). *J. Neurophysiol.*, **3**, 388.
 Lindsay, D. B. (1944). In Chapter 33 of *Personality and Behaviour Disorders*, edited by J. McV. Hunt. New York.
 Meyers, R., and Brecher, S. (1941). *J. abnorm. soc. Psychol.*, **36**, 413.
 Penfield, W. (1949). *Proc. int. Congr. Neurol. Paris*.
 — and Erickson, T. C. (1941). *Epilepsy and Cerebral Localization*. Springfield, Ill.
 Rosenbaum, M., and Maltby, G. L. (1943). *Arch. Neurol. Psychiat., Chicago*, **49**, 204.
 Sands, H., and Price, J. C. (1947). "Epilepsy." *Ass. Res. nerv. ment. Dis., Proc.*, **26**, 604.
 Stafford-Clark, D., and Taylor, F. H. (1949). *J. Neurol. Neurosurg. Psychiat.*, **12**, 325.
 Williams, D. (1941a). *J. Neurol. Psychiat., Lond.*, **4**, 107.
 — (1941b). *Ibid.*, **4**, 144.
 — (1941c). *Ibid.*, **4**, 257.
 — and Sweet, W. H. (1944). *Lancet*, **2**, 430.

IMPOTENCE*

BY

S. LEONARD SIMPSON, M.A., M.D., F.R.C.P.

Physician (with charge of Endocrine and Diabetic Clinics), Willesden General Hospital; Endocrinologist, Princess Louise Children's Hospital; Consultant Endocrinologist, St. Mary's Hospital

Potency, or implied male sexual potency, may be defined as the ability to initiate, sustain, and successfully conclude the act of sexual intercourse to the satisfaction of the male. If we include the words "and the female" some 50% of potent males might be classified as impotent. In the rat it is easier to measure potency mathematically. Thus in male rats (Stone, 1932) previously separated from females for a few days the number of copulations an hour in the first few hours of reunion was 33, 16, 12, and 18. These rats were said to have a powerful sexual drive. Less virile rats had 9, 3, 5, and 2 copulations an hour.

In man, Davis (1929) has endeavoured to get some data by questioning 1,000 women college graduates about the frequency of intercourse in the early years of married life. The figures were:

Twice or more a night	..	..	..	2%
Once a night	..	..	..	8%
Twice a week	..	..	..	33%
Once a week	..	..	..	45%
Once a month	..	..	..	12%

Average replies showed that intercourse took place twice nightly in the first three months; once nightly in the first year; and twice weekly in the first three years. After the first ten years the average male is less sexually active, and Davis noted a wane in the sexual drive after the age of 30.

Animal experiments suggest that the anterior pituitary gland is responsible for the sexual drive but that the gonad is essential for its completion. Castrated rats may continue to copulate for some months after castration. Of 27 castrated criminals (Stone, 1932) 14 became impotent, 12 had reduced potency, and 1 was unchanged. Some eunuchs and eunuchoids may copulate in some degree, and prove unreliable "guardians of the couch." The secretion of androgens by the testis seems to be the essential factor for full potency. Although androgens are secreted by the adrenal cortex in normal males and in eunuchoids, and in the latter to a level comparable to the lower total values in normals, most eunuchoids are impotent, so that adrenal androgens might seem to be less effective than testicular androgens. Nevertheless the sexual precocity of adrenal tumours is a very virile phenomenon.

The secretion of androgens at different phases of life is therefore very relevant to our subject. Hamburger (1948), measuring androgens in the urine as 17-ketosteroids, found that values rise abruptly at puberty, reach a maximum at the age of 25, and then fall gradually from 25 to 50 and more abruptly from 50 to 70. Women also show some fall in urinary 17-ketosteroids after the age of 25, so that the adrenals also come into the picture. However, some men at 60 may secrete the same quantities of androgens as others at 25. Hamilton and Hamilton (1948) found a progressive decline of 17-ketosteroids from the third decade onwards, the seventh decade giving values 40% of those of the third decade. The alpha and beta fractions both decreased with age.

*A paper read to the Medical Society of London on November 28, 1949.

Red spoons for measuring medicine are recommended by the British Standards Institution. In discussion with the Pharmaceutical Society and the Ministry of Health, the Institution has been considering the question of variations in the size of teaspoons and tablespoons. Different-sized spoons for cookery and for medicine are now proposed. The American fluid ounce is larger than the Imperial fluid ounce, so, to allow American and British recipes to be interchanged without confusion, it has been thought essential to specify American standards for spoons used in cookery. It is therefore suggested that spoons for medicines should be smaller than the others. The new size proposed for medicine tablespoons is one-fortieth and for cookery tablespoons one thirty-second of an imperial pint; for medicine teaspoons, one one-hundredth-and-sixtieth, and for cookery teaspoons one ninety-sixth of an imperial pint.