CARCINOMA OF ISLETS OF LANGERHANS WITH HYPOGLYCAEMIA

BY

B. F. BREARLEY, M.B., B.Ch., M.R.C.P.*Senior Medical Registrar, Royal Infirmary, Sheffield*

AND

J. W. LAWS, M.B., Ch.B.*Medical Registrar, Royal Infirmary, Sheffield*

[WITH SPECIAL PLATE]

Banting and Best isolated insulin in 1921, and Harris formulated his concept of hyperinsulinism in 1924. Since then insulin-secreting tumours have been increasingly recognized. The great majority of these are adenomata, often small and multiple. Frantz (1944), working with Whipple, was able to collect 149 cases of islet-cell tumour with hyperinsulinism: 28 of these were doubtfully malignant and 15 were proved malignant, metastases being found in adjacent organs. Many adenomata have been removed with complete relief of symptoms. The first case of carcinoma with hypoglycaemia was reported by Wilder *et al.* (1927). Since then, including our own, 19 cases have been reported in the literature. To the best of our

knowledge ours is the first case to be reported in this country.

In making our selection we have included only cases of carcinoma with metastases and indisputable hypoglycaemia. Cases in which metastases occurred but hypoglycaemia was known to be absent were described as follows: three by Duff and Murray (1942), one by Frantz (1944), and one by Sailer and Zininger (1946). Fabozzi (1903) described four cases, and Zanetti (1927), Hamdi (1932), Evangelisti (1935), Gordon and Olivetti (1947), and Aragona (1948) each described one case in which there was not sufficient evidence to prove that hypoglycaemia occurred. Willis (1936) described one case in which the primary tumour was thought at first, on histological evidence, to have been in the pancreas; metastases occurred, but there was no hypoglycaemia. Later, after further examination, the tumour was considered to be an argentaffin carcinoma.

Bickel *et al.* (1935) described the case of a male diabetic of 56 years who later developed a carcinoma of the islets of Langerhans with multiple metastases. His liver contained many large secondary growths and little normal tissue remained. In three months the patient's daily excretion of urinary sugar fell from 260 g. to 8 g. and his insulin requirements from 100 units to *nil*. The only blood-sugar level recorded was 211 mg. per 100 ml. Insulin assay showed no insulin to be present in the liver growths, but in the primary tumour it was present "en quantité notable."

TABLE I.—Summary of 20 Cases of Carcinoma of Islets of Langerhans with Hypoglycaemia

Year	Name	Sex	Age	Lowest Blood Sugar (mg. per 100 ml.)	Duration of Symptoms in Months	Insulin Assay	Site of Primary in Pancreas	Size of Primary (cm.)	Site of Secondary Deposits	Type of Operation	Result
1927	Wilder <i>et al.</i>	M	40	25	20	Present	Tail	8×6×6. Single	Liver and lymph nodes	Cholecystectomy and biopsy of liver	Died 4 weeks p. op.
1931	Fedoroff*	M	31	27	?	Not tested	"	"Large"	Liver (no necropsy)	Biopsy of liver	" 4 weeks p. op.
1934	Judd <i>et al.</i>	F	18	45	4	" "	Not stated	Not stated	" "	" "	" 4 months p. op.
1934	Jacobsen	M	36	25	12-24	" "	Head	Grapefruit. Single "Diffuse"	Liver only	Biopsy	" 4 months p. op.
1937	Cragg <i>et al.</i>	F	41	30	9	Present	Whole	" "	Liver and lymph nodes	Biopsy of liver	" 4 months p. op.
1938	Joachim and Banowitch	F	31	25	3	Not tested	Distal half	6×6×7. "Tangerine"	" (no necropsy)	Resection of spleen, portion of growth in tail of pancreas and lymph nodes	" 6 weeks p. op.
1939	Seckelt†	M	36	16	12	" "	Proximal half	4×2×2	Generalized	Exploratory laparotomy	" 10 weeks p. op.
1941	Ballinger	M	53	30	12	Present	Liver†	10×6×6	"	Biopsy—tumour of aberrant pancreas	" 4 months p. op.
1941	Flinn <i>et al.</i>	F	45	20	6	Not tested	Head	6	Liver and lymph nodes	No operation	—
1942	Gray§	F	48	27	4½	" "	Tail	1.8×1.8×1.3	Liver	Excision of tumour in tail	" 6 weeks p. op.
1942	Quarrier and Bingham	M	73	21	1	" "	"	3	Liver and lymph nodes	No operation	—
1943	Hanno and Banks ..	M	68	21	19	" "	"	8×8×6.5	Liver	Biopsy of liver metastases	" 2 days p. op.
1943	Holman <i>et al.</i>	M	45	27	16-24	" "	"	12×8×8	Liver and lymph nodes	Biopsy of tumour and liver metastases	" 27 days p. op.
1943	Browning	M	36	12	20	" "	Body and tail	9. Single	" "	Biopsy of liver metastases	" 14 months p. op.
1944	Brunschwig <i>et al.</i> ..	M	32	14	54	" "	" "	9. Multiple	Liver	Sept., 1939—Excision of two tumours. Jan., 1940—Excision large tumours in body and tail. Aug., 1942—Excision most of head and liver metastases. Dec., 1943—Pancreatico-duodenectomy and excision of liver metastases	Sept., 1942—June, 1943, x rays to epigastrium. June—Aug., 1943, treated with alloxan. Died 2 hours p. op.
1947	Lopez-Kruger and Dockerty	M	42	69	2	" "	Tail	8. Single	Generalized	No operation	—
1947	Gordon and Olivetti ..	M	46	10	5½	" "	Body and tail	10×7×5	"	" "	Treated with alloxan
1947	Flinn <i>et al.</i>	F	55	37	2½	" "	Tail	4×6	Liver and lymph nodes	Biopsy of liver metastases	" "
1948	Levrat and Brette ..	F	31	29	24	" "	Head	"Large"	" "	No operation	Died 24 days p. op.
1949	Brearley and Laws ..	F	73	25	19	" "	Tail	15×8×7. 760 g. Single	Generalized	" "	—

* Cited by Lopez-Kruger and Dockerty (1947). † Reported also by Brunschwig, Gomori, and Cannon (quoted by Gray, 1942). ‡ Later considered to have arisen in aberrant pancreatic tissue (see text). § Earlier reported by Joslin and later by Marble and McKittrick.

There seems to be little doubt that this was an insulin-secreting carcinoma of the islets of Langerhans with metastases, but those authors do not claim that hypoglycaemia ever occurred.

Twelve of the 20 cases listed in Table I occurred in males and eight in females. The ages vary from 18 years to 73 years. In 13 cases the primary tumour was present in the tail of the pancreas, in four cases it was present in the head, and in two it was recorded as being in the pancreas. Ballinger (1941) considered that in his case the origin of the growth was in aberrant pancreatic tissue in the liver. Frantz (1944), who later saw this patient and was able to review the operative findings, thought that the tumour arose in aberrant pancreatic tissue behind and above the head of the pancreas. This is supported by the work of Faust and Mudgett (1940), who described 370 necropsies with aberrant pancreatic tissue and found that it occurred in this position but not in the liver.

Case History

A woman aged 73 was first seen in the out-patient department on March 28, 1949. She stated that in March, 1948, she began to have frequent headaches and attacks of dizziness on bending down. A short time later she suddenly developed a sharp pain in the mid-epigastrium which radiated round the left costal margin into her back. The pain was intermittent and was sometimes related to movement. It was not related to food. She had anorexia from December, 1948, and in January, 1949, she developed attacks of vomiting. These attacks were also intermittent and did not seem to be related to the pain. She weighed 8 st. 10 lb. (55.3 kg.) and thought that she had lost about a stone (6.4 kg.) during her illness. Bowel movement was normal. There was nothing significant in her past or family history.

On examination she looked well, and there was no obvious wasting. There was no jaundice. Examination of the abdomen revealed a firm rounded mass extending about three finger-breadths below the left costal margin. This moved with respiration and could be pushed forwards from behind. The liver edge was not felt. There was no ascites. She had slight hypertension; her B.P. was 185/110, but her heart was not clinically enlarged. The other systems were normal.

When admitted to hospital on May 2, 1949, her condition had improved. There had been no vomiting for a month, and she was eating well. On clinical examination the mass in the left hypochondrium was the same size as before. On inspiration the liver edge could now be felt one fingerbreadth below the right costal margin. It was firm and smooth, and no nodules could be palpated.

Investigations.—A barium meal showed no evidence of disease of the stomach or duodenum. A radiograph of the chest revealed calcification of the aortic knuckle and a number of healed tuberculous lesions at the right apex. Heart size and diaphragm were normal. Intravenous pyelography revealed normal pelves and calices; enlargement of the left kidney was excluded. Examination of faeces for occult blood was negative on two occasions. A catheter specimen of urine was normal. A blood count showed: haemoglobin, 11.8 g. per 100 ml. (85%); leucocytes, 5,000 per c.mm. (polymorphs 70%); sedimentation rate (Westergren), 25 mm. in 1 hour. Fractional test meal: histamine-refractory achlorhydria. Liver-function tests: see Table II.

Difficulty was experienced in deciding the origin of the tumour in the left hypochondrium. In view of its position, its firm edge, its movement with respiration, its bimanual palpation, and the fact that enlargements of the stomach and left kidney had been excluded radiologically, it was thought to be an enlarged spleen. This, taken in conjunction with the firm

TABLE II.—Liver-function Tests

	May 11	September 27
Total serum proteins ..	6.2 g. per 100 ml.	5.5 g. per 100 ml.
.. .. albumin ..	4.6 " " 100 "	3.8 " " 100 "
.. .. globulin ..	1.6 " " 100 "	1.7 " " 100 "
Serum alkaline phosphatase ..	41 K.-A. units per 100 ml.	62 K.-A. units per 100 ml.
Thymol turbidity ..	2	2
Takata-Ara ..	Negative	Negative
Direct van den Bergh reaction ..		
Serum bilirubin ..	< 0.2 mg. per 100 ml.	0.4 " mg. per 100 ml.

enlarged liver, made the diagnosis of cirrhosis with spleno-megaly reasonable in spite of the equivocal liver-function tests. The patient was now symptom-free, and she returned home on May 18 to continue on a high-protein and low-fat diet.

On July 27 she was seen again in the out-patient department and was found to be still free from symptoms.

She was readmitted on September 21, 1949. She had been well until 7 a.m. that day, when she is stated to have collapsed. She had recovered from this at midday. At 5 p.m. she again became drowsy and relapsed into coma.

On admission she was drowsy. Her skin was moist, but she was not hot. Her temperature was 96° F. (36.5° C.) and her pulse rate 84. Examination of her abdomen showed that her liver had increased in size considerably. The edge was now four fingerbreadths below the costal margin; it was hard and irregular. The mass in the left hypochondrium was unchanged. There was no jaundice or ascites. Her blood pressure was 165/95.

FIG. A.—Chart of six-hour glucose-tolerance test (September 26). The patient received 12 g. of glucose in water every hour during the night preceding the test, up to 8 a.m.

A diagnosis of hypoglycaemic coma was made and a glucose-saline intravenous drip transfusion was started. The blood sugar at this time was found to be 25 mg. per 100 ml. The patient recovered consciousness rapidly. Treatment was continued with frequent small feeds and glucose drinks hourly.

A six-hour glucose-tolerance test was attempted (Fig. A), but this had to be abandoned after five hours owing to hypoglycaemia. The initial blood-sugar level was only 52 mg. per 100 ml. in spite of the fact that 12 g. of sugar had been given of necessity only two hours previously. Liver-function tests were repeated (see Table II). They showed little change except for a further rise in the serum alkaline phosphatase.

At first the patient responded to this treatment. After a few days it became less effective and she tended to lapse into deep coma more easily, especially in the early morning, but she still responded to intravenous hypertonic glucose solution. Later, even hourly glucose drinks failed to prevent coma, and on October 1, 1949, she went into a coma from which she did not recover. She died at 11.50 a.m. on October 2.

Post-mortem Report

Permission was obtained for examination of only the chest and abdomen. At necropsy, performed twenty-four hours after death, the body was found to be well nourished.

FIG. B.—Inferior surface of liver, showing metastases.

Internal Examination.—Abdomen:—The peritoneum was clean and shiny. There was no ascites. The liver was grossly enlarged (Fig. B) and was filled with metastatic deposits varying in size between 1/6 and 1 in. (0.4 and 2.5 cm.). On the left side under cover of the enlarged liver was a hard lobulated mass lying in the position of the spleen (size 15 by 8 by 7 cm., weight 760 g.). On dissection this was found to be the tail of the pancreas (Fig. C). This mass had displaced the spleen upwards and had the left kidney buried in its posterior surface. A few secondary deposits were present in the omentum and a larger number in the peritoneum, diaphragm, and surrounding lymph nodes. The right adrenal gland contained small metastases. The alimentary tract was followed through and found to be normal. **Thorax:**—There was evidence of old pleurisy on both sides and a large left and smaller right pleural effusion. The lungs were

FIG. C.—Anterior aspect, showing the tumour and its relationship to other abdominal organs.

oedematous and friable. Small subpleural deposits were found in both lungs. These were believed to be carcinomatous. The presence of small amounts of pus in the bronchioles was consistent with a terminal bronchopneumonia. The heart was normal. The ascending aorta was atheromatous.

Histology.—Sections of the primary growth (Plate, Fig. 1) and of the metastases in lymph node, suprarenal, and liver (Fig. 2) showed the growth to consist of alveolar masses of round clearish cells, separated by thin strands of fibrous tissue, closely resembling those of islets of Langerhans. All the cells proved to be of chromophobe type. No alpha or beta cells could be demonstrated by specific stains.

Summary of Post-mortem Findings.—Primary carcinoma of islets of Langerhans with secondary deposits in liver, right adrenal gland, peritoneum, diaphragm, pleura, and surrounding lymph nodes.

Discussion

The clinical diagnosis after the patient's first admission to hospital was cirrhosis of the liver with splenomegaly. In view of the course of the disease and the severity of the hypoglycaemic attacks which developed, this had now to be reconsidered. There were two probable mechanisms by which this hypoglycaemia could be brought about. One was that the glycogen storage power of the body was deficient because of lack of active liver tissue; the other was that there was an overproduction of insulin. Mann (1927) has shown that in dogs only 15% of normal liver tissue is necessary to maintain adequate carbohydrate metabolism. In man such extreme liver deficiency is usually accompanied by ascites and jaundice. But that hypoglycaemia may occur without these is shown by the cases of Crawford (1931) and Nadler and Wolfer (1929), who describe hypoglycaemia in negroes with large primary hepatic carcinomata. Symptoms of hypoglycaemia may also occur in glycogen disease.

Our patient had a large liver which had rapidly increased in size and a tumour which closely simulated a spleen. There was no jaundice or ascites, and the liver-function tests, apart from the serum alkaline phosphatase, were normal. The glucose-tolerance curve was suggestive of hyperinsulinism.

The differential diagnosis was carcinoma of the islets of Langerhans with gross liver metastases or hypoglycaemia due to cirrhosis of the liver. In either case laparotomy was clearly unjustified and treatment necessarily palliative. It was decided to obtain a biopsy by liver puncture, but by the time the necessary preliminaries had been carried out the patient was *in extremis* and the test had to be abandoned. Clinical evidence pointed to the diagnosis of an islet-cell carcinoma, but in view of the extreme rarity of this condition and the absence of any histological evidence this could not be made with any degree of certainty.

Summary

A case of metastasizing islet-cell carcinoma of the pancreas causing fatal hypoglycaemia is described.

The literature on this subject is reviewed. Nineteen other cases in which the diagnosis is indisputable have been collected. In all, twelve occurred in males and eight in females. The ages varied between 18 and 73 years.

The clinical diagnosis in our patient is discussed.

We wish to thank Dr. H. P. Brody, under whose care the patient was admitted, for his encouragement and advice, Dr. L. C. D. Hermitte for the pathological studies, and Mr. F. Battersby for the photographs.

BIBLIOGRAPHY

- Aragona, F. (1948). *Pathologica*, **40**, 94.
 Ballinger, J. (1941). *Arch. Path.*, **32**, 277.
 Banting, F. G., Best, C. H., Collip, J. B., Campbell, W. R., and Fletcher, A. A. (1922). *Canad. med. Ass. J.*, **12**, 141.
 Bickel, G., Mozer, J. J., and Junet, R. (1935). *Bull. Soc. méd. Hôp. Paris*, **51**, 12.
 Browning, J. S. (1943). *Ann. intern. Med.*, **19**, 669.
 Brunschwig, A. (1941). *Surgery*, **9**, 554.
 — Allen, J. G., Owens, F. M., and Thornton, T. F. (1944). *J. Amer. med. Ass.*, **124**, 212.
 Cragg, R. W., Power, M. H., and Lindem, M. C. (1937). *Arch. intern. Med.*, **60**, 88.
 Crawford, W. H. (1931). *Amer. J. med. Sci.*, **181**, 496.
 Duff, G. L., and Murray, E. G. D. (1942). *Ibid.*, **203**, 437.
 Evangelis, T. (1935). *Policlinico, sez. chir.*, **42**, 384.
 Fabozzi, S. (1903). *Beitr. path. Anat.*, **34**, 199. Cited by Lopez-Kruger and Dockerty (1947).
 Faust, D. B., and Mudgett, C. C. (1940). *Ann. intern. Med.*, **14**, 717.
 Federoff, P. S. (1931). *Vrach. Gaz.*, **35**, 585.
 Flinn, L. B., Beatty, G. A., Ginsberg, M., and Hemsath, F. A. (1941). *J. Amer. med. Ass.*, **117**, 283.
 — Minnick, E., and Gay, D. M. (1947). *Ann. intern. Med.*, **26**, 936.
 Frantz, V. K. (1944). *Ann. Surg.*, **119**, 824.
 Gordon, B. S., and Olivetti, R. G. (1947). *Gastroenterology*, **9**, 409.
 Gray, L. M. (1942). *Amer. J. Path.*, **18**, 633.
 Hamdi, H. (1932). *Z. Krebsforsch.*, **37**, 411.
 Hanno, H. A., and Banks, R. W. (1943). *Ann. Surg.*, **117**, 437.
 Harris, S. (1924). *J. Amer. med. Ass.*, **83**, 729.
 Holman, E., Wood, D. A., and Stockton, A. B. (1943). *Arch. Surg. Chicago*, **47**, 165.
 Jacobsen, V. C. (1934). *Arch. Path.*, **18**, 135.
 Joachim, H., and Banowitch, M. M. (1938). *Ann. intern. Med.*, **11**, 1754.
 Judd, E. S., Faust, L. S., and Dixon, R. K. (1934). *West. J. Surg. Obstet. Gynec.*, **42**, 555.
 Levrat, M., and Brette, R. (1948). *Pr. méd.*, **56**, 530.
 Lopez-Kruger, R., and Dockerty, M. B. (1947). *Surg. Gynec. Obstet.*, **85**, 495.
 Mann, F. C. (1927). *Medicine*, **6**, 419.
 Nadler, W. H., and Wolfer, J. A. (1929). *Arch. intern. Med.*, **44**, 700.
 Quarrier, S. S., and Bingham, C. T. (1942). *Ann. Surg.*, **115**, 363.
 Sailer, S., and Zininger, M. M. (1946). *Surg. Gynec. Obstet.*, **82**, 301.
 Seckel, H. P. G. (1939). *J. clin. Invest.*, **18**, 723.
 Wilder, R. M., Allan, F. N., Power, M. H., and Robertson, H. E. (1927). *J. Amer. med. Ass.*, **89**, 348.
 Willis, R. A. (1936). *J. Path. Bact.*, **42**, 203.
 Zanetti, G. (1927). *Arch. Sci. med.*, **49**, 505.

AUREOMYCIN IN BRUCELLSIS

BY

E. W. LINDECK, M.R.C.P.

Physician, Royal Sussex County Hospital

Four cases of *Brucella abortus* infection have been treated by "aureomycin," and this paper is a tentative assessment of its value.

Case Reports

Case 1.—A man aged 61 gave one month's history of a febrile illness before his admission to hospital. The agglutination for *Brucella abortus* was found to be positive up to a titre of 1/2,500. On the fifth day after admission he was given aureomycin, 0.25 g., twice a day, and as no reaction to this dosage

FIG. 1.—Case 1.

had occurred 0.25 g. was given three-hourly and continued for nine days (total amount, 18.5 g.). His temperature became normal in 72 hours (Fig. 1).

During the course of treatment he experienced some indigestion. There has been no relapse since discharge from hospital 23 weeks ago.

Case 2.—A youth aged 14 gave six weeks' history of frontal headaches; there had been no response to a course of a sulphonamide drug. His blood was found to agglutinate *Br. abortus* up to a titre of 1/5,000. The white cells numbered

6,200 per c.mm. (polymorphs 30%, lymphocytes 51%, monocytes 11%, eosinophils 7%, basophils 1%). On the sixth day after admission he was treated with 16 g. of aureomycin spread over 12 days; 0.25 g. being administered twice a day on the first two days, and then four-hourly. The temperature subsided permanently in 72 hours (Fig. 2). When seen eight weeks later

FIG. 2.—Case 2.

his spleen was still palpable, which suggests the possibility of some residual infection; his agglutination titre still remained 1/5,000. Twenty-three weeks later there had been no relapse.

Case 3.—A man aged 46 gave a history of a P.U.O. lasting several months; his blood was found to be positive to *Br. abortus* up to a dilution of 1/2,500. Leucocytes numbered 7,600 (polymorphs 32%, lymphocytes 63%, monocytes 5%). A full course of sulphadiazine produced no improvement. Aureomycin was then begun, 0.1 g. being administered in five divided doses on the first day; this was followed by divided doses of 0.6 g. on the second day, 1.5 g. on the third day, and then 2 g. for three more days—a total of 8.2 g. On the second day of treatment his temperature rose to a higher level than before, but on the fourth day (Fig. 3) it became normal, and remained so until a relapse took place four months later. On this occasion the agglutination titre was 1/500. Leucocytes numbered 5,500 (polymorphs 61%, lymphocytes 33%, monocytes 6%). Aureomycin, 6.2 g., was given on the same plan as before, the temperature subsiding in 48 hours. The patient has remained well for eight and a half months.

FIG. 3.—Case 3: first illness.

FIG. 4.—Case 3: relapse.

Case 4.—A woman aged 25 came with a three-months history of ill-health with intermittent pyrexia. She showed a rising titre of agglutination to *Br. abortus* of 1/250 to 1/500. Aureomycin, 0.75 g., was administered four-hourly for ten and a half days (total quantity, 47.25 g.). She experienced nausea and one vomiting attack during treatment. The temperature subsided on the fourth day and there has been no relapse in six months. Unfortunately a temperature chart is not available.

Discussion

Brucellosis is notoriously a relapsing disease, which makes any claim of cure difficult to substantiate dogmatically after such comparatively short follow-up periods, but in every case quoted aureomycin has at least interrupted the rhythm of the febrile paroxysms, and the patients have been symptom-free. To be quite certain that the disease has been completely eradicated repeated blood cultures would be necessary.

It is evident that there is no standardized dosage. Case 3 relapsed after a course of 8.2 g.; this is evidently too small a total dosage. Spink *et al.* (1948), who treated 24 cases of brucellosis due to *Br. melitensis* (which is apparently

J. W. WHITTICK: NECROPSY FINDINGS IN Q FEVER

FIG. 1.—Lung. Pneumonic consolidation with thickening of alveolar walls and perivascular tissues by round cells. ($\times 60$.)

FIG. 2.—Predominantly round-cell exudate in alveolus. ($\times 400$.)

FIG. 3.—Testis. Compact aggregation of macrophages forming nodule. ($\times 400$.)

FIG. 4.—Cerebrum. Margin of area of softening. ($\times 150$.)

FIG. 5.—(a) Extracellular rickettsiae in pulmonary alveolar exudate. (b) Intracytoplasmic rickettsiae in pulmonary alveolar exudate. (c) Rickettsiae in macrophage of testicular nodule. (d) Rickettsiae beside and in neuroglial cells. (All $\times 1,400$.)

B. F. BREARLEY and J. W. LAWS: CARCINOMA OF ISLETS OF LANGERHANS WITH HYPOGLYCAEMIA

FIG. 1.—Primary tumour. ($\times 150$.)

FIG. 2.—Secondary tumour in the liver. ($\times 150$.)