BIBLIOGRAPHY

- Aragona, F. (1948). *Pathologica*, **40**, 94.
 Ballinger, J. (1941). *Arch. Path.*, **32**, 277.
 Banting, F. G., Best, C. H., Collip, J. B., Campbell, W. R., and Fletcher, A. A. (1922). *Canad. med. Ass. J.*, **12**, 141.
 Bickel, G., Mozer, J. J., and Junet, R. (1935). *Bull. Soc. méd. Hôp. Paris*, **51**, 12.
 Browning, J. S. (1943). *Ann. intern. Med.*, **19**, 669.
 Brunschwig, A. (1941). *Surgery*, **9**, 554.
 — Allen, J. G., Owens, F. M., and Thornton, T. F. (1944). *J. Amer. med. Ass.*, **124**, 212.
 Cragg, R. W., Power, M. H., and Lindem, M. C. (1937). *Arch. intern. Med.*, **60**, 88.
 Crawford, W. H. (1931). *Amer. J. med. Sci.*, **181**, 496.
 Duff, G. L., and Murray, E. G. D. (1942). *Ibid.*, **203**, 437.
 Evangelis, T. (1935). *Policlinico, sez. chir.*, **42**, 384.
 Fabozzi, S. (1903). *Beitr. path. Anat.*, **34**, 199. Cited by Lopez-Kruger and Dockerty (1947).
 Faust, D. B., and Mudgett, C. C. (1940). *Ann. intern. Med.*, **14**, 717.
 Federoff, P. S. (1931). *Vrach. Gaz.*, **35**, 585.
 Flinn, L. B., Beatty, G. A., Ginsberg, M., and Hemsath, F. A. (1941). *J. Amer. med. Ass.*, **117**, 283.
 — Minnick, E., and Gay, D. M. (1947). *Ann. intern. Med.*, **26**, 936.
 Frantz, V. K. (1944). *Ann. Surg.*, **119**, 824.
 Gordon, B. S., and Olivetti, R. G. (1947). *Gastroenterology*, **9**, 409.
 Gray, L. M. (1942). *Amer. J. Path.*, **18**, 633.
 Hamdi, H. (1932). *Z. Krebsforsch.*, **37**, 411.
 Hanno, H. A., and Banks, R. W. (1943). *Ann. Surg.*, **117**, 437.
 Harris, S. (1924). *J. Amer. med. Ass.*, **83**, 729.
 Holman, E., Wood, D. A., and Stockton, A. B. (1943). *Arch. Surg. Chicago*, **47**, 165.
 Jacobsen, V. C. (1934). *Arch. Path.*, **18**, 135.
 Joachim, H., and Banowitch, M. M. (1938). *Ann. intern. Med.*, **11**, 1754.
 Judd, E. S., Faust, L. S., and Dixon, R. K. (1934). *West. J. Surg. Obstet. Gynec.*, **42**, 555.
 Levrat, M., and Brette, R. (1948). *Pr. méd.*, **56**, 530.
 Lopez-Kruger, R., and Dockerty, M. B. (1947). *Surg. Gynec. Obstet.*, **85**, 495.
 Mann, F. C. (1927). *Medicine*, **6**, 419.
 Nadler, W. H., and Wolfer, J. A. (1929). *Arch. intern. Med.*, **44**, 700.
 Quarrier, S. S., and Bingham, C. T. (1942). *Ann. Surg.*, **115**, 363.
 Sailer, S., and Zininger, M. M. (1946). *Surg. Gynec. Obstet.*, **82**, 301.
 Seckel, H. P. G. (1939). *J. clin. Invest.*, **18**, 723.
 Wilder, R. M., Allan, F. N., Power, M. H., and Robertson, H. E. (1927). *J. Amer. med. Ass.*, **89**, 348.
 Willis, R. A. (1936). *J. Path. Bact.*, **42**, 203.
 Zanetti, G. (1927). *Arch. Sci. med.*, **49**, 505.

AUREOMYCIN IN BRUCELLSIS

BY

E. W. LINDECK, M.R.C.P.

Physician, Royal Sussex County Hospital

Four cases of *Brucella abortus* infection have been treated by "aureomycin," and this paper is a tentative assessment of its value.

Case Reports

Case 1.—A man aged 61 gave one month's history of a febrile illness before his admission to hospital. The agglutination for *Brucella abortus* was found to be positive up to a titre of 1/2,500. On the fifth day after admission he was given aureomycin, 0.25 g., twice a day, and as no reaction to this dosage

FIG. 1.—Case 1.

treatment he experienced some indigestion. There has been no relapse since discharge from hospital 23 weeks ago.

Case 2.—A youth aged 14 gave six weeks' history of frontal headaches; there had been no response to a course of a sulphonamide drug. His blood was found to agglutinate *Br. abortus* up to a titre of 1/5,000. The white cells numbered

6,200 per c.mm. (polymorphs 30%, lymphocytes 51%, monocytes 11%, eosinophils 7%, basophils 1%). On the sixth day after admission he was treated with 16 g. of aureomycin spread over 12 days; 0.25 g. being administered twice a day on the first two days, and then four-hourly. The temperature subsided permanently in 72 hours (Fig. 2). When seen eight weeks later

FIG. 2.—Case 2.

his spleen was still palpable, which suggests the possibility of some residual infection; his agglutination titre still remained 1/5,000. Twenty-three weeks later there had been no relapse.

Case 3.—A man aged 46 gave a history of a P.U.O. lasting several months; his blood was found to be positive to *Br. abortus* up to a dilution of 1/2,500. Leucocytes numbered 7,600 (polymorphs 32%, lymphocytes 63%, monocytes 5%). A full course of sulphadiazine produced no improvement. Aureomycin was then begun, 0.1 g. being administered in five divided doses on the first day; this was followed by divided doses of 0.6 g. on the second day, 1.5 g. on the third day, and then 2 g. for three more days—a total of 8.2 g. On the second day of treatment his temperature rose to a higher level than before, but on the fourth day (Fig. 3) it became normal, and remained so until a relapse took place four months later. On this occasion the agglutination titre was 1/500. Leucocytes numbered 5,500 (polymorphs 61%, lymphocytes 33%, monocytes 6%). Aureomycin, 6.2 g., was given on the same plan as before, the temperature subsiding in 48 hours. The patient has remained well for eight and a half months.

FIG. 3.—Case 3: first illness.

FIG. 4.—Case 3: relapse.

Case 4.—A woman aged 25 came with a three-months history of ill-health with intermittent pyrexia. She showed a rising titre of agglutination to *Br. abortus* of 1/250 to 1/500. Aureomycin, 0.75 g., was administered four-hourly for ten and a half days (total quantity, 47.25 g.). She experienced nausea and one vomiting attack during treatment. The temperature subsided on the fourth day and there has been no relapse in six months. Unfortunately a temperature chart is not available.

Discussion

Brucellosis is notoriously a relapsing disease, which makes any claim of cure difficult to substantiate dogmatically after such comparatively short follow-up periods, but in every case quoted aureomycin has at least interrupted the rhythm of the febrile paroxysms, and the patients have been symptom-free. To be quite certain that the disease has been completely eradicated repeated blood cultures would be necessary.

It is evident that there is no standardized dosage. Case 3 relapsed after a course of 8.2 g.; this is evidently too small a total dosage. Spink *et al.* (1948), who treated 24 cases of brucellosis due to *Br. melitensis* (which is apparently

more resistant to aureomycin than *Br. abortus*), found that relapses occurred even after a course of 21 g., and in one case a blood culture was positive after a course of 25 g.; two other cases had positive cultures after 20 and 21 g. These authors at first combined a course of sulphadiazine with the aureomycin, but they discarded the former as it did not appear to influence results. Their final recommendation is that the dosage should be approximately from 4 to 6 g. daily for two weeks. They also describe a "Herxheimer" type of reaction in 12 out of the 24 patients with an abrupt rise of temperature sometimes accompanied by a fall of blood pressure and tachycardia some eight to twelve hours after the initial dose of aureomycin if this was as high as 0.5 g. Therefore they recommend that 0.1 g. should be given in divided doses on the first day, 0.6 g. on the second day, 1.6 g. on the third day, and the full dose on the fourth day. It will be observed from Fig. 3 that there was an abrupt rise of temperature to 103° F. (39.4° C.), in spite of the cautious dosage, on the second day of treatment; there do not seem to have been any serious clinical symptoms. The maximum follow-up in Spink's series of cases is a period of three months; they were well satisfied with the clinical response of patients to treatment.

Knight *et al.* (1949) report five cases of brucellosis treated with aureomycin, four being acute cases and one a bacteraemia; their standard course consisted of 6 g. on the first day in divided doses, and thereafter 4 g. daily for five days—a total of 26 g. The report suggests that the immediate beneficial result of aureomycin is very striking, but two of the five cases relapsed after 15 and 42 days respectively; the former patient had received 9.5 g. for seven days instead of the standard course; evidently the length of time the drug is administered, as well as the total dosage, has to be taken into consideration. Knight *et al.* (1949) also record a "Herxheimer" type of reaction in one of their patients. Side-effects consisting of gastric disturbances were unimportant. Their maximum follow-up period was five months.

Heilman (1949) reports that in mice aureomycin alone is not particularly effective in reducing the number of brucella in the spleen of infected mice; aureomycin combined with streptomycin or dehydrostreptomycin was the most effective, but even this combination did not eliminate all the organisms. Herrell and Barker (1949), at the same clinic, employed subsequently a combination of aureomycin (3 g. in divided doses daily) and dehydrostreptomycin (1 g. twice a day intramuscularly for 12 to 14 days) in four cases with initial success; the infecting organisms were *Br. suis* two cases and *Br. abortus* two cases. Their blood cultures after treatment in each case were negative, which suggests a permanent cure. Spink *et al.* (1948) had previously used streptomycin, but discarded it because of its unfortunate effect on the labyrinth. No such effect is recorded with dehydrostreptomycin.

In the present state of our knowledge it would therefore appear reasonable, in an average case of brucellosis in this country, to employ aureomycin in doses of 4 to 6 g. daily for a fortnight, with smaller amounts on the first three days (0.1 g., 0.6 g., and 1.6 g. in divided doses). The ease of administration by the oral route and the lack of important side-effects are important considerations. Should the case prove resistant or relapse frequently, dehydrostreptomycin and aureomycin in the dosage suggested by Herrell and Barker might well be employed.

It is to be hoped that further cases treated with aureomycin and controlled by subsequent blood cultures will soon be published by others.

My thanks are due to Dr. W. A. Bourne and to Dr. John Cumming, of Hove, for particulars of one case, to Dr. A. Head, of Storrington, Sussex, for particulars of another; also to Dr. Bremner, of Hove Hospital, and Dr. Pearson, of the Royal Sussex County Hospital.

REFERENCES

- Heilman, F. R. (1949). *Proc. Mayo Clin.*, **24**, 133.
 Herrell, W. E., and Barker, T. E. (1949). *Ibid.*, **24**, 138.
 Knight, V., Ruiz-Sanchez, F., Ruiz Sancho, A., and McDermott, W. (1949). *Amer. J. Med.*, **6**, 407.
 Spink, W. W., Braude, A. I., Costaneda, M., and Goyta, L. (1948). *J. Amer. med. Ass.*, **138**, 1145.

BRACHIAL NEURALGIA

BY

W. M. PHILIP, M.B., M.R.C.P.

Assistant Physician, Selly Oak Hospital, Birmingham

The syndrome of sciatica due to a prolapsed intervertebral disk is well recognized, and in essence consists of symptoms and signs resulting from irritation of a single nerve root. The muscles supplied by the root are tender, painful, and weak, while pain radiates down the dermatome, and frequently paraesthesia, numbness, sensory loss, and interference with the appropriate reflex arc are present. In addition, there is often a history of lumbago, and examination reveals protective spasm of the erectores spinae.

Brachial neuralgia is at present a less exact diagnosis than sciatica, and includes many diseases in which the presenting symptom is pain radiating down the arm; but there is a group of cases in which the symptoms are those of root irritation comparable to that found in sciatica, and in these a diagnosis of prolapsed intervertebral disk is commonly made. There are, however, reasons for believing that such a lesion is not the commonest cause of brachial neuralgia, and that, instead of a prolapse of the nucleus pulposus, arthritic changes in the neurocentral joint are the most usual source of root irritation in the neck (Lund, 1945; Bull, 1948).

A total of 66 cases has been selected from those seen personally in the out-patient department. Each has borne a superficial resemblance to brachial neuralgia, but of these cases 24 have been discarded as belonging to a miscellaneous group which defies classification. In all the others (42) it has seemed probable that the symptoms were the result of irritation of one or more cervical roots; the diagnosis was made primarily on clinical grounds, and the criteria on which individual cases were assessed were: (1) Radiation of pain into the arm. Usually this was referred accurately enough to one or more dermatomes to permit clinical location of the lesion. In only five cases was this not so. (2) Pain with or without tenderness in the muscles supplied by the appropriate root. (3) X-ray changes corresponding with the root involved.

A single root appeared to be involved in 24 cases, and it is in these in which the disease is localized that one may expect to find the characteristic x-ray appearances. The roots affected were: root 5, 1 case; root 6, 7 cases; root 7, 15 cases; root 8, 1 case.

With the multiplication of levels and the development of hypertrophic spondylitis, every variety of change may be found. The important point is that narrowing of the disk and the lesions that result from it are so constantly seen: in only two cases out of the 24 were the radiographs entirely normal. In the remaining cases the lesions were both clinically and radiologically multiple; in 10 the symptoms were bilateral, but one rather striking feature is the way in