SCARLET FEVER AND TONSILLITIS IN
RELATION TO ACUTE RHEUMATISM

BY

T. C. MACDONALD, A.F.C., M.D., D.P.H.

Wing Commander, R.A.F.

While the infectious nature of rheumatic fever was postulated by Hirsch the aetiological agent is still not positively identified. The various hypotheses have been set forth by Hadfield and Garrod (1947), who say: "It seems to us indisputable that more progress has been made towards identifying haemolytic streptococci as the cause of this disease than in any other direction."

Galabin (1932) and Longstaff (1904-5) showed the relationship between mortality rates of rheumatic fever, erysipelas, scarlet fever, septicaemia, and puerperal fever. Atwater (1927) published similar figures.

Accurate morbidity figures are not so easy to obtain. It was felt, therefore, that some information might be drawn from R.A.F. and W.A.A.F. sickness records, as these had certain advantages not usually found. The numbers were large, the population at risk was known, medical care was at all times easy of access, and the recording system is well tried. There is no difference in nutrition, whether in quality or in quantity of food; clothing is identical, and living conditions vary very little. There is a rough standardization for age by the fact that children, adolescents, and the aged are not in the Service. The figures for the sexes are given separately. On the other hand, one condition which facilitates infection by *Streptococcus pyogenes* is present. This is the communal living and, in particular, sleeping in barrack rooms. Other factors are overcrowding and constant influx of newcomers into each little community. I will consider only figures for the Forces in this country.

The figures (Tables I and II, Figs. 1 and 2) are obtained from the Air Ministry (1938-45) and from Department MA7 of the Medical Directorate General, which kindly supplied the figures for 1946 and 1947, which have not yet been published.

Comments

Glover (1942) discussed rheumatic fever in closed communities and showed its relationship with tonsillitis. It will be observed that when a very large number of samples are taken this relationship vanishes or almost vanishes. This is not surprising, as the diagnosis "tonsillitis" unsupported by bacteriological proof is a blunderbuss term. There is some hint of resemblance in the male statistics but none in the female. It will be seen, too, that the tonsillitis figures remain about the same level throughout, while rheumatic fever shows a tendency to drop.

"Tonsillitis" as an indicator of impending acute rheumatic cases would seem to be of little value.

The resemblance between the morbidity of scarlet fever and rheumatic fever is remarkable. This

FIG. 1.—Incidence of tonsillitis, scarlet fever, and acute rheumatism in the R.A.F. Log. scale.

FIG. 2.—Incidence of tonsillitis, scarlet fever, and acute rheumatism in the W.A.A.F. Log. scale.

TABLE I.—Incidence of Tonsillitis, Scarlet Fever, and Acute Rheumatism in the R.A.F. and W.A.A.F. at Home

Disease	Year	R.A.F.				W.A.A.F.			
		No. of Cases	No. of Deaths	Incidence per 1,000 p.a.	Average No. of Days Treatment	No. of Cases	No. of Deaths	Incidence per 1,000 p.a.	Average No. of Days Treatment
Tonsillitis	1938	2,305	—	36.00	8	—	—	—	—
	1939	4,436	—	35.94	8	45	—	19.6	6
	1940	9,396	1	32.10	9	667	—	50.97	8
	1941	14,048	—	24.02	9	2,110	—	43.79	8
	1942	15,752	—	23.93	9	6,271	—	49.08	9
	1943	19,505	—	29.44	8	9,403	—	52.62	9
	1944	25,131	1	36.75	8	8,980	—	51.88	9
	1945	19,230	—	30.08	8	6,670	—	48.65	9
	1946	12,012	1	35.19	9	2,537	—	50.31	9
	1947	8,715	—	35.24	9	1,401	—	61.23	9
Scarlet fever	1938	99	—	1.5	36	—	—	—	—
	1939	329	—	2.7	36	2	—	0.9	28
	1940	699	—	2.4	32	12	—	0.9	27
	1941	647	—	1.1	30	91	—	1.9	29
	1942	455	—	0.69	29	209	1	1.64	29
	1943	815	1	1.23	26	297	—	1.66	30
	1944	1,180	—	1.73	22	240	—	1.39	26
	1945	301	1	0.47	30	70	—	0.51	22
	1946	336	—	0.98	27	45	—	0.89	27
	1947	496	—	2.01	26	42	—	1.84	26
Acute rheumatism	1938	132	—	2.1	76	—	—	—	—
	1939	300	—	2.4	48	4	—	1.7	23
	1940	630	1	2.2	33	29	—	2.2	23
	1941	952	—	1.6	31	75	—	1.6	30
	1942	649	1	0.99	26	666	—	5.21	24
	1943	413	—	0.62	38	357	1	2.00	25
	1944	724	—	1.06	52	186	—	1.07	36
	1945	268	—	0.42	34	100	—	0.70	43
	1946	251	—	0.74	51	38	1	0.70	40
	1947	299	—	1.21	56	21	—	0.92	44

TABLE II.—Coefficients of Correlation Between Acute Rheumatism and Scarlet Fever and Tonsillitis

	R.A.F.	W.A.A.F.
Acute rheumatism and scarlet fever ..	+0.83	+0.45
Acute rheumatism and tonsillitis ..	+0.28	-0.14

is particularly so in the R.A.F. The W.A.A.F. statistics are less reliable, as the Force was expanded quicker and had always a higher rate of influx and egress. It will be seen that these rates not only are highly correlated but are of the same order of magnitude.

If the statistics are accepted as valid they must have some practical interest. I feel that too much emphasis in the past has been placed on the value in epidemic prediction of the known relationship of streptococcal tonsillitis with acute rheumatism. This is, of course, the result of laboratory influence. The point must also be made that scarlet fever is notifiable and the other two diseases are not.

In brief we can say: (a) Tonsillitis figures are of very little use on a national basis as a warning of an imminent rise in acute rheumatism morbidity. In any case they are not available for the civil population. (b) Scarlet fever figures are available, and it would appear that in the adult (but not aged) population there can be expected about the same number of acute rheumatism cases as there are of scarlet fever. (c) A rising national scarlet-fever rate should be a signal for medical officers of health to look for the haemolytic streptococcus in epidemics of sore throats, however small these epidemics may be. (d) The R.A.F. and similar statistics might be of use to public health authorities, as they include non-notifiable diseases.

I am indebted to the Director-General Medical Services, Royal Air Force, for permission to publish.

REFERENCES

- Air Ministry (1938-45). Statistics on the Health of the Royal Air Force and Women's Auxiliary Air Force. (Air Publication 875.) London.
 Atwater, R. M. (1927). *Amer. J. Hyg.*, 7, 343.
 Galabin, A. L. (1902). *Lancet*, 1, 1671.
 Glover, J. A. (1942). Rheumatic Fever; in *Lancet*, Control of Common Fevers, London.
 Hadfield, G., and Garrod, L. P. (1947). *Recent Advances in Pathology*. London.
 Longstaff, G. B. (1904-5). *Trans. epidem. Soc. Lond.*, 24, 33.

A report in the *New York Times* of testimony given before the Joint Congressional Committee on Atomic Energy describes the United States' preparations for a national blood transfusion service to be used in the event of atomic attack. It has been recommended that a peacetime blood donor system should be set up that would be ready if necessary to meet extraordinary wartime requirements. Dr. Norvin C. Kiefer, from the Office of Civilian Mobilization, estimated that "several million" pints of atomic might be needed within the first few weeks of an atomic attack. He said that a huge reserve of standard bottles and other supplies must be ready to obtain blood in large quantities from persons outside the attacked area. It was possible to devise on paper an organization which could recruit millions of blood donors within a period of a few days, but the technical problems would be enormous and without actual experience of a comparable peacetime service the task would be almost impossible. Such a service would answer a peacetime need and offer a strong foundation on which an expanded wartime programme could be built with relative ease, safety, and speed. "Standardization of equipment is imperative," added Dr. Kiefer, "and in addition great efforts must be made to extend the period in which whole blood can be preserved, and to develop more effective blood substitutes." Conferences among the interested agencies are going on to this end.

Medical Memoranda

Cranial Arteritis Presenting as Meningitis

[WITH SPECIAL PLATE]

Temporal or cranial arteritis was first described by Jonathan Hutchinson in 1890. In 1932 Horton and Magath, in America, rediscovered the condition and described two cases, and in 1938 Jennings submitted the first report of two cases in England. Since then many cases have been described, and Harrison (1948) collected and summarized the outstanding features of 75 cases.

The condition has been reviewed and its relationship with periarteritis nodosa discussed by Miller (1949). He points out the widespread nature of the disease, which may affect intracranial vessels and larger arteries of the chest, trunk, and limbs.

Two features of interest in the case reported below were the marked signs of meningeal irritation and the successful therapeutic effect of infiltration of the affected areas with 2% procaine. This measure was suggested by Dantes (1946) and was successfully used in four cases by Roberts and Askey (1949).

CASE REPORT

A housewife aged 68 was admitted to West Middlesex Hospital on April 9, 1949. Her chief complaints were of headache and photophobia. She also complained of tender lumps on her face, forehead, and scalp, of long-standing malaise, and of recent feverishness and vomiting.

She had last been quite well nine months previously. During the winter she had noticed stiffness and pain in her knees and ankles, which massage and heat only aggravated. In April, 1949, she began to get headaches; these were shooting stabs of pain "from ear to ear," numbering 20 to 30 a day. Soon she began to get pain across her shoulders, made worse by changes in posture; pain appeared deep in the front of the thighs, and her calf muscles would ache badly on exercise. Generally off-colour and with poor appetite, she went to a convalescent home, where three weeks' rest brought much subjective improvement and lessening of pain. A few days after her return home the pains and general symptoms recurred. She noticed loss of weight and her increasingly pallid complexion.

Two weeks before admission, lumps appeared in her neck, just in front of both ears, and on the back of the head. They were extremely tender to touch and the overlying skin was reddened. Over the subsequent week her temperature rose to 101° F. (38.3° C.), and she began to get a constant throbbing headache, made worse by movements of the head or on bending down. She began to vomit four or five times a day, her jaws became stiff, and the headache invaded the back of her neck. She lay sick and miserable in bed with the curtains drawn, as "the light hurt her eyes."

On admission the patient was drowsy, flushed, and perspiring. She resented interference, and lay on her side with her head retracted and her eyes tightly closed. Her temperature was 99.2° F. (37.3° C.) and pulse rate 90.

On examination the most noteworthy feature was the presence of tender inflamed pulsatile lumps along the course of both superficial temporal arteries, involving the forehead and areas in front of and behind both ears. Moreover, the external carotid areas below the angles of the jaw were tender and inflamed, and there were numerous tender lumps in the scalp in the occipital and vertical regions. There was tenderness on palpation along the course of the common carotid and the right brachial arteries. Palpation of the abdominal aorta gave rise to pain, though this is often found in normal subjects. Neck rigidity was present and photophobia was marked,