REFRESHER COURSE FOR GENERAL PRACTITIONERS

HYPERTENSION—I

BY

ROBERT PLATT, M.D., M.Sc., F.R.C.P.*Professor of Medicine, University of Manchester*

Hypertension is a syndrome and not a disease. A large number of cases with high blood pressure fall into the group which is known as primary or essential hypertension. This means that no pre-existing disease which might have caused the hypertension can be discovered. In other words, as Wakerlin (1949) has said, "‘Essential hypertension’ is a designation born of ignorance and is equivalent to ‘fever of unknown cause.’" Although we are ignorant of the cause of essential hypertension I believe that it is much more than a merely negative diagnosis, and that it will turn out to be a definite entity with a recognizable pathogenesis. A family history of hypertensive disease is present in a high proportion of cases of essential hypertension. They tend to conform to a certain bodily type, and many believe that they also have certain psychological characteristics.

Essential hypertension tends to run a definite course; it probably begins quite early in life, but is usually seen by the physician in middle-aged or elderly persons.

Hypertension secondary to renal or endocrine disorders is, on the other hand, commoner in young persons. The most important renal causes of secondary hypertension are nephritis, pyelonephritis, congenital abnormalities of the kidney, and pregnancy toxæmia. The endocrine causes are Cushing's syndrome and phaeochromocytoma, and occasional cases are due to coarctation of the aorta.

In these two brief articles I shall not deal with the rarer of these causes of hypertension, nor shall I discuss the systolic hypertension of old people. As the aorta and large vessels become less and less elastic a rise in the systolic pressure without any similar increase in the diastolic pressure is the natural result according to simple physical laws. The finding of a blood pressure of 180/90 in an arteriosclerotic person approaching 70 years of age should simply be regarded as part of the process of growing old.

The Mechanism of Hypertension

It is natural to suppose that an increase in blood pressure is a process of adaptation, and it may therefore have a purpose in overcoming resistance in some part of the circulation. There is considerable evidence that this is the case.

It is known that in uncomplicated human hypertension there is no change in the cardiac output, in the total blood volume, or in the viscosity of the blood. The increased pressure is unquestionably due to a constriction of the arterioles, causing an increase in the peripheral resistance. Although the cardiac output remains normal the work of the heart is of course increased.

The modern work on experimental hypertension begins with the experiments of Goldblatt *et al.* (1934), who showed that by constricting the renal arteries and thereby reducing the blood supply to the kidneys a rise in blood pressure was promptly and constantly produced. If the constriction was moderate a condition resembling benign essential hypertension was produced, and if the constriction was severe the state of the animals resembled that of malignant hyper-

tension in that they developed arteriolar necrosis in various organs, though not, however, in the kidneys, because by the very nature of the experiment the renal arterioles were protected from the effects of the high blood pressure. If the renal artery of only one kidney was constricted, hypertension was usually more transient and less severe, and if the affected kidney was removed the blood pressure returned to normal.

Wilson and Byrom (1939) found that severe hypertension of the malignant type could be produced in rats by constriction of only one renal artery, in which case the pathological changes of malignant hypertension, with their characteristic arteriolar necrosis, could be found in the opposite kidney. They speak of a vicious circle in hypertension (Wilson and Byrom, 1941) in the sense that ischaemia of the kidney sends up the blood pressure, and if the rise in diastolic hypertension is severe enough arteriolar changes are produced which damage the kidney still further and so produce an even greater state of renal ischaemia. That the arteriolar injury is in fact due to increased intravascular tension is shown by Byrom and Dodson (1948) in further experiments in which they raised the intravascular pressure acutely by injection of saline and produced arteriolar necrotic lesions in the kidney. The bearing of these findings upon human hypertension will be discussed later.

Soon after the original experiments had been made it was found that the hypertension produced by renal ischaemia was not brought about by a nervous mechanism. It had long been known that a substance called "renin," which was capable of raising the blood pressure, could be extracted from the kidney, and it is now known that renin is an enzyme which acts on some substance present in the plasma to produce another substance now known as hypertensin, which raises the blood pressure. Other enzymes (hypertensinases), which seem to be widely distributed in the body, are known to destroy hypertensin.

Since experimental hypertension in animals so closely resembles essential hypertension in man, it is tempting to assume that the initial pathology of essential hypertension is a narrowing of renal arterioles, leading to ischaemia of the kidney and an excessive production of renin. This would suggest a compensatory purpose to ensure that the kidney received an adequate blood supply. An enormous amount of subsequent investigation, however, has failed to prove the connexion between the kidney and essential hypertension. There are, in fact, certain rather important objections to the hypothesis. The first is that biopsies of renal tissue removed at operation in patients with essential hypertension show no significant renal vascular lesions in nearly 30% of cases (Castleman and Smithwick, 1943). Against this objection one may ask what is a significant vascular lesion, and one may also point out that a very small reduction in arteriolar calibre may produce a very considerable obstruction to blood flow when the whole vascular bed of the organ is being considered.

The second objection to the renin theory is that excess of renin cannot be discovered in the blood in chronic

hypertensive states in man, but here it must also be admitted that it cannot be discovered in similar chronic states of hypertension in animals, despite the fact that the hypertension has in that case been caused by interference with the renal circulation. In acute hypertension in animals and in man (pregnancy toxæmia and acute nephritis) Dexter and Haynes (1944) have shown that an excess of renin may be present, so that it is possible that in the chronic stage another mechanism is brought into play, or else the amount of renin present is too small to be detected by methods at present in use.

We know that hypertension, apparently indistinguishable from essential hypertension, occurs in Cushing's syndrome, in which the primary defect is in the pituitary or the adrenal gland, and in the treatment of Addison's disease hypertension has often been produced by deoxycortone and sodium chloride. Selye *et al.* (1943) have produced nephrosclerosis and hypertension in animals by the administration of corticoid compounds and salt. Shorr *et al.* (1945) have produced a vaso-excitor material by the anaerobic incubation *in vitro* of normal kidney. They believe that it is distinct from hypertensin, that it plays a part in the response of the vascular system to shock, and that it may have some significance in the production of hypertension. It is thus possible, if not probable, that hormonal factors other than renin play an important part in the production of hypertension in man, and we must leave the controversy at present unsettled.

The fact that the blood pressure is sometimes reduced by sympathectomy operations does not in any way prove that essential hypertension in man is produced by a nervous mechanism, since lowering of sympathetic tone might reduce the blood pressure whatever its initial mechanism.

That hypertension in man may have a renal origin is shown by its frequent occurrence in nephritis, and by the fact that occasional cases are known in which the removal of a pathological kidney has reduced the blood pressure to normal. The question at issue is therefore not whether hypertension in man may sometimes be due to kidney disease, but whether all forms of human hypertension, especially the essential cases, are renal in origin.

Essential Hypertension

Essential hypertension is usually seen in the so-called benign form. Some authorities estimate that about one-quarter of the population suffer from some degree of hypertension in later life, and in many of these it is the direct or indirect cause of death. In all probability the condition starts quite early in life but, being symptomless, is not usually recognized. Anyone who has taken the blood pressures of a large number of students or recruits realizes that a proportion of them tend to have a blood pressure higher than average, especially if they are nervous of the examination. There is considerable evidence that this group is largely composed of people who will tend to have hypertension later in life.

Ayman (1948) has shown that this tendency for the blood pressure to rise to abnormal levels is much commoner in the children of hypertensive parents, and Platt (1947) has shown that, whereas a family history suggestive of high blood pressure is present in most cases of essential hypertension, this does not apply in cases of hypertension secondary to renal disease. Persons who suffer from essential hypertension in middle age are often rather broadly built, with broad faces, fair hair soon becoming grey, and a rather ruddy, healthy-looking complexion. They tend to look rather more than their age and often show a

tendency to nervous flushing, not only of the face but of the neck and upper chest. Their blood pressures are usually extremely labile, going up with the least anxiety, and coming down almost to normal levels with rest and reassurance. An initial blood pressure of 230/120 may become 150/90 after a few days' observation in hospital.

Usually, hypertension remains symptomless until middle age or later. It may be discovered for the following reasons.

First, during a routine medical examination undertaken for life insurance or other purposes.

Secondly, the blood pressure is usually taken as a routine during pregnancy and the early stages of essential hypertension may thus be revealed.

Thirdly, women at the menopause who complain of flushing sensations frequently ask their doctors to test the blood pressure.

Fourthly, hypertension may be revealed by the onset of symptoms caused by the disease.

Cardiac failure is the cause of death in most cases of essential hypertension. The commonest symptoms therefore are those of cardiac origin. They may take the form of progressive exertional dyspnoea, later accompanied by ankle oedema; or of acute left ventricular failure, with attacks of paroxysmal nocturnal dyspnoea; or of angina pectoris. Hypertension predisposes to coronary occlusion, an attack of which may be the first indication of illness. Cerebral complications, next to heart failure, are the commonest cause of death, and cases may be seen for the first time because of a sudden hemiplegia. Benign essential hypertension never leads to renal failure, though renal-function tests may be slightly depressed in elderly hypertensive subjects, and occasionally there is a little albumin in the urine, especially if cardiac congestion is present.

The relation of headache and dizziness to hypertension is very difficult to define. It is certain that many patients with extreme degrees of high blood pressure live and die without complaining at any time of either symptom. If dizziness is described it is never rotatory or labyrinthine in type. Patients who give a clear history of vertigo of this kind will always be found to have some upset of labyrinthine function, whatever may be the height of their blood pressure. The so-called dizziness of the hypertensive is a feeling of light-headedness, especially on regaining the upright position after stooping.

Many cases of malignant hypertension (described below) develop severe paroxysmal incapacitating headache, often accompanied by vomiting. Frequent or incapacitating headache in benign hypertension is unusual except in those who have had a tendency to headache most of their lives—many of them have had typical migraine in former years—and in the very many hypertensive subjects who have been given an anxiety neurosis by the discovery of their high blood pressure. It would not be far from the truth to say that the majority of symptoms of hypertension are iatrogenic. True hypertensive headache may be frontal or occipital and is usually most prominent in the early morning.

Malignant Hypertension

Malignant hypertension, which is probably only a more severe variety of essential hypertension, is distinguished by the development of papilloedema and other manifestations of hypertensive retinopathy, a very high diastolic pressure (140 or over), and a rapidly downhill course leading to renal failure and death within two years of the onset of symptoms. The age at which symptoms commonly appear is between 40 and 50. Mostly they take the form of severe

paroxysmal headache with vomiting, but in some cases cardiac symptoms are prominent. In other cases it is loss of vision which brings the patient first to the optician and then to the doctor. Usually the patient when first seen is clearly in the malignant phase of hypertension; sometimes he is known to have had a high blood pressure for several years.

This article will be concluded next week.

REFERENCES

- Ayman, D. (1948). *Arterial Hypertension*. New York.
 Byrom, F. B., and Dodson L. F. (1948). *J. Path. Bact.*, **60**, 357.
 Castleman, B., and Smithwick, R. H. (1943). *J. Amer. med. Ass.*, **121**, 1256.
 Dexter, L., and Haynes, F. W. (1944). *Proc. Soc. exp. Biol., N.Y.*, **55**, 288.
 Goldblatt, H., Lynch, J., Hanzal, R. F., and Summerville, W. W. (1934). *J. exp. Med.*, **59**, 347.
 Platt, R. (1947). *Quart. J. Med.*, **16**, 111.
 Selye, H., Hall, C. E., and Rowley, E. M. (1943). *Canad. med. Ass. J.*, **49**, 88.
 Shorr, E., Zweifach, B. W., and Furchgott, R. F. (1945). *Science*, **102**, 489.
 Wakerlin, G. E. (1949). *Ann. intern. Med.*, **31**, 312.
 Wilson, C., and Byrom, F. B. (1939). *Lancet*, **1**, 136.
 — (1941). *Quart. J. Med.*, **10**, 65.

FORM AND FUNCTION IN THE RENAL TRACT

INAUGURAL MEETING OF RENAL ASSOCIATION

The newly formed Renal Association held its first meeting on March 30 at the Ciba Foundation. Mr. John Sophian was elected secretary, and an executive committee was formed consisting of Mr. F. J. F. Barrington, Professor K. J. Franklin, Mr. D. F. Ellison Nash, and Dr. W. W. Payne. It is hoped to hold quarterly meetings.

Mr. ELLISON NASH gave an address on "Form and Function in the Renal Tract," discussing mainly renal lesions in the child. He first described the embryology of the renal tract, and pointed out that in post-natal life the nephrons continued to form, and the Wolffian duct to grow and to be taken up into the bladder. Gross deviations from the normal were well known, and he mentioned a few of them—the solitary kidney, arrest of the kidney in ascent or in rotation, the double pelvis, the double ureter with one opening, and the ectopic kidney opening outside the bladder, probably into the posterior urethra. The last-named abnormality was one of the commonest causes of enuresis. A child might be labelled a bed-wetter, and it might be some years before the condition was diagnosed. The clinical significance of such a developmental abnormality depended upon the powers of evacuation of the accessory ureter. If it was able to excrete the urine there was not much risk of infection, but if stasis occurred, infection nearly always followed.

Great strides had been made in the understanding of the pathology of these conditions during the last few years; the literature was voluminous and the picture fascinating but confusing. Mr. Nash went on to discuss derangements of function and their causes. He mentioned some recent work suggesting that nicotine stimulated the pituitary in its task of regulating renal flow. The output of urine per pound of body weight fell after the ninth month. Renal function was not mature until the second year. Tubular function in the infant might be extremely poor, and the slightest upset of fluid balance affected the kidneys and prevented a proper clearance of chlorides and urine. In the infant there was only a narrow range of alkali reserve. The kidney could not cope with anything beyond that range, and failure was even more likely if there was any renal damage, which was not uncommon.

The newborn infant's bladder emptied itself automatically at a certain capacity. As the capacity of the bladder increased by natural growth and physiological maturity, the contractions

became inhibited at a higher nervous level; some of the contractions were accompanied by a desire to void and others were not. The majority of children with enuresis had an immature bladder. Psychiatric or psychological causes of enuresis were probably very few and far between. Of the children coming to hospital with enuresis 30% had gross organic disease—ectopic ureter, gross infections, and other conditions. It was unfair offhand to label a case of enuresis "psychological" when there was a one-in-three chance that there was organic disease.

MEDICAL ART SOCIETY EXHIBITION

[FROM A SPECIAL CORRESPONDENT]

Calm festivity is the first impression conveyed by the Medical Art Society's exhibition at Walker's Galleries, Bond Street, London (April 17–29), and the enjoyment with which these medical men and women have evidently filled their canvases is communicated with skill and charm to the visitor. Members of the French Salon des Médecins are also showing some of their work, and reveal an *expertise* that not all their British colleagues attain. Two landscapes catch the eye immediately (17 and 73). Both convey an effortless vigour that exemplifies the Taoist text "Perfect activity leaves no track behind it." In the first, "Swiss Morning," Dr. Gladys Ward depicts snow-clad peaks with the enthusiasm of one who has enjoyed walking among them, and in the other Dr. L. D. Berger, of the Salon, has spread a glistening autumn scene on the canvas with a palette knife.

Landscapes predominate, and Mr. T. Holmes Sellors and Dr. Hugh Stannus depict some characteristically English scenes with charm and skill, while Sir Philip Manson-Bahr, the president of the society, exhibits several tropical scenes. Dr. R. A. Beaver is exceptional in preferring still-life—a type of painting at which professionals are more apt to succeed—but his pictures of flowers (42 and 57) have a harmonious splendour that reveals both freshness of vision and assimilation of the work done since the Impressionists. Portraiture again is elusive for the amateur—indeed, few professionals to-day go further than an expensive likeness—but No. 63, by Dr. André Boissier, is a perceptive study, and No. 4, by Dr. Eric Topham, is expressive if not quite so accomplished.

Most of the work reveals strong Impressionist influence, and when some of the painters have tried something in later vein they have not always managed to avert their gaze from modern poster art. But Dr. A. B. Hewlett has succeeded in doing so with his "Boat House" and "The Tree," in both achieving clarity and vigour without superficiality. More subdued are Sir Henry Bashford's compositions, and his "Champagne" shows a sober tranquillity that must be rarely inspired by that refreshment. All are delicate well-finished works, especially the contemplative "Trout Stream." Mr. Frank Forty has neatly captured in his "Hastings" the wateriness of boats even when beached, and Dr. Francis Roles's carefully painted street scene (26) shows individuality. Nearly all the pictures reveal unexpectedly high accomplishment and convey a holiday spirit—whether of gusto or relaxation—that is a delight to share.

The Report of the Central Council for Health Education for 1948–9 singles out four of its activities for special mention. Improvement of cleanliness in the preparation of food is given first importance because of recent deficiencies in this direction. Education in the principles of diet has been studied, with recollections of the lessons taught by the war, and a campaign on this subject will be made as soon as food supplies improve. Health education in industry has been accepted as an important topic and is to be increased. The Council has been studying the Health Service Act in connexion with the prevention of illness and the aftercare of the sick, which are now the responsibilities of the local health authorities.