idiopathic steatorrhoea. They were young adults of different age and of both sexes (two female and two male), and all had faulty fat absorption as shown by the fat-balance test. Other cases gave a history suggestive of a hereditary element. Thus a man aged 62 who had suffered from steatorrhoea all his life, starting with coeliac disease in infancy, had a father and sister with "chronic diarrhoea."

Bearing the above cases in mind, we investigated the possibility that a genetic factor was of aetiological importance in the sprue syndrome. Our method of investigation was similar to that used previously when inquiring into the relationship of idiopathic steatorrhoea to coeliac disease.

In the first instance the records of approximately 200 cases of coeliac disease were selected and the appropriate questionnaire was sent to the family doctor in charge of the patient. Up to date 130 replies have been received (see Table XI). It will be seen that 9.2% gave a history of steatorrhoea in some members of the family—in most cases it was a brother or sister with coeliac disease—whilst 3.8% had relatives who gave a history of a condition highly suggestive of one or other type of the sprue syndrome.

Similarly, a questionnaire was sent to family practitioners in charge of cases of idiopathic steatorrhoea. In a significant percentage (10.6) of the 75 replies received relatives had definite steatorrhoea or symptoms suggestive of such (see Table XI).

TABLE XI

	No. of Cases	Relative with Steatorrhoea	Relative with Symptoms of Malabsorption Syndrome
Coeliac disease	130	12 (9.2%)	5 (3.8%)
Idiopathic steatorrhoea ..	75	3 (4%)	5 (6.6%)

In both instances the figures as they stand suggest that a genetic factor is involved—probably a recessive gene yielding a diathesis. It should be pointed out, however, that the familial incidence might be related to environmental factors operating on the family. In 8 out of 12 cases of coeliac disease a brother and a sister were affected. In order to exclude an environmental factor it would be necessary to inquire closely into the history of the syndrome in more distant relatives. We consider, however, that enough information of importance has been collected to warrant a thorough study of the possibility that the syndrome is conditioned by a genetic factor which may be accentuated and brought to light by a variety of predisposing environmental agents.

Inquiry from several of our colleagues interested in the sprue syndrome has elicited the information that they also had families with more than two members suffering from steatorrhoea. Thus, Professor Graham, of Glasgow, has given us six family histories in which two or more members of the family suffered from the sprue syndrome. Two similar family histories have come to our notice in Edinburgh and one in Leeds.

Summary and Conclusions

The unique opportunity afforded by priority food rationing has enabled us to study certain features of clinical interest in 2,044 cases of the sprue syndrome, consisting of 1,303 cases of coeliac disease, 403 cases of tropical sprue, 311 cases of idiopathic steatorrhoea, and 27 unclassified cases.

Evidence is produced to suggest that the average number of new cases of coeliac disease and idiopathic steatorrhoea developing annually in Great Britain is approximately 117 and 44 respectively. In both diseases males and females are equally affected, and in 34% of cases of idiopathic steatorrhoea a history

was obtained which suggested that the disease started in childhood.

Information obtained from the Army and the R.A.F. in regard to the incidence of tropical sprue, and discharges from the Services related thereto, clearly shows that the prognosis of tropical sprue as it affected personnel in the 1939-45 war was extremely good.

Lastly, records were obtained of a number of families in whom two or more relatives were affected with idiopathic steatorrhoea or coeliac disease, a finding which suggests that a genetic element may be of great importance in the aetiology of the syndrome. The problem of the aetiology of the sprue syndrome, with especial regard to the genetic factor, will be dealt with in a subsequent communication.

We wish to express our thanks to the Ministry of Food and to the Food Rationing (Special Diets) Advisory Committee of the M.R.C. for permission to undertake this investigation; to various officials of the Ministry of Food in London and Edinburgh, and in particular to Miss Hollingsworth, for the large amount of work undertaken in collecting and classifying food priority certificates; to the Directors-General of the R.A.M.C., R.A.F., and the Indian Army Medical Services for permission to publish the data relating to Service personnel; to Major R. B. Stalbow, R.A.M.C., and Wing Commander G. Williams, R.A.F., who were mainly responsible for providing us with the requisite information regarding Service personnel published in this paper; and to many general practitioners in charge of cases of the sprue syndrome who assisted us in our inquiry into the relationship of coeliac disease to idiopathic steatorrhoea and the family incidence of the sprue syndrome.

REFERENCES

- Davidson, L. S. P., Girdwood, R. H., and Innes, E. M. (1947). *Lancet*, **1**, 511.
Hutchison, Robert (1940). *Ibid.*, **1**, 194.

STEATORRHOEA IN A FAMILY

BY

C. W. BARTLEY, M.D., M.R.C.P.

Medical First Assistant, the London Hospital

Steatorrhoea is not a very uncommon disorder, and has been recognized for many years. Garrod and Hurlley (1913) quote Tulpus in the seventeenth century as being one of the first to describe fatty stools, and credit the earliest use of the term "steatorrhoea" to Kunzmann in 1824, while Gull (1855) described it in association with diseased mesenteric glands. Gee in 1888 published his classical monograph *On the Coeliac Affection*, and subsequently there have been many papers describing the clinical picture of steatorrhoea in both children and adults.

In his monograph Gee remarked that "naked-eye examinations of dead bodies throw no light upon the nature of the coeliac affection," and it is still true that in the great majority of patients dying from steatorrhoea little information about the fundamental pathology can be obtained at necropsy. Such cases occurring in adults have been called idiopathic steatorrhoea, and Bennett, Hunter, and Vaughan (1932) have indicated that in many of these cases the onset could be traced back to childhood or adolescence; and they expressed the view that coeliac disease and idiopathic steatorrhoea were variants only of the same disease. In neither of these two conditions has a familial incidence been noted.

In only a small number of patients with steatorrhoea is any underlying disorder of structure discovered. Cases have been described from time to time associated with chronic relapsing pancreatitis (Dornberger *et al.*, 1948). Other cases have shown involvement of the mesenteric lymphatics and small intestines by new growth (Fairley

and Mackie, 1937), by tuberculosis (Hurst *et al.*, 1942), and by lipophagia granulomatosis (Apperly and Copley, 1943). Increased fat content of the stool has also been observed in cases of gastro-colic fistula and in other acquired abnormalities of the gut (Brown, 1949). No familial incidence has been recorded in any of these conditions.

There remains, however, a small group of patients with steatorrhoea in whom necropsy has revealed a specific fibrosis of the pancreas with cystic dilatation of the acini. These changes were first described by Landsteiner in 1905, and were subsequently reported in association with coeliac disease by Passini (1919), Parmelee (1935), and others.

In 1938 Blackfan and May described 35 well-defined examples of this cystic fibrosis of the pancreas in a series of 2,800 routine necropsies on children. The lungs showed bronchiectasis and terminal bronchopneumonia in all. Five of them had died shortly after birth; others had presented a clinical picture of failure to thrive, with frequent attacks of diarrhoea and repeated pulmonary infections; while seven had presented as coeliac disease and later had developed chronic pulmonary infection. Blackfan and May also described 200 cases in the same series in which smaller degrees of cystic fibrosis of the pancreas were found in children dying of some other disease.

In the same year, Andersen (1938) reviewed 49 cases of this condition, which she called fibrocystic disease of the pancreas. Of these, 22 were fresh examples, 20 of them being her own cases. This last group represented 3.3% of a total of 605 examinations of the pancreas obtained from 1,000 routine necropsies at a children's hospital. She estimated that in every one of her cases 90% or more of the pancreas was involved by the fibrosis. She referred to a further 21 cases encountered in these 605 examinations in which she estimated that between 20% and 90% of the pancreas was involved. Of this group, only three cases presented clinical signs of the disease, the others dying from other conditions. Finally, she described a group of 69 cases in which a few cysts and minimal fibrosis were observed. None of these were affected clinically.

The clinical manifestations of the disease fell into three groups: (1) cases of meconium ileus; (2) cases dying of pulmonary infection before the age of six months; and (3) cases presenting with steatorrhoea, later developing bronchiectasis and dying of bronchopneumonia before the age of 14½ years. By way of comparison she referred to 11 patients with coeliac disease encountered in the same series, of whom none died of pulmonary disease.

In this and subsequent series it has appeared that the pulmonary infection is characteristic of the disease. Farber (1946) considered that the disease affected all mucus-secreting glands, and the loss of normal mucus from the bronchial epithelium accounted for the particular susceptibility to infection. For this reason the mortality is high, and no case has been recorded as living after the age of 14½ years (Matheson, 1949).

A second characteristic of fibrocystic disease of the pancreas is its strong familial incidence. Blackfan and May remarked upon it in their original paper, and Andersen and Hodges (1946) reviewed 20 families affected by the disease and suggested that it might be inherited as a recessive trait. More recently, Matheson (1949) concluded that it might even be inherited as an incomplete dominant. This incidence contrasts strongly with the surprising lack of case reports of idiopathic steatorrhoea occurring in families.

A third characteristic of the disease which is of use in the clinical diagnosis is the very low concentration or

absence of trypsin in the duodenal juice of affected children. Andersen (1942), using the viscosimetric method of assaying trypsin, found that unaffected children, including cases of coeliac disease, gave a concentration of over 100 units per ml., whereas in 15 cases of fibrocystic disease of the pancreas diagnosed clinically or confirmed at necropsy the reading was never higher than 10 units, and in the majority of cases trypsin was absent entirely.

Present Investigation

Below is described a group of five cases of steatorrhoea occurring in the same family (see Diagram), with a brief

Diagram showing the relationship of the various members of the family. Those affected are shown in black.

discussion of the possible aetiology. The affected members included the father, aged 48, his eldest daughter, aged 22, and three younger children. In each case, apart from the purely clinical and routine laboratory investigations, the diagnosis has been confirmed with a three-day fat-balance test as suggested by Cooke *et al.* (1946). In Cases 1 and 2 the test diet contained 50 g. of fat a day, and in the other three 30 g. of fat. The test period began with a saline enema which was discarded, and ended with a saline enema which was collected.

The lower limit of normal has been estimated as 91% absorption, the normal range being 95% $\pm$ 4%. In each case, to exclude any anatomical abnormality, a barium meal and follow-through was carried out. A routine glucose-tolerance test was also performed. For this the fasting blood sugar was estimated, and blood-sugar levels were determined at three-quarters of an hour and at two hours after the ingestion of 50 g. of glucose. Finally, in every case an attempt was made to obtain a specimen of pancreatic juice by duodenal intubation. This was performed under radiological control. Any fluid so obtained was assayed for trypsin by the casein digestion method (Bray, 1946), the normal range being 25 to 50 units of trypsin per ml.

Case 1

The father of the family, a lorry driver aged 48, had always been rather small and under-weight. Between the ages of 14 and 46 he had suffered from recurrent attacks of carpo-pedal spasm, usually in the winter. At the age of 28 he had developed dyspepsia, and at 38 the presence of a duodenal ulcer had been proved radiologically. His bowels have always been opened regularly twice a day, the stools being large, pale, and often putty-like.

On examination he was seen to be thin but not wasted. There was pigmentation of the face, neck, and hands. His height was 5 ft. 4 in. (162.5 cm.) and weight 109 lb. (49.4 kg.). The tongue was completely bald and magenta-coloured, and there was angular stomatitis. The abdomen was not distended, but there was slight epigastric tenderness, and the liver was palpable three fingerbreadths below the costal margin. The heart, lungs, and central nervous system were normal. No tetany was elicited, but there was slight genu varum.

Investigations.—Blood count: Hb, 102%; white cells, 6,700, with a normal differential count. Fractional test meal: normal curve. Plasma proteins: total, 6.6 g.% (albumin 4.2 g., globulin 2.4 g.); all flocculation tests were negative. Serum calcium, 9.5 mg. per 100 ml.; serum phosphorus, 4.5 mg. per 100 ml. Alkaline phosphatase, 6 units (King-Armstrong). A radiograph of the chest was normal, and a barium meal and follow-through, apart from showing a scarred duodenal cap, was normal. A radiograph of the long bones revealed slight osteoporosis. Glucose-tolerance curve: fasting level, 87 mg. per 100 ml.; at $\frac{1}{2}$ hour, 118 mg.; at 2 hours, 143 mg. Three-day fat-balance test: 85% absorption. Duodenal aspiration failed owing to the scarring of the duodenal cap.

The diagnosis of steatorrhoea was suggested by the history of recurrent attacks of tetany and by the appearance of the stools. On examination there was evidence of a low-grade avitaminosis and a flat glucose-tolerance curve. The barium meal revealed scarring of the duodenal cap but no delay in emptying. The fat-balance test showed that he was suffering from a low-grade failure of fat absorption.

Case 2

The eldest daughter, aged 22, had been seen to be pale six years previously, and two years later she had begun to lose her appetite and had become apathetic. An iron tonic had been prescribed by her doctor, and she had taken it regularly ever since. Two years ago spooning and brittleness of her finger-nails occurred. Her bowels have always been opened twice daily, the stools being of normal consistency, although occasionally light yellow in colour. Diarrhoea occurred if she ate fruit. The catamenia were scanty, but her growth and development were normal.

On examination the skin, physical development, and distribution of fat were seen to be normal. There was pronounced pallor of the mucous membranes and gross koilonychia, but no jaundice. The tongue was smooth and atrophic, and there was slight angular stomatitis. The abdomen was not distended, but the spleen was palpable two fingerbreadths below the costal margin. There was a soft, blowing mitral systolic murmur. The lungs and central nervous system were normal.

Investigations.—Blood count: Hb, 41%; red cells, 4,000,000; C.I., 0.52; M.C.V., 72 μ^3 , with anisocytosis, poikilocytosis, and slight polychromasia; reticulocytes, 0.8%; white cells, 6,900, with a normal differential count. Fractional test meal: normal curve. Sternal marrow: normoblastic hyperplasia. Plasma proteins: total, 6.4 g.% (albumin 4.3 g., globulin 2.1 g.). Serum calcium, 8.5 mg. per 100 ml.; serum phosphorus, 3.8 mg. per 100 ml. A radiograph of the long bones showed some osteoporosis. X-ray examination of the chest and a barium meal and follow-through were normal. Glucose-tolerance curve: fasting, 100 mg. per 100 ml.; at $\frac{1}{2}$ hour, 125 mg.; at two hours, 100 mg. Three-day fat-balance test: 74% absorption. Duodenal aspiration: 25 units of trypsin per ml. (normal, 25 to 50 units).

This case presented as an idiopathic hypochromic anaemia which either was iron-resistant or had been inadequately treated. A trial course of ferrous sulphate, 15 gr. (1 g.) four times a day for three weeks, while in hospital failed to improve the condition. It responded later to a course of saccharated iron administered intravenously. The presence of osteoporosis suggested the diagnosis of steatorrhoea, and the fat-balance test gave unequivocal confirmation of this in spite of the fairly normal appearance of the stool.

Case 3

A son aged 14 $\frac{1}{2}$ had recurrent attacks of diarrhoea for nine years: each lasted two to three weeks, during which his bowels were opened four times a day, with very bulky clay-coloured but inoffensive stools. The attacks were usually precipitated by excessive fats in the food. In the last year he has had no attacks, but the stools, although formed, have remained rather pale and greasy. Except for some "weakness" he has been well.

On examination he appeared healthy, though rather small for his age. His height was 4 ft. 7 in. (140 cm.), and weight 69 lb. (31.3 kg.)—normal average height 5 ft. 1 in. (155 cm.), weight 108 lb. (49 kg.). The mucosae were pale, and the genital development was prepubertal. Otherwise his heart, lungs, abdomen, and central nervous system were normal.

Investigations.—Blood count: Hb, 67%; red cells, 5,300,000; C.I., 0.63; M.C.V., 78 μ^3 ; white cells normal. Fractional test meal: normal curve. Sternal marrow: normoblastic erythropoiesis. Plasma proteins: total, 6.4 g.% (albumin 3.8 g., globulin 2.6 g.). Serum calcium, 7.3 mg. per 100 ml.; serum phosphorus, 4.3 mg. per 100 ml. Alkaline phosphatase, 12 units (King-Armstrong). A radiograph of his chest and a barium meal and follow-through were normal. Glucose-tolerance curve: fasting, 143 mg. per 100 ml.; at $\frac{1}{2}$ hour, 145 mg.; at two hours, 154 mg. Three-day fat-balance test: 83% absorption. Duodenal aspiration: no trypsin detected.

The history of the diarrhoea and the presence of a hypochromic anaemia were suggestive of steatorrhoea, but no other gross stigmata could be found clinically. His serum calcium was low and the glucose-tolerance curve flat. The fat-balance test supplied evidence of moderate failure of fat absorption, and confirmed the diagnosis.

Case 4

A daughter aged 11 has had continuous diarrhoea for six years, her bowels being opened three times a day. The stools were of moderate size, and partly formed, but noticeably pale and fatty. Her appetite was good, but she was intolerant of fats. She has always been very small in stature for her age. As an infant she had had two generalized convulsions.

On examination she appeared infantile. Her height was 3 ft. 6 in. (107 cm.) and weight 46 lb. (20.9 kg.)—minimum normal height for age is 4 ft. (122 cm.) and minimum weight 53 lb. (24 kg.). There was wasting of the buttocks, with slight genu valgum and bowing of the tibiae, and the fingers were clubbed. There was angular stomatitis, and the mucosae were pale. The abdomen was distended and tympanitic. The heart and lungs were normal. There was a positive Chvostek sign, otherwise the central nervous system was normal.

Investigations.—Blood count: Hb, 43%; red cells, 5,000,000; C.I., 0.43; white cells, 5,350—differential count normal. Fractional test meal: normal curve. Sternal marrow: normoblastic erythropoiesis. Plasma proteins: total, 5.5 g.% (albumin 3.6 g., globulin 1.9 g.). Serum calcium, 10 mg. per 100 ml.; serum phosphorus, 3.3 mg. per 100 ml. A radiograph of the chest and a barium meal and follow-through were normal. A radiograph of the long bones showed rachitic changes, osteoporosis, and retardation of development. Glucose-tolerance curve: fasting, 178 mg. per 100 ml.; at $\frac{1}{2}$ hour, 194 mg.; at two hours, 178 mg. Three-day fat-balance test: 74% absorption. Duodenal aspiration on two occasions: no trypsin detected.

On clinical grounds there was no doubt that this child was suffering from steatorrhoea. The fat-balance test was used to assess the severity of the condition.

Case 5

A son aged 9 has had "diarrhoea" for four years; his bowels were opened two to four times a day, the stools being formed but fatty, clay-coloured, and slightly bulky. He was "upset" by fats and fruits, and had always been pale, rather listless, and underdeveloped, with a protuberant abdomen. His appetite was good. Four weeks before admission he developed a rash. He had had three or four fits as an infant, and one attack of bronchopneumonia.

On examination his growth was seen to be moderately retarded, his height being 3 ft. 7 in. (109 cm.) and his weight 43 lb. (19.5 kg.)—normal average height 4 ft. 4 in. (132 cm.) and weight 69 lb. (31.3 kg.). There was slight wasting of the buttocks and bowing of the tibiae. The skin showed patchy, red, raised, desquamating lesions between $\frac{1}{4}$ in. and 2 in. (0.6 and 5 cm.) in diameter over arms and legs and left pinna; with one patch of hyperkeratinization, 3 by 2 in. (7.5 by 5 cm.).

on the anterior surface of one shin. There was angular stomatitis, and the abdomen was distended and tympanitic. The heart, lungs, and central nervous system were normal.

Investigations.—Blood count: Hb, 48%; red cells, 4,900,000; C.I., 0.5; M.C.V., 70 μ^3 ; white cells normal. Fractional test meal: normal curve. Plasma proteins: total, 6 g.% (albumin 4.2 g., globulin 1.8 g.). Serum calcium, 8 mg. per 100 ml.; serum phosphorus, 4.8 mg. per 100 ml.; alkaline phosphatase, 4 units (King-Armstrong). A radiograph of his chest and a barium meal and follow-through were normal. A radiograph of his long bones showed delayed ossification. Glucose-tolerance curve: fasting, 125 mg.; at $\frac{1}{2}$ hour, 182 mg.; at two hours, 143 mg. Three-day fat-balance test: 86% absorption. Duodenal aspiration: no specimen obtained.

The general impression this child gave was that of a "near coeliac," as Andersen termed it. The history and general appearance suggested steatorrhoea, and there was clinical evidence of a low-grade avitaminosis, a considerable anaemia, and slight hypocalcaemia. Again the fat-balance test helped to confirm the diagnosis.

Discussion

Attention was at first drawn to this family by the discovery of anaemia in four of the children. The obvious appearance of coeliac disease in one of them, and the suggestive histories of the others, prompted a more thorough investigation. The three-day fat-balance test provided an apparently useful and reliable method of confirming the diagnosis in every case.

The aetiology of this familial failure of fat absorption seemed somewhat obscure. It is almost axiomatic that idiopathic steatorrhoea is not a familial disease. On the other hand, there is a strong familial incidence of fibrocystic disease of the pancreas, and this disorder may present as a steatorrhoea indistinguishable from coeliac disease. However, in the cases of fibrocystic disease of the pancreas which have been described there is no record that any of them lived beyond the age of 14½ years (Wiglesworth, 1946; Matheson, 1949). Furthermore, pulmonary infections have been so constant a finding in this condition that Farber (1946) and Bodian (1946) think they are a manifestation of the disease process itself, and not merely, as Blackfan and May (1938) suggested, a result of dietary insufficiency. On the other hand, Andersen (1949) is still of this latter opinion, and has stated that, with adequate chemotherapy and a correct dietary regime, cases under observation have remained free of respiratory infections for many years.

There are further points which seem worthy of consideration. The original papers of Blackfan and May and of Andersen indicated that practically all degrees of involvement of the pancreas by the fibrocystic changes might be encountered pathologically in routine necropsies on children. The pathological findings have also suggested that it is only when pronounced changes are so encountered that clinical features are recognized. In Andersen's paper the degree of pancreatic involvement has been related to the number of cases presenting signs. Where it was estimated that 90% or more of the pancreas was involved every case showed clinical features. In the group in which between 20% and 90% was involved only three cases out of a total of 21 examples manifested the disease. In the final group of 69 cases, in which it was estimated that less than 20% of the pancreas was involved, all the children had died from purely incidental causes. It is not unreasonable to suppose that other children with a similar degree of minimal fibrosis do not die from incidental causes and might survive until adult life. In support, Hellerstein (1946) records this pathological change as an incidental finding at necropsy on an adult.

It was in an endeavour to discover clinically whether there was some impairment of pancreatic function in this family that duodenal intubation was attempted in every case. In Case 1, in spite of the use of a variety of methods, the deformed duodenal cap seemed to present an impassable barrier, and in Case 5 no specimen could be obtained owing to lack of co-operation of the patient. In two further cases (Nos. 3 and 4) adequate specimens were obtained under strict radiological control, but no trypsin could be detected in either. These findings might be interpreted as confirming the diagnosis of fibrocystic disease of the pancreas in those two cases. From the remaining case (No. 2) a specimen was obtained which showed a concentration of trypsin that was just within the lower limit of the estimated normal range. This patient had survived until the age of 22, and, except for the disturbance of iron and fat absorption, was in good health. Therefore it might be surmised that the degree of fibrosis of her pancreas was only moderate—in which case trypsin might be expected to be present in the duodenal juice in moderate quantities.

In conclusion it seems hardly necessary to emphasize the fact that with the evidence available it is impossible to be dogmatic concerning the aetiology of the steatorrhoea occurring in this family. Fibrocystic disease of the pancreas seems to be the only condition producing steatorrhoea that has a pronounced familial incidence, and there is circumstantial evidence that it may occur in adults. It is suggested that the severity of the condition may vary with the extent of the pancreatic involvement.

Summary

An outline of the development of our knowledge of the aetiology of steatorrhoea is presented.

Reference is made to the finding of fibrocystic disease of the pancreas in children at routine necropsy. Its clinical manifestations and its marked familial incidence are noted.

Five cases of steatorrhoea, including two adult cases, occurring in one family are described, and the possible aetiology is discussed briefly.

My thanks are due to Dr. K. M. A. Perry, under whose care the patients were admitted to hospital, for his permission to publish the case reports, and for his help and encouragement. I also wish to thank Dr. J. R. Tasker and Dr. H. C. Hamilton for their invaluable aid throughout, and for much constructive criticism.

REFERENCES

- Andersen, D. H. (1938). *Amer. J. Dis. Child.*, **56**, 344.
 — (1942). *Ibid.*, **63**, 643.
 — (1949). *Proc. R. Soc. Med.*, **42**, 25.
 — and Hodges, R. G. (1946). *Amer. J. Dis. Child.*, **72**, 62.
 Apperly, F. L., and Copley, E. L. (1943). *Gastroenterology*, **1**, 461.
 Bennett, T. I., Hunter, D., and Vaughan, J. M. (1932). *Quart. J. Med.*, n.s. **1**, 603.
 Blackfan, K. D., and May, C. D. (1938). *J. Pediat.*, **13**, 627.
 Bodian, M. (1946). *Arch. Dis. Child.*, **21**, 179.
 Bray, W. E. (1946). *Synopsis of Clinical Laboratory Methods*. C. V. Mosby Co., St. Louis.
 Brown, A. (1949). *British Medical Journal*, **1**, 1073.
 Cooke, W. E., Elkes, J. J., Frazer, A. C., Parkes, J., Peeney, A. L. P., Sammons, H. G., and Thomas, G. (1946). *Quart. J. Med.*, **15**, 141.
 Dornberger, G. R., Comfort, M. W., Wollaeger, E. E., and Power, M. H. (1948). *Gastroenterology*, **11**, 691.
 Fairley, N. H., and Mackie, F. P. (1937). *British Medical Journal*, **1**, 375.
 Farber, S. (1946). *Arch. Path.*, **37**, 238.
 Garrod, A. E., and Hurlley, W. H. (1913). *Quart. J. Med.*, **6**, 242.
 Gee, S. (1888). *St. Bart's Hosp. Rep.*, **24**, 17.
 Gull, W. (1855). *Guy's Hosp. Rep.*, **38**, 1, 369.
 Hellerstein, H. K. (1946). *Ohio St. med. J.*, **42**, 616.
 Hurst, A., Wright, G. P., and Ryle, J. A. (1942). *Guy's Hosp. Rep.*, **91**, 25.
 Landsteiner, K. (1905). *Zbl. allg. Path. path. Anat.*, **16**, 903.
 Matheson, W. J. (1949). *British Medical Journal*, **2**, 206.
 Parmelee, A. H. (1935). *Amer. J. Dis. Child.*, **50**, 1418.
 Passini, F. (1919). *Dtsch. med. Wschr.*, **45**, 851.
 Wiglesworth, F. W. (1946). *Amer. J. med. Sci.*, **212**, 351.