ulceration are among the most obvious toxic signs, and their development, presumably due to interference with the growth of the lining epithelial cells, is an indication to withhold the drug. Haemorrhage is a special and serious problem and repeated blood transfusions may be necessary, but may prove ineffective. The cause of the bleeding is obscure; it has been claimed in thrombocytopenic purpura and in untreated leukaemia that the amount of heparin-like substances in the blood is increased (Allen, Bogardus, Jacobson, and Spurr, 1947). Whether additional changes occur as the result of giving aminopterin, other than the depression of platelet numbers, is uncertain.

Indications and Limitations

In view of the temporary nature of the remissions and the ultimately disappointing results of the use of folic-acid antagonists in human acute leukaemia, and of the very small therapeutic margin and extreme toxicity of the drugs, it may well be asked what, if any, should be the role of aminopterin or allied compounds in the treatment of acute leukaemia. Although our experience is small it has been large enough for initial enthusiasm to be tempered and for us to realize the serious limitations of our attempts at treatment. It is impossible to be enthusiastic; the hopes of patients, parents, and doctors are raised only to be dashed by later experience.

Nevertheless, that remission can be observed at all is remarkable, and it seems difficult to withhold the drugs, should they be available, just because relapse will inevitably follow remission, any more than transfusions can be withheld just because they also are temporary expedients. It is always possible, if improbable, that a particular patient may have a prolonged and really worth-while remission, and further experience and different schemes of dosages may produce better results.

Summary

The results of treatment of 13 cases of acute and subacute leukaemia in children and adults with the folic-acid antagonist aminopterin are reported.

Clinical and haematological remissions were obtained in all of seven children treated, and in two of six adults. In those cases which responded there was a clinical return to normality and the peripheral blood was rendered normal, or almost normal, for varying periods. Although striking improvements were obtained in the bone marrows, leukaemic cells were never completely eradicated from them.

All patients who originally benefited have relapsed, and have eventually died after becoming more or less refractory to treatment.

The action of aminopterin in any given case is unpredictable and the toxic effects of the drug are severe. Great caution is necessary in its use.

We are indebted to Professors A. Haddow and E. Boyland, of the Chester Beatty Institute, for supplies of aminopterin for the early cases, and to Lederle Laboratories Division, Cyanamid Products, Ltd., for further supplies. We are most grateful to the staffs of the departments of medicine and paediatrics for allowing us to treat patients under their care. The photomicrographs were prepared by Mr. E. V. Willmott.

REFERENCES

- Allen, J. G., Bogardus, G., Jacobson, L. O., and Spurr, C. L. (1947). *Ann. intern. Med.*, **27**, 382.
- Angier, R. B., et al. (1945). *Science*, **102**, 227.
- Berman, L., Axelrod, A. R., Heide, E. C. V., and Sharpe, E. A. (1949). *Amer. J. clin. Path.*, **19**, 127.
- Birge, R. F., Jenks, A. L., and Davis, S. K. (1949). *J. Amer. med. Ass.*, **140**, 589.
- Dameshek, W. (1948). *Blood*, **3**, 1057.
- (1949). *Ibid.*, **4**, 168.
- Farber, S. (1949). *Ibid.*, **4**, 160.
- Diamond, L. K., Mercer, R. D., Sylvester, R. F., and Wolff, J. A. (1948). *New Engl. J. Med.*, **238**, 787.
- Fildes, P., and Rydon, H. N. (1947). *Brit. J. exp. Path.*, **28**, 211.
- Karnofsky, D. A., Patterson, P. A., and Ridgway, L. P. (1949). *Proc. Soc. exp. Biol., N.Y.*, **71**, 447.
- Meyer, L. M. (1948). *Trans. N.Y. Acad. Sci.*, **10**, 99.
- Fink, H., Sawitsky, A., Rowen, M., and Ritz, N. D. (1949). *Amer. J. clin. Path.*, **19**, 119.
- Pierce, M., and Alt, H. (1948). *Proc. cent. Soc. clin. Res.*, **21**, 89.
- Rogers, L. L., and Shive, W. (1948). *J. biol. Chem.*, **172**, 751.
- Seeger, D. R., Smith, J. M., and Hultquist, M. E. (1947). *J. Amer. chem. Soc.*, **69**, 2567.
- Smith, J. M., Cosulich, D. B., Hultquist, E., and Seeger, D. R. (1948). *Trans. N.Y. Acad. Sci.*, **10**, 82.
- Stickney, J. M., Hagedorn, A. B., Mills, S. D., and Cooper, T. (1948). *Proc. cent. Soc. clin. Res.*, **21**, 31.
- Stokstad, E. L. R., Pierce, J., Page, A. C., jun., Hoffman, C. E., Franklin, A. L., and Jukes, T. H. (1949). 1st Int. Cong. Biochem., Abstr. No. 415/1.
- Regan, M., Franklin, A. L., and Jukes, T. H. (1948). *Fed. Proc.*, **7**, 193.
- Thiersch, J. B. (1949). *Cancer*, **2**, 877.
- and Philips, F. S. (1949). *Amer. J. med. Sci.*, **217**, 575.
- Whitby, L. E. H., and Christie, J. M. (1935). *Lancet*, **1**, 80.
- Wolman, I. J., Eglick, P., Dickstein, B., and Githens, J. (1949). *Penn. med. J.*, **52**, 474.
- Woolley, D. W. (1947). *Physiol. Rev.*, **27**, 308.

EXTERNAL ILIAC ARTERY THROMBOSIS

BY

A. M. BOYD, F.R.C.S.

Professor of Surgery

AND

R. P. JEPSON, F.R.C.S.

Lecturer in Surgery

The Professorial Surgical Unit, Royal Infirmary, Manchester

[WITH PHOTOGRAPHURE PLATE]

On attendance at the neurovascular clinic at the Manchester Royal Infirmary, patients with the symptoms of intermittent claudication undergo a routine clinical examination which enables the underlying vascular disease to be assessed (Boyd *et al.*, 1949). In several cases the examination failed to reveal any underlying pathological condition. Subsequent investigation showed that in two cases the lesion was a unilateral thrombosis of the external iliac artery. As a search of the literature has not produced any description of such a lesion and its sequelae, a detailed account of these two cases is given. No discussion is made of other and now well-recognized causes of "proximal" arterial block such as the aortic bifurcation syndrome (Leriche and Morel, 1948).

Case 1

Whilst felling a tree on December 15, 1947, a man aged 28 was struck across the lower part of the abdomen and thigh by a large branch, causing extensive bruising. Subsequent x-ray examination did not show any bony injury. The accident occurred at 10 a.m., and at 1 p.m. his right foot felt numb and colder than the left. The sensation spread upwards to the knee within the next hour, but after about twelve hours it gradually receded. The patient noticed no weakness or colour change in the right leg. After ten days in hospital the contusions resolved and he was discharged.

When he resumed his ordinary activities he found that on walking about 100 yards (91 metres) a deep sickening ache was felt in the right calf and, to a less degree, in the right thigh: this disappeared after a short rest. He noticed nothing else abnormal apart from a vague sense of weakness in the leg; his general health was excellent and he continued to work as a market gardener. Because of the pain in the leg he attended the local hospital as an out-patient and was given an extensive course of short-wave diathermy and massage—without relief. He was referred to the neurovascular clinic on February 1, 1949.

The patient's general condition was that of a healthy young man; nothing abnormal was detected on systemic examination; the blood pressure was 150/90 mm. Hg. On inspection the legs

appeared normal; measurement showed slight wasting of the right thigh, the mid-thigh circumference being 17½ in. (44.5 cm.) on the right and 18½ in. (47.6 cm.) on the left. Oscillometer readings were normal on the left leg but much reduced on the right; the arterial pulses were palpable in the left leg, but though the common iliac and femoral arterial pulses could be detected on the right side the popliteal and tibial could not. The temperature was the same in both feet.

An exercise-tolerance test was carried out on the "claudicometer"—a motor-driven moving platform fitted with an automatic recorder measuring the speed of walking and the distance traversed. The patient walked at 2½ miles (4 km.) an hour; he first felt pain in the right calf and thigh at 120 yards (110 metres) and he halted at 300 yards (274 metres). On examination immediately afterwards the right foot was paler than the left; no pulsations or oscillations were detectable in the right leg, but those in the left leg were increased from the resting level. Decrease in oscillations after exercise—the "inverse reaction"—has been demonstrated to occur below a main arterial block (Ejrup, 1948).

From the accumulated evidence the diagnosis of a traumatic thrombosis of the right external iliac artery was made.

Operation.—On February 4, 1949, an arteriectomy was performed. The right external iliac artery, which was about 3 mm. in diameter, was found to be thrombosed from the division of the common iliac artery to the level of Poupart's ligament. The thrombosed segment was removed; it dissected freely from its bed without any evident periarteritis. There were no signs of thrombosis in the right internal iliac artery, which appeared to be larger than normal. Neither the internal nor the external iliac vein was affected. A right lumbar sympathectomy was then carried out, the first three lumbar ganglia and the intervening chain being removed.

Histology.—The histological report on the resected segment of artery was as follows: "The artery is considerably smaller than normal. The internal and external elastic lamellae are abnormally folded. The lumen is filled with incompletely organized thrombus. There is no evidence of active arteritis, periarteritis, or arteriosclerosis."

Post-operative Review.—After a period of uneventful convalescence the patient returned to work. His progress was reviewed on February 5, 1950, twelve months after the operation. The patient said he was able to walk as far as he wished—on one occasion 15 miles (24 km.)—with no discomfort beyond, rarely, the mildest of aches in the right calf. On an exercise-tolerance test he walked at 4 miles (6.4 km.) an hour for a distance of one mile (1.6 km.). A slight paling of the right foot was seen after the test. Table I gives a summary

TABLE I

	Right		Left	
	At Rest	After Exercise	At Rest	After Exercise
Pulses:				
Femoral	+	+	+++	+++
Popliteal	—	+	++	++
Dorsalis pedis ..	—	+	+	++
Posterior tibial ..	—	+	+	++
Oscillations (scale divisions):				
Thigh	1	1	8	9
Calf	3	1	6	7
Supramalleolar ..	1½	—	4	5

of the observations made on the arterial pulses and oscillations at rest and after exercise. These observations are indicative of an established collateral circulation, the apparent discrepancy of the oscillometer readings being due to the phase relationship of the pulse waves.

Case 2

This patient, a man aged 23, gave no history of any definite traumatic incident, but explained that his hobby was sprint cycling. In August, 1944, after walking a distance estimated at

less than 100 yards (91 metres), his left thigh and calf developed an ache, which quickly passed off when he rested. He did not remember there being any coldness or numbness in the left leg or foot. He had found also at this time that when he was cycling the left leg quickly tired and the right leg was forced to do most of the effort of pedalling. The claudication onset distance had gradually increased, and when he was referred to the neurovascular clinic on April 14, 1949, he suffered discomfort only when he walked at a fast pace or during a cycle race, the factor which had caused him to seek medical advice. There was no relevant previous medical history.

On general examination nothing abnormal was detected; the blood pressure was 130/80 mm. Hg. The nutrition of

FIG. A.—Diagrammatic arteriogram of Case 2.

the legs was excellent, but there was slight wasting of the left leg, the thigh being ¼ in. (1.9 cm.) and the calf 1 in. (2.5 cm.) less in circumference than on the right side. Oscillometer readings were normal on the right leg but reduced on the left; the arterial pulses were palpable on both legs; the femoral pulse, however, was less marked on the left side.

On the exercise-tolerance test the patient walked at 3 miles (4.8 km.) an hour for a distance of one mile (1.6 km.), at which point he complained of a feeling of discomfort in the left thigh. The left foot was seen to be slightly paler than the right. Oscillometer readings were increased on the right leg and still further reduced on the left side; the arterial pulses were scarcely palpable on the left side. A left external iliac artery thrombosis was diagnosed.

Operation.—An arteriectomy was performed on April 19, 1949. A thrombus was found extending from the division

of the left common iliac artery to a point $\frac{1}{2}$ in. (1.25 cm.) below Poupart's ligament, thus involving the whole of the external iliac and upper part of the common femoral arteries. The internal iliac, external pudental, superficial epigastric, and external circumflex iliac arteries on the left side were hypertrophied but otherwise appeared normal. An arteriogram taken after the injection of 10 ml. of thorotrast into the common iliac artery showed the excellence of the collateral circulation (Fig. A). The thrombosed segment dissected free without difficulty—there being no periarterial reaction—and was excised.

A left lumbar sympathectomy was also undertaken, the first three lumbar ganglia and intervening chain being removed.

Histology.—The histological report on the excised segment of artery read as follows: "The artery is much decreased in size, and in consequence the internal and external elastic lamellae are tortuous and thickened. There is a slight increase in the amount of medial elastic tissue. The lumen is filled with fibrous tissue which has incorporated the intima. In this fibrous tissue are a few blood spaces which are lined by a simple epithelium. There is no evidence of any inflammatory reaction. There is nothing to suggest the cause for this now quiescent thrombosis" (Plate, Figs. 1, 2, and 3).

Post-operative Review.—A review of the patient's condition was made four months after operation, and the original tests were repeated. The intervening period was too short for the final results of the operation to be assessed. On the exercise-tolerance test the patient walked at 3 miles an hour for a distance of one mile without discomfort. The left foot did not appear pale. The observations are summarized in Table II.

TABLE II

	Left	
	At Rest	After Exercise
Pulses:		
Femoral	+	+
Popliteal	+	+
Dorsalis pedis	++	++
Posterior tibial	+	++
Oscillations (scale divisions):		
Thigh	1	< $\frac{1}{2}$
Calf	4	< $\frac{1}{2}$
Supramalleolar	2	< $\frac{1}{2}$

Literature

The occurrence of intermittent claudication with no evidence of arterial disease has been thought to be due to arterial spasm. Simmons (1936) states: "A proportion of cases of intermittent claudication are due to pure spasm of the arteries in neuropathic individuals, and no structural disease of the vessels is demonstrable either at the time or in later years." Such cases have also been described under the term "paradoxical spasm."

Pearl (1937) reported six cases under the heading of "angiospastic claudication." When these patients took exercise the foot on the affected side became pale and the dorsalis pedis pulse ceased to be palpable. These phenomena did not occur after a lumbar sympathetic block with procaine; this fact was regarded as evidence of arterial spasm.

The possible existence of a high arterial block does not seem to have been considered by these authors, and in consequence their cases were not investigated from this point of view.

A case of "spastic claudication" was described by Lindqvist (1945a). Further investigation showed this to be due to "an organic constriction of the lower part of the aorta" (Lindqvist, 1945b).

Some doubt is felt with regard to the existence of "angiospastic claudication" as a separate entity. Diagnosis

of such a condition is considered unjustifiable until a block proximal to the femoral artery has been excluded.

Clinical Features

The presenting symptom in both of our cases was intermittent claudication. During exercise the patients noticed vague discomfort in the calf and thigh followed by typical claudication pain, relieved by a short rest. The degree of claudication, as usual, depended on the balance between blood supply and demand, becoming less severe as the collateral circulation developed.

The arterial pulses of the limb were diminished, though not to the same degree in the two patients. In Case 1 only the femoral pulse was present; in Case 2, with less severe claudication, all the pulses were present but the femoral pulse was diminished. After exercise the arterial pulses of the affected limb disappeared completely in Case 1 and almost completely in Case 2.

Oscillometric readings were taken while the patients were resting. In comparison with those of the normal side the readings on the side of the lesion were much reduced in Case 1 and moderately so in Case 2. After the patients had walked a sufficient distance to produce the claudication pain it was found that the oscillometric readings were increased on the normal leg and still further reduced on the painful leg. The foot on the painful side also became paler.

Other investigations revealed no significant changes. Temperature gradients and electrical skin-resistance readings were normal; straight radiographs of the abdomen, pelvis, and limbs showed no vascular calcification; the clotting time and platelet counts were within normal limits.

The diagnostic features which must be emphasized are the disappearance or marked diminution of pulses and oscillometric readings after exercise—Ejrup's (1948) "inverse reaction"—followed by a rapid return to normal on resting.

Possible Causes of the Arterial Thrombosis

Arterial Disease

Segmental thrombosis of a main vessel in young men is generally regarded as due to "thrombo-angiitis obliterans." The onset of this condition is usually acral, and a history of phlebitis is common. In older subjects thrombosis of a main vessel is a frequent complication of arteriosclerosis. Of the patients with this condition seen in our clinic approximately 50% have a thrombosis of one or both popliteal arteries and 10% of one or both femoral arteries. The histological findings and the lack of progression in the case histories quoted is thought to exclude either of these diseases.

Idiopathic Thrombosis

Nygaard and Brown (1937) have described five cases with "non-reactive" thrombosis of the main limb vessels under the term "essential thrombophilia." In addition to the limb vessels there was more widespread arterial thrombosis in the cerebral, renal, and coronary vessels; a deep or superficial venous thrombosis was also found. Four cases showed an increase in the coagulability of the plasma.

Examples of a "bland arterial thrombosis" have been reported by Leriche and Stricker (1933) under the heading of "spontaneous localized monarteritis of indeterminate origin." Learmonth, Blackwood, and Richards (1944) have described a similar condition by the term "localized arterial thrombosis of indeterminate origin."

Trauma

Arterial thrombosis is sometimes seen in healthy arteries after severe injury—"run-over" accidents or heavy objects falling across the limb. The accompanying vein is usually thrombosed; fractures are common. An extensive diffuse arterial haematoma, resulting in a tense swelling of the limb and distal ischaemia, is often encountered. Thrombosis of a healthy artery by recurrent crutch pressure has been reported by Platt (1930). Vessels already damaged by arteriosclerosis may be thrombosed by a lesser degree of trauma.

The most common cause of arterial damage resulting in thrombosis is indirect trauma. The site is either where the artery is in close relation to a joint or where it lies between bone and the tendinous insertion of a muscle—for example, the popliteal artery where it enters the popliteal tunnel at the level of the upper border of the femoral condyle, or the subclavian artery as it lies on the first rib behind the tendon of the scalenus anticus. The pathology of the lesion is not clear, but contusion of the arterial wall causing rupture of the vasa vasorum and an intramural haematoma is a possibility. A similar lesion or an intimal tear may be due to a "stretch-strain" or may be the result of traction on a muscular branch.

The popliteal artery is the most common site of thrombosis due to indirect trauma. Boyd and Wilde (1950) have shown that this artery is firmly fixed to the oblique ligament of the knee-joint. The artery enters a fibromuscular canal at the level of the upper border of the femoral condyles, emerging 2 cm. below the knee-joint. During its passage through the "tunnel" the artery may be damaged by contraction of the surrounding muscles or by stretching from the attachment to the oblique ligament on repeated flexion of the knee—especially when fixed to the "tunnel" walls through inflammatory changes in the accompanying vein or lymphatic vessel. Stretch-strain of the artery at the level of the upper border of the femoral condyles occurs with sprains of the knee-joint of the type producing internal derangement. Arteriograms show that the prethrombotic lesion is always at this level, where the artery enters the popliteal canal.

The external iliac artery has been shown to be fixed to the ilio-psoas muscle at the level of Poupart's ligament by the fusion of the iliac and transversalis fascia with the adventitia of the artery (Panton, 1923).

The causes of the thrombosis in the above two cases would seem to be direct trauma in Case 1, who was struck by the falling tree, and indirect trauma due to stretch-strain from repeated flexion of the hip-joint in Case 2, the sprint cyclist.

Treatment

The important collateral circulation after the occlusion of the external iliac artery is by way of the gluteal and obturator arteries and the spinous and cruciate anastomoses. The afferent vessels to these anastomoses arise from the internal iliac and occasionally the common iliac arteries, and receive sympathetic innervation from the lumbar sympathetic chain through the somatic nerve bundles. The common iliac artery is also innervated from the presacral plexus. A lumbar sympathectomy therefore denervates not only the limb vascular bed but also the main collateral vessels.

The incision for the lumbar sympathetic chain is made as small as possible in order to avoid excessive damage to the loin collateral vessels. It is also made a little lower than the standard extraperitoneal loin incision so that the iliac vessels may be more easily exposed. More bleeding

than usual is experienced during the muscle section, and careful haemostasis must be secured. The lumbar chain from L1 to L3 is removed, and the diaphragmatic crus is split to ensure section of the highest lumbar ganglion. Only by the removal of L1 ganglion can the necessary high thigh denervation be obtained (Ratcliffe and Jepson, 1950).

In each of the cases quoted a segment of thrombosed external iliac artery was removed. This was done primarily for histological verification, but there is also reason to believe that removal of the thrombotic zone may discourage further spread. Leriche has suggested that the thrombosed artery initiates reflex vasospasm, and hence should, for this reason alone, be excised.

Prognosis

In both cases, at the post-operative review, exercise caused the limb pulses to become palpably stronger, whereas pre-operatively the converse had been observed.

Subjectively, both patients considered their condition to be much improved. The follow-up period cannot, however, be taken as sufficient for judgment of the long-term results.

If the analogy between primary popliteal thrombosis and these cases is just, it may be assumed that the prognosis is excellent. With increasing age, however, the effect of arteriosclerosis may be more severe in the leg in which the circulation is already impaired by the occlusion of the main artery.

Summary

Two cases of thrombosis of the external iliac artery are reported.

The syndrome of a "high arterial block," as illustrated by these cases, is described. Stress is laid on the "inverse reaction" of the oscillometric readings following exercise.

The possible aetiology of the condition is discussed and reasons for a traumatic origin are given.

Lumbar sympathectomy was performed on these cases and details of the post-operative improvement are given.

Our thanks are due to Dr. A. Hall Ratcliffe for his help with the script, to Miss Dorothy Davison for the diagram, and to Messrs. Ward and Kilshaw for the photographs.

REFERENCES

- Boyd, A. M., et al. (1949). *J. Bone Jt Surg.*, **31B**, 325.
 — and Wilde, F. R. (1950). In press.
 Ejrup, B. (1948). *Acta med. scand.*, Suppl. 211.
 Learmonth, J. R., Blackwood, W., and Richards, R. L. (1944). *Edinb. med. J.*, **51**, 1.
 Leriche, R., and Morel, A. (1948). *Ann. Surg.*, **127**, 193.
 — and Stricker, P. (1933). *L'Arteriectomie dans les Artérites Oblitérantes*. Masson, Paris.
 Lindqvist, T. (1945a). *Acta med. scand.*, **121**, 32.
 — (1945b). *Ibid.*, **121**, 409.
 Nygaard, K. K., and Brown, G. E. (1937). *Arch. intern. Med.*, **59**, 82.
 Panton, J. A. (1923). *J. Anat.*, **57**, 106.
 Pearl, F. L. (1937). *Amer. J. med. Sci.*, **194**, 505.
 Platt, H. (1930). *Arch. Surg.*, **20**, 314.
 Ratcliffe, A. H., and Jepson, R. P. (1950). *J. Neurosurg.*, **7**, 97.
 Simmons, H. T. (1936). *Lancet*, **1**, 73.

A scholarship of £100 to £150 will be awarded by the National Association for the Prevention of Tuberculosis to a registered female nurse, working at the time of her application in Scotland, whose name is on the Queen's Roll of the Queen's Institute of District Nursing. Preference will be given to a nurse working in the Highlands. The scholarship will enable her to spend a period of from three to six months in post-graduate study in hospitals or clinics in Scotland, England, or Scandinavia. Candidates should state age, qualifications, and previous experience; reasons for wishing to do postgraduate work in tuberculosis; and should affirm their intention to continue in tuberculosis work after attaining the scholarship. Application should be made to Miss A. J. Weir, Scottish Secretary, N.A.P.T., 65, Castle Street, Edinburgh, 2, by November 1.

E. WILSON HALL, A. GEOFFREY SHERA, AND E. OWEN FOX: CARCINOMA SIMULATING HERNIA

FIG. 1.—Inguinal gland. Mitoses, vesiculation, and inflammatory cells; remnants of lymphoid tissue. ($\times 220$.)

FIG. 2.—Inguinal gland. Oat cells. The hyper-chromatic nuclei are almost universal. ($\times 290$.)

FIG. 3.—Increased density in right lower lobe; probable bronchial carcinoma. Adhesions are present at the left costophrenic angle.

G. SMITH: EFFECT OF DIATHERMY CURRENTS ON METAL IMPLANTS

FIG. 4.—Primary growth of right lung (lower lobe). ($\times 100$.)

FIG. 1.—Site of implant (tantalum gauze). ($\times 30$.)

FIG. 2.—Tissue reaction to implant (tantalum gauze). ($\times 200$.)

A. M. BOYD AND R. P. JEPSON: EXTERNAL ILIAC ARTERY THROMBOSIS

FIG. 1.—Case 2. Haematoxylin and eosin. ($\times 12$.)

FIG. 2.—Case 2. Haematoxylin and eosin. ($\times 60$.)

FIG. 3.—Case 2. Weigert's elastic stain. ($\times 60$.)