Chronic Iridocyclitis

There is a form of uveal inflammation which manifests itself in a manner likely to remain unnoticed by the patient for some time. The first symptom is diminution of vision, or maybe rapid increase in presbyopia. On examination only very fine vitreous opacities and fine K.P. are found. With a slit-lamp it is possible to see cells circulating in the aqueous. There is probably no circumcorneal injection (ciliary flush).

In spite of local treatment the disease tends to remain active; very often a secondary glaucoma supervenes, and this is a most serious matter. Very few of such cases resolve, and the eye is usually lost after a period of relapses. Little or nothing is known about this disease. It is attributed to such possibilities as septic absorption, allergic reaction, and tuberculosis—but rarely is anything definite found, elimination of which cures the local disease. Treatment therefore consists of atropine, rest, and complete overhaul; and if secondary glaucoma occurs operative treatment may become urgent.

EXPEDITION TO NIGERIA

M.R.C. SEARCH FOR TROPICAL PLANTS

BY

R. K. CALLOW, M.A., D.Phil.

Research Chemist on the Staff of the Medical Research Council

In November, 1949, an expedition was sent by the Medical Research Council to Nigeria to investigate certain tropical plants as possible sources of raw materials for the preparation of cortisone. The members of the expedition have recently returned, and, although evaluation of results must wait upon the completion of detailed laboratory investigation, some account of the background of their work and of its progress can be given.

The expedition, under the title of the Medical Research Council Medicinal Plants Survey, consisted of myself and a technical assistant, Mr. W. Read, from the National Institute for Medical Research, with a botanist, Mr. R. D. Meikle, of the Royal Botanic Gardens. During our stay in Nigeria we had our headquarters in University College, Ibadan, and grateful acknowledgment is made of the invitation from the Principal, Dr. Kenneth Mellanby,

and of the laboratory facilities which were made available by Professor R. J. McIlroy and Professor F. W. Sansome, in the Chemistry and Botany Departments respectively.

Origin of the Expedition

To explain the need for the expedition it is necessary to recall the sensational announcements, made just over a year ago, of the therapeutic activity of cortisone ("Compound E," 11-dehydro-17-hydroxycorticosterone; Formula I) in rheumatoid arthritis, and the fantastic discrepancy between the quantity of this compound which could be made available and the potential demand for its use. Although the compound is present in extracts of adrenal cortex, to obtain any considerable amount from this source is plainly impossible. A complicated many-stage synthesis from bile acids is known but is expensive, and the United States journalists have emphasized sufficiently the almost astronomical number of cattle which would be required for the maintenance of a single patient.

Synthetic routes to the adrenocortical hormones have been, and are being, busily explored; but a major difficulty is the introduction of the oxygen atom in position 11 in the steroid ring system (* in Formula I). This structure seems to be essential for the required type of biological activity. Were it possible to begin a partial synthesis

from a natural product with this bit of the molecule "ready made," the problem would be immensely easier. Attention was early directed to sarmentogenin, one of the group of heart poisons, the constitution of which, with an 11-hydroxyl group (Formula II), had recently been proved by Katz, in Switzerland.

Sarmentogenin was believed to occur in the seeds of *Strophanthus sarmentosus* P.D.C., a climbing plant of West Africa, and United States journalists rapidly sketched visions of vast areas of Texas planted with crops which would supply the rheumatoid arthritics of the world. The bubble burst when two of Professor Reichstein's colleagues collected in West Africa authentic seeds of *S. sarmentosus*, and it was shown that they contained little or no sarmentogenin but mainly a new heart poison, sarverogenin, the constitution of which is still undetermined. The origin of sarmentogenin now presented a fascinating scientific mystery.

Sarmentogenin had been isolated by three groups of workers, on each occasion from commercial samples of seed coming on to the market described as seeds of *Strophanthus hispidus*. In each instance pharmacognostic examination indicated that the seeds were not those of *S. hispidus*, but in general corresponded most closely in properties, particularly in colour reaction, with seeds of *S. sarmentosus*. If now, from the Swiss results, the source of sarmentogenin were not *S. sarmentosus*, what seed

FIG. 1.—Bush resthouse at Kagara. The M.R.C. expedition on tour.

could it be? It was a seed which occasionally came on the market in large quantity. Could it be the seed of a botanically unrecognized species, of a species hitherto considered rare, of a strain or variant of either *S. hispidus* or *S. sarmentosus*, or a seed used as an adulterant by native collectors? Precise information, even of the distribution

FIG. 2.—Near Zaria, N. Nigeria. An African forest guard with blossoms of *S. sarmentosus*.

African species of *Strophanthus* is being described—so far without sarmentogenin being found except as a minor constituent.

An Important Objective

A second, and potentially perhaps more important, object of the Medical Research Council expedition was to investigate plants botanically allied to *Strophanthus* which might contain useful steroids, not necessarily heart poisons. Plants of the Apocynaceae, the family which includes the genus *Strophanthus*, and the Asclepiadaceae were given special attention, and the Director of Research of the Colonial Products Research Council was able to suggest

FIG. 3.—*Strophanthus sarmentosus*: an unripe pod.

importance, but the subject is not yet sufficiently developed for anticipatory discussion.

The botanical work by Mr. Meikle was of prime importance in the unambiguous identification of the species collected. Moreover, his observations have added a great deal to our knowledge of the distribution and habitats within Nigeria of *Strophanthus* and other species. Botanically, the Nigerian flora is but partially known and recorded

as lists of species from scattered localities. Questions of distribution, habitat, and oecology—all that lies beyond herbarium botany—await investigation.

In its botanical exploration and collecting the expedition had the invaluable help of officers of the Agriculture and Forestry Departments with their local knowledge or guidance to sources of information. *Strophanthus* is neither a crop nor a forest product, and in this actively—even feverishly—developing country these officers are busy men. As one put it, "Our people are too busy to sit under bushes waiting for pods to ripen, but we shall be delighted if you come out to do that job, and we will give you all the help we can." The Emir of Katsina, awake to the potential importance of a new natural product, instructed district and village heads to organize collections of "kwankwani," and an almost overwhelming quantity of 10,500 pods of *S. sarmentosus* was obtained within four days. Elsewhere, the expedition visited villages of the primitive "pagan" tribes and gathered the crops from their bushes cultivated for the hunters' supply of arrow poison.

Potentialities for Research

These occupations were by no means sedentary: the botanist and the chemist covered over 9,000 miles by road in the course of their survey and, incidentally to their main task, had a great opportunity of appreciating the wealth of scientific and medical subjects calling for research in West Africa. The difficulties are to find housing, equipment, and money for work to be done by people in the country with knowledge of conditions there. Nigeria is a healthy region for Europeans now that specifics against malaria are available, but the high temperatures and humidities, particularly in the south, are trying, and exasperations and frustrations are magnified. Nevertheless scientific work is beginning—for instance, the recently organized scheme for investigation of Nigerian medicinal plants and native medicines at University College, Ibadan—and the African, with the heavy handicap of lack of any tradition of science, and sometimes even literacy, is demanding and obtaining European education, and efforts are being made to attract him to technical and scientific work for his country.

In Nigeria, the expedition's programme of laboratory work was subordinated to collecting, with the result that a large quantity of material, in the form of seeds and dried plants or crude extracts, which was brought back has now to be examined more closely. This will take time, and, though if all goes well the scientific results will emerge in due course, it must not be thought that a solution of the problem of getting a precursor of cortisone is just round the corner. Even if a plant rich in such a substance be found, its cultivation and the methods of extracting the raw material will need further long research.

A circular from the Ministry of Education discussing the education of girls before they start nursing courses takes the view that the best preparation that most students can have for nursing is a broad general education at school, preferably up to the age of 18. On the other hand, some secondary schools have introduced full-time pre-nursing courses which have proved of considerable value, and the Minister hopes that local authorities will stimulate interest in the courses and establish new ones where necessary. Part-time pre-nursing courses are sometimes held for girls who leave school at 15 and begin earning a living. The wastage from these courses is often high, but local authorities are asked to provide them where needed. The wastage may be reduced if the girls obtain advice from the youth employment officer.