It would seem that failure of forceps delivery caused by the cervix not being fully dilated arises mainly through the inability to recognize that the cervix must be completely effaced and retracted over the presenting part before forceps may be safely applied in domiciliary midwifery. Occasionally the general anaesthetic given preparatory to the application of forceps causes the head to retract, and examination then reveals a rim of cervix. By mistaking the tight rim of the primigravid cervix four-fifths dilated for the soft dilatable cervix of the multigravida, which is no bar to delivery, the forceps are applied but without delivery of the head.

It was formerly held that caesarean section, even by the lower-segment route, had little or no place in the treatment of failed forceps cases owing to the serious maternal risk from infection. It is now submitted that the results of this series discredit that view, and that the introduction of chemotherapy has revolutionized the treatment of failed forceps cases caused by brim disproportion when the baby is still alive.

In view of the high incidence of puerperal infection it is advisable to begin penicillin therapy immediately in all cases. In spite of chemotherapy the puerperium is often prolonged, with urinary complications much in evidence; and as these cases are potentially infected and therefore a risk to normal patients isolation is recommended.

The case needing forceps delivery when the head is not showing at the vulva is the one that may lead to difficulty. Practitioners who are not experienced obstetricians would be wise to confine themselves to delivering at home only those patients needing a low-forceps delivery.

Summary and Conclusions

The records of 100 cases of failed forceps have been investigated.

Primigravidae comprise almost three-quarters of the patients.

Occipito-posterior positions and deep transverse arrest of the head account for over 50% of the cases; the cervix was not fully dilated in 20%.

Since the introduction of chemotherapy the maternal mortality has been materially lowered, as has the foetal mortality, but to a less extent.

Internal version has little or no place in the treatment of failed forceps delivery.

Brim disproportion should be treated by lower-segment caesarean section; the results from this operation were uniformly good in spite of the previous manipulations.

My thanks are due to the honorary staff of the Birmingham Maternity Hospital for constructive criticism and permission to publish.

BIBLIOGRAPHY

- Baird, D. (1935). *Lancet*, **2**, 801.
 Bourne, A. W., and Williams, L. H. (1932). *Recent Advances in Obstetrics and Gynaecology*, p. 394. London.
 Browne, F. J. (1946). *Ante-natal and Post-natal Care*, p. 290. London.
 Claye, A. M. (1931). *Practitioner*, **126**, 611.
 De Lee, J. B., and Greenhill, J. P. (1947). *Principles and Practice of Obstetrics*, pp. 657, 907. Philadelphia.
 Dobbie, B. M. Willmott (1948). *Obstetrics and Gynaecology*, p. 220. London.
 Feeney, J. K. (1947). *Irish med. J.*, **6s.**, 190.
 Hendry, J. (1928). *British Medical Journal*, **2**, 185.
 Kerr, J. M. Munro (1937). *Operative Obstetrics*, p. 471. London.
 Lane-Roberts, C. S. (1935). *Practitioner*, **134**, 731.
 Miller, Douglas (1927). *British Medical Journal*, **2**, 685.
 — (1928). *Ibid.*, **2**, 183.
 Morris, W. I. C. (1947). *Edinb. med. J.*, **54**, 90.
 Phillips, L. (1926). *Lancet*, **1**, 113.
 Shaw, Sir William Fletcher (1928). *British Medical Journal*, **2**, 188.
 Stabler, F. (1947). *Ibid.*, **2**, 878.
 Stacey, J. E. (1931). *Ibid.*, **2**, 1073.
 Wrigley, A. J. (1946). *Proc. roy. Soc.*, **39**, 569.

USE OF RECTAL THIOPENTONE IN CHILDREN

BY

JOHN LORBER, M.B., B.Chir., M.R.C.P., D.A.

(From the Department of Child Health, University of Sheffield)

Many diagnostic or therapeutic procedures in paediatrics are painful or so unpleasant that the child will not co-operate. This is especially so in the first years of life, when even some painless procedures cannot be performed satisfactorily or without undue loss of time.

The commonest methods of sedation, such as the oral administration of chloral hydrate and of various barbiturates, are both unreliable and insufficient for many purposes. Thomson *et al.* (1949) found that with children over 1 year of age the inhalation of "trilene" gave satisfactory results during minor diagnostic procedures of short duration, such as lumbar puncture or aspiration of the chest. This method is not practical in infants or when access to the face or prolonged sedation is desired. Further, most children dislike the smell of anaesthetic agents and are afraid of a mask over their face.

In paediatric neurological practice complete examination of the fundus oculi and encephalography are procedures for which suitable sedatives are required. For the latter procedure the sedative action should last throughout the injection of the air and the subsequent radiological examination.

Rectal thiopentone has been used as the routine sedative in the streptomycin unit of the department of child health of the University of Sheffield since the beginning of 1949. This method was also used on a few occasions in other departments of the Children's Hospital, Sheffield, during the same period. It is felt worth while to publish the results to date, not because the method is original but because its value is not generally known.

The use of thiopentone by the rectal route was described by Weinstein (1939), Hooper (1942), and Weinstein and Light (1943). The last-named authors used the method in 1,500 cases, some 50 of them being children. They adopted a dosage scheme of 1 g. per 50 lb. (22.7 kg.) of body weight, and used thiopentone as a basal anaesthetic before inhalation anaesthesia or spinal analgesia. They were satisfied with its therapeutic action and noted no toxic effects. Burnap *et al.* (1948) reported similar findings in 300 children, and noted the absence of complications even after repeated administrations. Hunter (1943) used smaller doses (0.75 g. per 50 lb.), but his results in a small series were not so consistently satisfactory.

Material and Method

The results of 113 consecutive rectal administrations of thiopentone to 64 children between the ages of 3 months and 12 years are reported. Eleven were under 1 year of age. Fifty-six of the children were suffering from some form of tuberculosis, mostly meningitis and miliary tuberculosis; the other eight suffered from various neurological disorders. The examinations were usually carried out shortly after admission, when many of the children were extremely ill, and were often repeated on two to six occasions.

The indications for rectal thiopentone sedation were examination of the fundus oculi on 54 occasions, encephalography on 47, and other procedures (cystoscopy, laryngoscopy, etc.) on 12.

The following method was used. A small enema was given to constipated children a few hours before the proposed administration, to help the absorption of the drug and minimize the risk of its return after injection. No food was given for at least three hours before the thiopentone was due. A hypodermic injection of atropine, 1/150 or 1/100 gr. (0.43 or 0.65 mg.), was given half an hour before the thiopentone in almost all cases. This is not essential, but is useful in controlling salivation.

The thiopentone was dissolved in 20 to 40 ml. of warm tap-water and injected with a syringe slowly into the rectum by a rubber catheter. This injection took at least five minutes to complete. Afterwards the buttocks were held or strapped together to prevent some of the drug being returned. When this was not prevented, half the original dose was given half an hour later, if the child was not sufficiently sedated. The dose employed was 1 g. per 50 lb. of body weight. A detailed schedule is given in the accompanying Table.

Dosage Schedule of Thiopentone for Children Weighing up to 52½ lb.

Weight of Child		Dose of Thiopentone
lb.	kg.	
7.5-12.5	(3.40-5.67)	0.2 g.
12.6-17.5	(5.71-7.93)	0.3 g.
17.6-22.5	(7.98-10.20)	0.4 g.
22.6-27.5	(10.25-12.47)	0.5 g.
27.6-32.5	(12.51-14.74)	0.6 g.
32.6-37.5	(14.79-17.01)	0.7 g.
37.6-42.5	(17.06-19.28)	0.8 g.
42.6-47.5	(19.32-21.54)	0.9 g.
47.6-52.5	(21.59-23.81)	1.0 g.

The child was usually asleep by the time the injection was completed, and then was left undisturbed lying on its side for at least another five minutes to ensure adequate sedation.

When examination of the fundi was planned the pupils were previously dilated with 1% homatropine drops.

Results

With two exceptions the results were uniformly satisfactory. Systematic examination of the eyegrounds, with a detailed description of all abnormalities and their precise location, could be completed in five minutes. As the desired effect of the thiopentone lasted for nearly an hour in all cases, it was possible for several observers to complete the examination or to demonstrate the findings to a class of students without tiring the patient. There was just enough time to complete an encephalographic examination. Although the children were sufficiently awake to take meals after some three hours they usually slept most of the next 24 hours. This was of great benefit in diminishing headache following encephalography. There was complete amnesia for 24 hours even if the child reacted to a painful stimulus, such as lumbar puncture.

In a few cases cystoscopy or laryngoscopy was performed successfully without any additional sedation, but if any surgical incision was needed—as, for example, the opening of an abscess or the making of cranial burr-holes—local analgesia had to be used as well.

In the absence of painful stimuli complete muscular relaxation was obtained. This was useful for abdominal palpation, usually carried out in addition to the original purpose of the sedation. An enlarged spleen or liver was easily felt. This method could therefore be used in particularly difficult cases for the diagnosis of abdominal disorders, though I have had no personal experience of rectal thiopentone given specifically for this purpose.

In no case was any general or local, immediate or remote toxic effect observed. Nevertheless, it is advisable to have

oxygen and an injection of nikethamide or picrotoxin in readiness in case of excessive sedation.

One infant was in danger of dying after aspirating stomach contents, but made a complete recovery. Another child had severe spasms of coughing with temporary cyanosis. These two accidents were due to failure to observe the correct technique: one was fed just before thiopentone was given, and the other did not receive atropine.

Rectal thiopentone should not be given within three hours of a meal. It is contraindicated in the presence of respiratory obstruction, painful conditions of the rectum, or diarrhoea. Otherwise it is safe even for extremely ill and semiconscious children. It is an excellent method for ophthalmoscopic examination of babies.

Summary

Rectal thiopentone was given on 113 consecutive occasions to 64 children aged from 3 months to 12 years. The dose was 1 g. per 50 lb. of body weight. Complete and satisfactory sedation was obtained for examination of the fundus oculi, encephalography, and other procedures not requiring surgical incision.

No local or general toxic effects were observed.

Detailed technical instructions are given.

Acknowledgments are due to Professor R. S. Illingworth and Dr. T. Colver for access to their cases.

REFERENCES

- Burnap, R. W. Gain, E. A., and Watts, E. H. (1948). *Anesthesiology*, **9**, 524.
 Hooper, C. L. (1942). *Ibid.*, **3**, 458.
 Hunter, A. R. (1943). *Brit. J. Anaesth.*, **18**, 126.
 Thomson, J., Grewar, D. A. I., and Grounds, J. G. (1949). *British Medical Journal*, **2**, 1449.
 Weinstein, M. L. (1939). *Curr. Res. Anesth.*, **18**, 221.
 — and Light, G. A. (1943). *Ibid.*, **22**, 68.

CONVULSIONS IN CHILDHOOD AN EPIDEMIOLOGICAL NOTE ON THEIR ASSOCIATION WITH EPILEPSY, ECLAMPSIA, AND STILLBIRTH

BY

T. W. BUCHAN, M.B., D.P.H., F.R.F.P.S.

Nathan and Ransford (1949) have adduced evidence in favour of the claim that convulsions in infancy are the early manifestation of epilepsy, and Grantly Read (1950) has drawn attention to the possible relationship which they may have to certain types of convulsion in pregnancy and labour. Since the commonest cause of convulsions in pregnancy and labour is eclampsia, which is not infrequently associated clinically with stillbirth, it is of interest to note that certain epidemiological kinships exist between infantile convulsions and epilepsy, eclampsia, and stillbirth. These kinships may be considered under (1) secular trend, (2) seasonal variation, and (3) regional distribution, limiting the investigation to mortality data relating to England and Wales.

Secular Trend

Mortality from these diseases has been declining for many years, and the latest available figures show that unprecedentedly low rates now obtain. Mortality from convulsions in infants under 1 year has fallen from 9.68 per 1,000 births in 1911 to 0.35 in 1947, and deaths at all ages from epilepsy show a decline—interrupted, however, during the two world wars—from 72 per million in 1911 to 29 in 1947. Mortality from eclampsia has fallen sharply since