REFRESHER COURSE FOR GENERAL PRACTITIONERS**POLIOMYELITIS**

BY

W. HOWLETT KELLEHER, M.D., D.P.H.*Physician Superintendent, Western Hospital, London*

The current outbreak of this baffling disease should soon be reaching its peak, July, August, and September being almost invariably the months of greatest prevalence. In Britain the latter part of September may confidently be expected to show a decline in case incidence, although cases will continue to crop up sporadically during the winter months. Winter epidemics, however, have been encountered in some countries, although rarely. An instance of this was a recent severe outbreak in an Eskimo village during an exceptionally hard winter.

It is likely that the disease has been with us for thousands of years, though its ravages have not been apparent until comparatively recent times. Scandinavia had the unenviable distinction of providing the first real outbreaks, and the disease still remains a serious problem in Northern Europe. Somewhat later the United States suffered from serious outbreaks, notably that in New York in 1907, with more than 2,000 cases. But it was nine years before that city had its most serious outbreak, with over 9,000 cases and a mortality of more than 20%.

It is not within the scope of this article to discuss the theories held to account for the epidemic conflagration of poliomyelitis after it has been known to smoulder for many years. What we must accept as ominous is the expectation, borne out by the experience of other countries, that it will remain with us as a serious menace until more is known about its source and methods of control.

Causal Agent : Prophylaxis

A filterable virus, one of the smallest known, is the cause. There are several groups of poliomyelitis virus, the great majority of human infections being due to one group only, which can be transmitted to monkeys and contains several antigenic strains. The virus is inactivated by a temperature of 55° C. for 30 minutes, and by a 1 : 5,000 solution of potassium permanganate in normal saline. The virus in infected spinal cord can be inactivated by 0.3% formalin in 3½ days. Virus in suspensions of cord in acid solution is inactivated in 10 minutes in the presence of 0.05 p.p.m. of free chlorine. The usually adopted water-chlorination methods are believed to be ineffective in practice, especially in the presence of organic matter: if there is none, however, 0.2 p.p.m. of chlorine should be effective.

The virus has been isolated from sewage and sewage-contaminated water, and from flies, and may occur on unwashed fruit and vegetables. It can be recovered from the patient's oropharynx in the first week or so of the clinical disease and from the faeces in at least the first month or two. Healthy carriers are also known, while some people have been found to excrete the virus for some weeks before the onset of symptoms. These facts help in planning prophylaxis.

During the incubation period (see below) contacts should be kept under close observation and visits to infected households forbidden. During epidemic periods in localities where cases have occurred all unnecessary gatherings, especially of children, should be avoided. Exhausting physical exercise by children or young adults may facilitate

infection or lead to more extensive paralysis if undertaken when the disease is beginning.

Some authorities have expressed the view that during a poliomyelitis outbreak inoculations which are not urgently required should be postponed. Operations such as tonsillectomy, adenoidectomy, and dental extraction should also be held over. A damaged mucous membrane may facilitate the entrance of infection.

Age Incidence and Immunity

Although the earlier epidemics in Britain showed up to a 90% attack rate in infants and young children, this group now accounts for little more than one-third of the cases, the other two-thirds being in children of school age and in the adolescent and adult population. Infants under 1 year old enjoy a relative immunity or may suffer a mild attack only. An example of this is the experience on more than one occasion in the Western Hospital that young nursing trainees in infant nurseries have suffered from non-paralytic or from severe paralytic attacks while their protégés appeared to escape entirely.

The mechanism of immunity is as yet unexplained. Plenty of evidence exists of the presence of a neutralizing antibody in the blood of both convalescents and infected adults in communities frequently affected by epidemics, but its presence does not guarantee immunity from attack. Second attacks of poliomyelitis can occur, although the risk of a second attack may be somewhat less for a year or two. Some, if not all, of the recorded cases may have been instances of infections with strains differing from the original infection. It is said that immunity is strain-specific. Serum therapy and prophylaxis have proved a failure.

Incubation Period

This has been variously estimated as between a few days and four or five weeks, but is commonly believed to be between 7 and 14 days. The disease experimentally produced or following injury or operation generally shows itself after 7 to 10 days. Where it has followed local tissue damage from inoculation, it may be noted in from two to four weeks.

Clinical Manifestations

The majority of clinical attacks show two phases. In the first, a stage of general systemic disturbance, it will be impossible in the majority of instances, particularly in the absence of an outbreak, to do more than keep the possibility of poliomyelitis virus infection in mind.

First Phase

The initial constitutional symptoms may closely simulate a virus respiratory tract infection. Cough, however, is only infrequently met with, but pharyngitis is common. Temperature varies within wide limits and no positive conclusions may be drawn from it. More significant may be severe headache, particularly frontal, and vomiting, probably of nervous rather than toxic origin. General muscular

pains and lethargy are more noticeable than irritability at this stage. This phase of the disease is most variable in duration, and although it may occupy several days it is important to emphasize now that the briefest and slightest disturbance, a mild malaise and a temperature of 98.8° to 99.6° F. (37.1° to 37.6° C.), may be the only manifestation of poliomyelitis. That, of course, is the true abortive type. It is obvious that its identity, apart from laboratory assistance, can only be inferred—particularly if previous contact is known.

Second Phase

This may be separated from the initial phase by several days of comparative well-being, but in many cases it either merges into it or it may predominate from the start. Where the phases are clearly separated the temperature chart will characteristically show a "camel" (dromedary) curve. It is obvious both clinically and from histological studies that the second phase represents the inflammatory reaction due to viral attack on the central nervous system. It is a true meningo-encephalomyelitis, and the clinical signs vary little from those of any other infective or post-infective viral meningo-encephalitis. This true meningeal phase is almost invariably present in some degree before paralysis, but the paralysis may come on very rapidly, especially in infants, even within a few hours of onset of the first general symptoms. This is particularly so in those cases in which a monoplegia has been suspected to have been influenced by inoculation reaction.

Except in the rare encephalitic types, one of the most striking objective signs in the majority of patients, particularly in the presence of considerable pyrexia and other discomforts, is alertness merging into irritability, at least in young children (the adult is usually more apprehensive than irritable). Frontal headache is often very severe, and vomiting may recur. Generalized hyperaesthesia or pain and tenderness may be complained of, but, apart from that evoked through movement of muscles, painful spasm is not generally a common feature of cases in the pre-paralytic stage seen in the Western Hospital, however troublesome this may be at the onset or after the paralysis has become evident. Tendon and superficial reflexes may be temporarily exaggerated, and there may be fibrillary twitching of muscles.

It is obvious that where, as quite commonly happens, an inflammatory reaction in the mid-brain is marked, any degree of objective signs indicative of encephalitis may be encountered. Nystagmus is quite common and should always be looked for. Complaint is often made of insomnia, even in young persons, and when this is obviously not due to pain it should excite suspicion of early irritative encephalitic involvement.

Almost invariably present at this stage is some spasm of muscles with stiffness of posterior cervical and spinal groups and of the hamstrings, but it is a very variable sign. The elicitation and significance of this sign are amplified below. Retention of urine due to temporary bladder dysfunction is common, particularly where the lumbar cord is heavily damaged. Constipation may also be troublesome.

Methods of Examination and Assessment

For practical purposes and simplicity the following broad classification of cases may be adopted:

I. *Non-paralytic*: Abortive, meningeal, peripheral neuritic, cortical.

II. *Paralytic*: Bulbar, spinal, or combinations of both (bulbospinal).

The term "bulbar" is rather loosely used for any clinical type resulting from inflammation in any part of the brain-stem, although only applicable directly to the much more limited area of the medulla oblongata.

Abortive and Non-paralytic Poliomyelitis

"Abortive" is the name generally given to cases which have not shown signs of inflammation of the central nervous system, whereas non-paralytic cases have progressed to that stage but show no muscle weakness. It is obvious that the abortive case is normally diagnosed only by inference as a contact or with laboratory help. The only symptom may be a slight pyrexia of brief duration, and an abnormal cerebrospinal fluid (a protein level of more than 55 mg. per 100 ml.) alone aids the diagnosis. The real importance of diagnosis lies only in the potential infectivity of the abortive case, which is often a carrier. There is certainly no danger to life or health. Since abortive and non-paralytic infections are said to be 100 times more frequent than paralysed cases, the impracticability of hospital isolation of every abortive case is clear. A very large proportion of cases admitted to our special hospitals to-day on suspicion of poliomyelitis cannot be confirmed as even abortive attacks. Hospitalization is therefore not justified where definite neurological signs are absent. The practitioner's difficulties in face of the possibilities and of family apprehension can be understood, but both the patient and his family may best be served by allowing him to remain under daily supervision at home, and the burden on the hospitals will be considerably lightened, often by more than 50%.

Non-paralytic poliomyelitis includes all cases with the subjective and objective disturbances of the central nervous system known as meningeal irritation. This also is a large group. The hospitalization of such cases is fully justified and indeed important unless their removal creates physical or mental trauma with subsequent worsening of paralysis. Strict rest can be prescribed and a look-out maintained for the earliest signs of dangerous complications. If paralysis ensues it is often evident within a day or two of the onset of meningeal signs, but it may be delayed for many days. Although paralysis becomes maximal within 48 hours of its appearance, instances not infrequently occur where fresh lesions may continue to crop up for as long as seven days after the initial paralysis. For example, respiratory muscle involvement often takes some days to become apparent, a further reason for the hospitalization of the suspected case with meningeal signs.

Paralytic Poliomyelitis

As stated above, this may be spinal, bulbar (brain-stem), or combinations of both. As it still accounts for the largest number of paralytic cases, the spinal will be considered first.

It must be remembered that spinal paralysis includes not only cases of limb paralysis but trunk muscle lesions, including diaphragmatic and intercostal. Constipation and retention of urine probably come into a different category: function generally recovers in a few days.

The symptoms which usher in an attack differ in no respect from those enumerated above in relation to non-paralytic and abortive poliomyelitis. An amplification of the indications of muscle spasm is, however, called for.

(1) *Neck Rigidity*.—This is rarely so marked as in acute suppurative meningitis, but the differentiation from that seen in early tuberculous meningitis may be difficult. Only very rarely is true head retraction noted. Infants and young children

will often naturally resist any attempt at flexion, simulating a true rigidity. By attracting their attention so as to cause neck flexion or by raising them by the shoulders from the supine position (when they should normally flex the head) it is possible to remove the doubt.

(2) *Back Rigidity*.—This varies greatly. In some cases the condition may be so extremely painful as to make it impossible to get even slight flexion, and in others it may be so slight as only to be elicited by asking the sitting-up patient to kiss his knees, which are fully flexed, the thighs also being fully flexed and in the adducted position. Intermediate degrees are often seen, as where the patient can get himself into the sitting posture but because of rigidity or shortening cannot sustain it without using his extended arms placed behind him as in a tripod (Amoss's sign). Very often the difficulty in maintaining the sitting position is not due to stiffness of the spinal muscles but to hamstring muscles in spasm or shortened, and this can readily be demonstrated by sitting him on the edge of the bed with flexed knees, when he is found to be able to straighten his back fully. Incidentally this is an excellent position in which to make tests of the serratus and rhomboid muscles. Spasm of the neck and back muscles may, if prolonged and severe in a patient who is allowed to assume any comfortable position, result in considerable shortening with marked opisthotonus.

(3) *Kernig's Sign*.—This is frequently very marked in poliomyelitis, but it may be absent, especially in infants. Indeed, any or all of the muscular rigidities described above may be absent.

It should be emphasized that all these signs commonly occur in non-paralytic poliomyelitis as well as in all degrees of the paralytic disease. No information regarding the paralysis may be deduced from their intensity.

Paralysis or weakness when it occurs may be so slight as to require the finest tests for its detection, or it may be extreme. It is of lower motor neurone type with diminished or absent tendon reflexes. Even in the most extensively paralysed cases the asymmetry which is typical of poliomyelitis may be detected. Sensory loss should not be encountered, but there may be subjective disturbances—i.e., a feeling of numbness—and in later stages poor circulation and trophic changes may give rise to local sensory disturbances.

The commonest sites of paralysis are the lower limbs and the shoulder girdle, in that order. In the leg the quadriceps, the anterior tibial, and the peroneal groups are most commonly affected, in the arm the deltoid and triceps muscles. When a whole limb is involved the lesion is, except possibly in infants, easily recognized, and one will be concerned merely with making a differential diagnosis. It is, however, with the lesser degrees of dysfunction that detection is difficult, and this makes a meticulous examination necessary. A suggested scheme of examination has been set out in detail by the present author.*

Useful Clinical Points

The examination should be systematic, commencing with the head and neck and proceeding to the upper extremities and shoulder girdle, the trunk muscles and the lower extremities. Certain points must be noted:

- (1) With shoulder-girdle involvement look out for diaphragmatic and/or intercostal weakness.
- (2) With lower extremity weakness watch for retention.
- (3) Immobilization due to pain may give rise to an erroneous impression of the severity of the paralysis.
- (4) The abdominal recti are often asymmetrically attacked.
- (5) Even in very young children substitute movements to correct the defect of paralysis can very quickly be learned. A frequent example of this is rotation of the scapula to

produce abduction of the arm, and tilting the pelvis to help the dorsiflexion of the foot.

(6) In examination of infants accurate estimates of power may be made by the active resistance of the infant to handling. The more annoyed he is the better for one's purpose. Tempting with a toy or a sweet, or stimuli such as plantar tickling, will help greatly. Even assessment of minor weakness may quite accurately be made in this way.

(7) No single examination in a suspected case will suffice. Quite apart from the possibility of an extension of the paralysis or weakness, there are the factors of uncooperativeness and pain which may distort the picture. Continued raised temperature in the pre-paralytic phase or an unsettled one after the onset of the paralysis should make one alert to the possibility of extension. Following such stages a settled temperature for several days almost invariably denotes quiescence, with little further risk.

(8) Paresis when slight may often be detected only when the suspected muscle or group is compared with the opposing limb.

(9) Wasting may be hidden, in babies particularly, by fat. It is usually evident after about a fortnight.

(10) Absent or sluggish reflexes do not necessarily denote the presence of paralysis or even of paresis, but where an asymmetry of response is undoubted it is rarely misleading. The triceps and ankle jerks of normal infants are often difficult to elicit.

(11) Rapid deformity—e.g., foot-drop—may develop through spasm and shortening of opposing muscle groups; hence early diagnosis is of importance in prevention. Over-stretching of paretic muscles may seriously limit their recovery.

(12) Early recognition of respiratory muscle damage is necessary in order to avoid fatigue and further, perhaps irreversible, neurone damage through anoxia.

In any event an early consultation with a specialist colleague will help to avoid occasional pitfalls. Even those with extensive experience are not infrequently humiliated.

Prognosis in Spinal Paralysis

A flail limb in which no partial recovery has occurred in a few weeks is unlikely eventually to show even reasonable recovery. If there is no sign of return of power after four or five months the prognosis is almost negative. Where some recovery is evident early one may expect that with proper supervision the recovery will continue up to 18 months after onset. If neuronal recovery is to take place it does so within a few weeks.

The prognosis in recovery of movement is naturally greatly influenced by physical and orthopaedic treatment; full mobility of joints must be present before any power which remains can be utilized.

Bulbar (Brain-stem) Lesions

This is a most important group, to which full justice cannot be done in this necessarily condensed review. The 1947 admissions to the Western Hospital showed cranial nerve lesions in about one-third of all cases, which was considerably more than previously recorded. As in the spinal group, asymmetry and variability in symptomatology are the rule, ranging from the slightest of attacks to the most virulent. Rarely severe encephalitic symptoms are seen, but in the great majority of bulbar cases paralysis or weakness of certain cranial nerves is the only manifestation. Lesions of the upper group of cranial nerves (third to eighth) must be carefully looked for. For the detection of fifth-nerve involvement it may be necessary to make tests against resistance. Seventh-nerve lesions are relatively common and may occur alone. They are almost always of lower motor neurone type. Eighth-nerve (vestibular) disturbance is not uncommon, but frequently of brief duration.

*Kelleher, W. H., *Med. Pr.*, 1947, **218**, 452.

The lower group of cranial nerves is far more important because of the threat to life which extensive lesions involve. The ninth, if affected alone, and especially if unilateral, may cause little inconvenience. The more extensive combinations of the ninth, tenth, and twelfth cause great inconvenience and danger, due to obstruction of the airway by unswallowed buccal secretions. Another serious threat to life arises from complete laryngeal abductor paralysis, the approximation of the vocal cords causing great difficulty in inspiration. The difficulty is increased with spasm of the adductors. A common enough occurrence is eleventh-nerve paralysis, often detected only by muscle resistance tests.

By far the most fatal of all complications arises from inflammation of the respiratory centre in the medulla, giving rise to variations in depth, rate, and rhythm of breathing, or from defects of the vasomotor centre in the same area leading to circulatory failure.

No examination of a poliomyelitis patient is complete, therefore, until a careful test of the cranial nerves has been made. The pharynx should always be carefully inspected, as prompt treatment (e.g., postural drainage) may avert disaster.

Respiratory Failure

In addition to the commonly fatal type of respiratory failure arising from bulbar damage, it is most important to be able to make an early diagnosis of the more treatable muscle (spinal) paralysis or weakness. Careful and repeated inspection is necessary, particularly where shoulder-girdle paralysis is present. Serious anoxaemia may have occurred long before mechanical defect is obvious. When dyspnoea is present the patient is in urgent need of artificial respiration. Among the earlier signs are an increasing restlessness, anxiety, sweating, dilatation of pupils, and an increasing pulse rate. Later, accessory muscles may be brought into play. Dilatation of alae nasi may be noted, and speech tends to become staccato or jerky. Stethoscopic examination should reveal poor air entry to the lungs, especially at the bases. Defective epigastric excursion may be seen or such excursion may be entirely absent. Often only the upper intercostals may be found active, aided by accessory muscles. Attempts at coughing will be noted as extremely ineffectual.

Unfortunately, a not uncommon combination is that of pharyngeal or laryngeal dysfunction with a blocked airway and a respiratory (spinal) failure. The prognosis here is extremely serious. Central (medullary) respiratory failure is almost invariably fatal.

Differential Diagnosis

This bristles with difficulties as judged by hospital admissions of suspects. In the initial stages with general systemic disturbance many cases of tonsillitis, otitis media, rheumatism, and catarrhal respiratory infections arrive in hospital. There is also an increasing number of cases of pyrexia of unknown origin. Admittedly, many of these could be abortive poliomyelitis and justify diagnostic lumbar puncture, but in inter-epidemic periods or where there is no question of contact it can be justified only on academic grounds.

In the meningeal stage and in the absence of paralysis the examination of the cerebrospinal fluid is fully justified, particularly where tuberculous meningitis must be excluded, since early treatment of that condition is now imperative; this is true also of pyogenic meningeal infections and cerebral abscess. Subarachnoid haemorrhage can be a cause of confusion, in the author's experience.

When paralysis develops the toxic neuropathies must be excluded, rarely with difficulty. Cerebral vascular lesions even in children (thrombosis) are not rare. Trauma of nerve tissue, myelitis, and tumours have all caused confusion. Acute toxic polyneuritis (Guillain-Barré syndrome) is often difficult to differentiate from severe forms. Pseudo-paralysis may simulate poliomyelitis, and here mistakes are less excusable.

Pain fixation from any cause (rheumatism, arthritis, or injury) must be excluded. Spasm of muscles due to pelvic inflammation may immobilize the hip. Erythema nodosum and acute osteomyelitis occasionally cause confusion. Again, less excusable are fractures and dislocations.

Treatment

There is no specific for this disease. None of the newly developed antibiotics appear to have any activity against the virus. There is almost universal agreement that the claims for serum (convalescent or gamma-globulin fraction) cannot be supported, even if administered in the pre-paralytic stage. The value of any method is most difficult to assess owing to the very large number of abortive and non-paralytic cases, and also because spontaneous recovery may occur in paralytic cases.

Physical and Orthopaedic Treatment.—Strict rest is necessary for pre-paralytic and early paralytic cases, both for avoidance of discomfort and for the prevention of the spread of paralysis as a result of fatigue. Sedatives and analgesics have their uses in diminishing pain and irritability, but must be cautiously used in the presence of respiratory paralysis. Apart from dealing with mechanical defects (e.g., respiratory dysfunction) and the correction of physiological upset, treatment will be directed to the prevention of deformity in paralysed limbs and the encouragement by various physical methods of the action of muscles or parts.

In former days the idea of treatment by strict immobilization held the field. This is now regarded by many orthopaedists as harmful and by others as certainly no better than methods involving freedom of movement of limbs. The important thing is to transfer responsibility for the treatment of the paralysed condition very early to the physical-medicine department of a hospital where the advice of an orthopaedic surgeon can be obtained. The majority of the authorities believe that passive movement with or without hot moist packs is ultimately of the greatest benefit in preventing deformities, wasting, and trophic changes. Gradually introduced and carefully graded active exercises are commenced quite early. These measures are greatly assisted when carried out in a warm bath or with the aid of supportive slings.

The Kenny Treatment.—The principles of treatment advocated by Miss Elizabeth Kenny make a heavy demand on staff and time. A modified technique is in use in many orthopaedic departments in this country. Few, however, can afford to omit splinting entirely. The majority combine a modified splinting technique with passive and active movements assisted by hot packs in the earlier stages. If no splints are employed the part must be frequently examined and the position corrected, measures hardly obtainable in the comparatively sparsely staffed physiotherapy departments of to-day.

Respiratory Failure.—This can be treated only in hospital, spinal paralysis cases being placed in a respirator, while postural drainage and aspiration, perhaps combined with tracheotomy, may help the bulbar group.