REFRESHER COURSE FOR GENERAL PRACTITIONERS**THE FAILING HEART**

BY

K. SHIRLEY SMITH, M.D., F.R.C.P.*Physician, and Physician in Charge of Cardiac Department, Charing Cross Hospital*

While dyspnoea is constantly present in heart failure, the symptoms and signs as a whole vary according to the cause of the failure. Thus heart failure in mitral stenosis is very different clinically from that in hypertensive disease. The recognition of these divergences and of their importance dates from the improved methods of determining circulation time and cardiac output during the past 20 years; moreover, the better understanding of the circulatory derangements in the various syndromes of failure has led to a more discriminating therapy. Although digitalis still holds an honoured place as a specific cardiac remedy, it is by no means in every case the most important, or even an advisable, medicament. For these reasons it will be best here to consider first the recognition of heart failure in general and in its various forms; the basic principles of prevention and treatment will then be reviewed. Each important syndrome of failure in relation to its cause will be considered and its treatment and management described. The various forms of acute failure, such as occur in cardiac infarction, diphtheria, and pulmonary embolism, will not be dealt with.

Recognition of Early Heart Failure

Breathlessness on exertion is the most constant indication of diminishing cardiac reserve, although this symptom obviously has many other possible organic causes, notably diseases of the lungs. But it may also derive from disorders of function such as the effort syndrome. When difficulty is encountered in distinguishing an organic from a functional dyspnoea the following points in symptomatology will help: breathlessness due to disease of the heart will be proportionate to the amount of exertion and may be accompanied by cyanosis; it will be constant from day to day and the patient will often claim that he feels quite well when the symptom is not being produced. Shortness of breath due to functional causes (e.g., in effort syndrome and sighing breathing, or suspirious) varies greatly from day to day and even from hour to hour, bearing no precise relation to the degree of effort; and, apart from this symptom, the patient has other complaints, often numerous, such as dizziness, "nerves," and insomnia.

The failing heart usually implies a diminishing cardiac output, and thus a reduced blood supply to muscles and vital organs. Physical weakness and disinclination for activity result. Sleep may become disturbed from no obvious cause, and the patient may have found benefit from using extra pillows. Cough is not uncommon, particularly on lying down at night. This symptom and paroxysmal nocturnal dyspnoea will be referred to later in the section on left ventricular failure. Instrumental or other special methods of investigation find little application in the diagnosis of cardiac failure, but radiographs will reveal engorgement of the branches of the pulmonary artery when the left ventricle is failing and at a time when there may be no other manifestation except paroxysmal dyspnoea. Some enlargement of the heart will be found in almost every case in which dyspnoea is of cardiac origin.

While in adults the first signs of congestive failure are rales at the bases of the lungs and oedema at the feet, in children the most important indication is the respiratory rate. This has been pointed out by Taussig, who states that when the rate exceeds 30 a minute the liver begins to enlarge. She draws the inference that rales at the bases in children cannot be taken as being due to failure unless a rapid respiration rate and hepatic engorgement coexist.

Treatment of Early Heart Failure

When the cardiac reserve is waning but before congestion has developed, adjustments in the regime and mode of life are of greater importance than medicines. The first requirement is to lessen the work of the heart. The obvious means of doing this is by curtailing the patient's daily activities and increasing his hours of rest, both at night and in the daytime. A laborious occupation must be discontinued and lighter work sought. If insomnia impairs the night's rest a sedative at bedtime, such as phenobarbitone 1 gr. (65 mg.) or sodium amytal 2 gr. (130 mg.) should be given. At this stage digitalis therapy is not required. The work of the digestive system may be lessened by avoidance of heavy meals. A search for contributory causes of overburdening of the heart should be made, and, if found, remedied. Thus coughing should be checked by a codeine linctus. Obesity, which is a common and sometimes heavy burden for the damaged heart, should be corrected by adjustments of diet—chiefly the restriction of starchy foods and the avoidance of fat, fried foods, and sugary foods.

Infections, and particularly those of the respiratory tract, are often immediate causes of heart failure. As shown below, this is especially the case when chronic pulmonary disease overburdens the right heart; but it is also true in other forms of heart disease, and many a patient with advanced hypertension or mitral stenosis holds his own until an attack of bronchitis or influenza ushers in heart failure with increasing breathlessness, which may be paroxysmal. This serious consequence is attributable to toxæmia, the strain of coughing, and the heightened metabolism in feverish illnesses. For these reasons persons with heart disease should do their best to avoid contact with colds or influenza, and if such infections are contracted they should be treated with especial care: anything more than a slight cold will necessitate confinement to the house or perhaps to bed. The infection itself will need prompt treatment according to its nature, and, particularly, measures should be taken for the relief of cough.

Anaemia is a frequent additional burden on the heart, because in this condition there is a great increase in the output and therefore in the work of the organ. If anaemia exists, prompt steps should be taken to determine its nature by blood examination and the appropriate remedy—iron or liver extract—administered. The repair of anaemia may so greatly lessen the work of the heart that failure is averted.

Use of Digitalis

If, in spite of the adjustments of regime and the lightening of the work of the heart, congestive failure develops,

the use of digitalis becomes necessary. Digitalis is then of value irrespective of the cause of the failure, of the rhythm of the heart, the level of the blood pressure, and the presence or otherwise of mitral or aortic lesions. This drug, however, cannot be effectively administered unless the facts concerning excretion and action are known. The body can eliminate about 20 minims (1.2 ml.) of the tincture or 2 gr. (130 mg.) of the leaf preparation in 24 hours; only when the daily dosage exceeds these amounts does digitalis begin to accumulate in the body. When a certain concentration is reached in the body the dyspnoea of heart failure is relieved, diuresis sets in, and the discomforts due to oedema of the legs and engorgement of the liver disappear. Digitalization is the process of securing this beneficial result while avoiding toxic symptoms due to overdosage; these are headache, nausea, vomiting, and slowing of the heart with pairing of the beats.

The quantity of digitalis necessary to bring about digitalization in a patient of 10 st. (63.5 kg.) weight is about 20 ml. of the tincture or 30 gr. (2 g.) of digitalis folia. For lighter patients less will be needed in proportion, and for heavier patients more. The digitalizing total may be given in divided doses over a period of 24 hours (rapid digitalization); thus an initial dose of 90 minims (5.4 ml.) of the tincture could be followed by 60 minims (3.6 ml.) after six hours, and 30 minims (1.8 ml.) six hours later, and thereafter 20 minims (1.2 ml.) thrice daily, this amount being increased or diminished. Sometimes in very rapid fibrillation with a heart rate of 150 a minute or more, or when a speedy result is essential, "digoxin" may be used in doses of 1.5 mg. orally or 1 mg. intravenously. These doses will produce full effects in six and two hours respectively, but they must never be given if the patient has been taking digitalis during the previous 10 days.

In general, these very rapid methods of digitalization are not necessary, and satisfactory effects upon congestive failure are produced in several days by the use of an initial dose of 6 gr. (0.4 g.) of the leaf followed by 3 gr. (0.2 g.) at six-hourly intervals, until the symptoms of dyspnoea and congestion are alleviated or toxic symptoms are produced, when the administration should be stopped for 12 hours and continued subsequently at a dose of, say, 2 gr. twice or thrice daily. Thereafter a maintenance dose of digitalis must be determined; this may range from 2 gr. of the leaf thrice daily to 1 gr. daily.

Treatment of Left Ventricular Failure

This is a very common form of heart failure, because the chief causes—namely, hypertensive disease and aortic valve disease—occur so frequently. The patient becomes aware of breathlessness on exertion, perhaps slight, and then experiences a paroxysm of dyspnoea which usually wakens him from sleep. The distress may be short-lived, so that sleep is soon resumed, or it may become a protracted struggle to breathe, lasting for an hour or more. For this disorder morphine is a sovereign remedy and should be given subcutaneously in doses of $\frac{1}{4}$ to $\frac{1}{2}$ gr. (16–22 mg.), repeated in half an hour if necessary. The patient should be propped up well with a back rest or pillows in bed or in an arm-chair. The latter is especially helpful in severe cases, as the dependent posture of the legs slows venous return and helps to lower the pressure in the right auricle; in the overloaded heart this results in an augmented cardiac output, so that pulmonary congestion is relieved. If, in spite of the use of morphine, dyspnoea persists or pulmonary oedema or Cheyne-Stokes breathing appears, an intravenous injection of theophylline ethylenediamine ("cardophyllin") should be given—0.48 g.

in 10 ml. of distilled water, administered slowly. Some patients with failure of this kind who have high blood pressure and distended veins with normal rhythm respond more dramatically to venesection than to any other procedure. This may be carried out either by slitting the antecubital vein direct through the skin with a sharp scalpel or by introducing a wide-bore Record needle or French's needle into the vein. In either case, after 15 to 20 oz. (420–570 ml.) of blood has been withdrawn, a pad and firm bandage are applied and the arm allowed to lie in a horizontal position.

In some patients with left ventricular failure there is, in addition to the usual features, conspicuous bronchospasm aggravating the respiratory distress. The safest means of treating this will be by the oral or, if necessary, intravenous administration of one or other of the theophylline derivatives, such as cardophyllin or "mono-theamin." Ephedrine and adrenaline are, in general, contraindicated in heart failure owing to their liability to increase the work of the heart; however, if the measures mentioned above fail to relieve bronchial spasm a subcutaneous injection of adrenaline hydrochloride, not exceeding 4 minims (0.25 ml.) in a 1 in 1,000 solution, might be tried.

When pulmonary oedema develops and is not quickly relieved by the methods described above, oxygen administration should be started without delay. The method of choice is the transparent oxygen tent which forms a canopy over the head of the bed. There must be no open fire in the room if this method is used, owing to risk of explosion.

Episodes of paroxysmal dyspnoea are due to failure of the left ventricle to discharge fully its contents, so that blood begins to accumulate in the left auricle and the pulmonary circulation. The attack should be a signal for a thorough review of the patient's mode of life, the state of the heart, and the general condition. Rest in bed is ordered for a week or two, the patient being placed on a light diet with no added salt; the intake of fluids during the latter part of the day is avoided so far as is possible. If oedema exists three or four injections of 2 ml. mercurial diuretic are given intramuscularly or intravenously at four-day intervals. Even if there is no oedema one or two such injections may help to relieve paroxysmal dyspnoea. Apart from the longer-term measures, one of the best preventives for nocturnal dyspnoea is a draught containing chloral hydrate 10 gr. (0.65 g.), with nepenthe 20 minims (1.2 ml.), given at night. In chronic left ventricular failure, digitalization, preferably by one of the slower methods described above, is often of value, and may bring good results when other therapy has failed. Large doses in hypertensive disease may induce auricular fibrillation.

Heart Failure in Chronic Rheumatic Heart Disease

Heart failure in mitral stenosis is compatible with a longer tenure of life than failure in any other chronic heart disease. I have records of a patient whom I have seen in failure at intervals during eight years, and it is not uncommon for failure to recur at intervals for four or five years. Failure in these patients is ushered in often by bronchitis or other respiratory tract infection, or sometimes by physical strain; fibrillation may make its appearance at this time or it may have existed previously. It is particularly in failure associated with fibrillation of rheumatic origin that digitalis effects its most brilliant results, and the rapid digitalization method, with massive or intravenous dosage, finds its chief application when such failure has developed suddenly with a very high heart rate.

Digitalization will be employed in conjunction with mercurial diuretics. Venesection is not advisable in these

patients. On account of the vascular engorgement and the oedema of the lungs, oxygen administration will often be of value in the initial stages of treatment of severe failure. Pulmonary infarction (generally due to embolism) is perhaps commoner in this variety of failure than in any other; pleural pain, sudden dyspnoea, or jaundice otherwise unexplained should arouse suspicion of this complication, which should be treated with morphine injection and oxygen therapy.

The Failing Heart in Chronic Pulmonary Disease

This form of failure, usually following long-standing bronchitis and emphysema, is one in which the cardiac output is substantially more than normal. Recognition of this fact is fundamental in treatment, because the raised output is a compensation for the reduced oxygenating power of the lungs. Any treatment, such as venesection, that depletes blood volume will be detrimental. For the reason that it may lessen cardiac output by lowering venous pressure, full digitalization may be prejudicial to these patients.

Failure in patients with chronic pulmonary disease most commonly develops in the winter months, and the first cold or foggy spell of the autumn reduces many of these patients to a serious condition in a few days. Such breakdown is due principally to recrudescence of infection in the respiratory tract. The lung symptoms become worse, with severe dyspnoea, wheezing, and cough; cyanosis deepens, while the cervical veins and liver become engorged; headache and mental confusion develop. This is one of the few forms of heart failure in which pyrexia may coexist, but fever is not always present. Normal rhythm may be maintained or fibrillation may supervene.

Treatment is based upon the existence of cyanosis and infection: oxygen therapy should be instituted at an early stage and maintained for days or a week or two, and at the same time penicillin treatment (200,000 units I.M. every six hours, or "prolophen" 1 ml. daily) should be started, whether or not obvious infection or fever exists. Sulphadimidine may be employed in addition. The necessity for immediate effective treatment in these cases makes it impracticable to await sputum examinations and tests of penicillin sensitivity of suspected causal organisms. The measures described will often retrieve situations that are well judged to be desperate.

Digitalis may be employed at a later stage, but only in moderate dosage, such as tab. digitalis folia 2 gr. twice or thrice daily. Morphine, if it is used, should be given in small doses and in conjunction with atropine. A few injections of mercurial diuretic should be given if oedema is more than slight. While salt should be restricted, too great a reduction of fluid intake should not be made, three pints (1.7 l.) in the 24 hours being a suitable quantity.

Some Less Common Forms of Failure

In thyrotoxic heart disease, heart failure develops only after many years, and generally only when no adequate treatment has been given for the thyroid disorder. Indeed, by the time heart failure sets in evidence of active thyrotoxicosis may be lacking; the ruins of the cardiac structure and its chaotic function are the relics of a fire that is smouldering or extinct. Such patients have hearts that are enlarged and fibrillating rapidly; they often have hypertension; the evidences of failure point to both ventricles, so that there is often pulmonary oedema and hydrothorax, while the liver and the cervical veins are engorged. The usual regime of bed-rest and fluid and salt restriction

is necessary, coupled with digitalization, which should be by fairly rapid oral methods. If swift control is gained by these methods it is an indication that little or no thyroid influence remains. Sometimes digitalis alone fails to control the heart rate, and success following the simultaneous administration of Lugol's iodine points to persisting thyrotoxicosis. In this case the treatment of failure must be followed by the treatment of the thyrotoxicosis either by operation or by methyl or propyl thiouracil.

Failure develops often in later middle life among patients who have severe thoracic deformity. In these cases cervical venous engorgement may be masked and hepatic engorgement may be difficult to assess. Increasing dyspnoea and oedema of the legs, with signs of pulmonary oedema, are the important indications. Treatment will be based on the general lines already described in conjunction with digitalization that is not too rapid. The evidences of right ventricular failure ultimately predominate and call for oxygen therapy, and the antibiotics if respiratory tract infection is suspected.

Treatment of Advanced Chronic Failure with Anasarca

This condition may be the final stage of failure due to any cause. The duration of life must in any case be short, and the aim of treatment is to secure comfort by dispersing the accumulation of fluid. In the first place salt intake must be reduced to less than 0.5 g. daily, milk being restricted and salt-free bread and margarine being used. Fluids are cut down to 25-30 oz. (0.7-0.8 l.) in the 24 hours. Any accumulation of fluid in the pleural cavities must be removed. If the legs have become greatly swollen with fluid and mercurial diuretics are diminishing in effect, then drainage of the legs by Southey's tubes is called for. It is often found that removal of fluid collections by these various means allows some restoration of cardiac function. Thus kidney function is improved and the mercurial diuretics regain their efficacy. Slowly the longer-term measures of salt restriction take effect and the patient may ultimately be quite rid of the oedema. He thus may become able to leave his bed by day and sleep undisturbed at night. Measures of this kind may be effective also in the even more intractable cases where there is recurrent ascites due to chronic congestive changes in the liver. Fluid accumulation may by these means be confined to the abdomen and the intervals between paracenteses be considerably lengthened. Mercurial diuretics function well when given intraperitoneally, and 1.5 ml. in 10 ml. normal saline introduced at the conclusion of tapping will sometimes produce a better result than intramuscular injection in disorders of this kind.

Dr. I. C. Fang, of the World Health Organization, has recently gone to Bangkok, where he will temporarily direct W.H.O. health activities in the Western Pacific. A regional organization, grouping all countries in that area, is to be established early next year. It is expected that the regional committee of W.H.O. member States in the Western Pacific will hold its first meeting in February or March to recommend the permanent location of the regional office, designate a director, and outline health campaigns needed in the various countries. Several international health programmes are now under way in the Western Pacific region: they include W.H.O. nursing, sanitation, hospital, and tuberculosis services in China, and joint W.H.O./Unicef programmes for mother and child health in Sarawak, Brunei, North Borneo, Hong Kong, and South Korea, as well as B.C.G. vaccination campaigns against tuberculosis in Singapore, Malaya, Formosa, and the Philippines. It is expected that the temporary W.H.O. office will be in Hong Kong until a permanent site is approved by the United Nations.