REFRESHER COURSE FOR GENERAL PRACTITIONERS**CARCINOMA OF THE STOMACH**

BY

C. F. W. ILLINGWORTH, C.B.E., M.D., Ch.M., F.R.C.Sed.*Regius Professor of Surgery, University of Glasgow*

Carcinoma of the stomach is one of the commonest of neoplasms, and one of the most malignant. It arises insidiously and in its early stages gives rise to no striking or characteristic symptoms; it spreads rapidly and soon extends beyond the limits of operative extirpation. Often the disease is already far advanced by the time the patient consults his doctor, and in some cases, though operation is performed without delay, the prospect of a permanent cure is small.

Gloomy though the picture appears, it is not without relieving features. In recent years advances in medical care and in surgical technique have enlarged the scope of operation and at the same time reduced its dangers. Improvements in pre-operative preparation, particularly the more ready recourse to intravenous administration of fluids, protein, and blood, now enable all but the most feeble patients to be brought into a condition suitable for operation, while technical advances, particularly the extended use of the thoracic approach, have made it possible to remove growths which formerly were regarded as inoperable. It is now practicable to remove growths originating in any part of the stomach, including the fundus and cardia, while the operation of total gastrectomy has lost much of its former danger.

As in all other forms of malignant disease early diagnosis is all-important, and it is with this aspect of the problem that this paper is mainly concerned.

Predisposing Factors

In those forms of malignant disease which develop on the basis of recognized predisposing factors the most promising line of attack is in the precancerous stage. It has long been recognized that chronic gastric ulcer may predispose to carcinoma, and, though the proportion of cases in which such a malignant change takes place is now regarded as quite small, this group is nevertheless of particular importance, for here we have an opportunity to establish the diagnosis and to carry out radical treatment at a really early stage. Indeed, the improved end-results that have been claimed by some workers in recent years have been due in large part to the increasing number of cases coming to operation at a stage when the chronic ulcer has only just begun to develop malignant characteristics.

There is some evidence that achlorhydric gastritis may similarly predispose to carcinoma. This term ("gastritis" is a misnomer) implies atrophy of the gastric mucosa accompanied by complete histamine-fast achlorhydria. The condition is well known as a concomitant of pernicious anaemia, but it may occur alone. Sometimes it is a congenital abnormality but more often is acquired; it is a common finding in the later decades of life.

The proportion of patients with complete achylia who ultimately develop carcinoma is probably not large; nevertheless it is large enough to emphasize the need for particular attention in such cases in order to ensure that the development of malignancy will not be missed.

Early Symptoms

It should be emphasized that the predisposing factors just mentioned are found only in a small proportion of cases of carcinoma of the stomach, and from the point of view of early diagnosis in the generality of cases there is a danger that their importance may be overstressed. It is far more important to remember that in something like 95% of cases the patient has been in good health and entirely free from digestive disturbance prior to the development of malignant disease.

The initial symptoms of gastric carcinoma are by no means constant or characteristic, and often they are at first of quite insignificant character. In general they fall into two groups: (1) impairment of appetite with some loss of energy and a vague feeling of deterioration of health; and (2) minor digestive disturbances with vague epigastric discomfort, fluctuant distension, nausea, and perhaps vomiting. In some cases a sudden distaste for an article of food, particularly butcher's meat, is the first feature to attract the patient's attention. While such symptoms are of course far from characteristic, when they develop in a middle-aged or elderly patient the possibility of carcinoma should always be considered. The onset of vague digestive disturbances in a middle-aged man with a clear gastric history should always arouse suspicion.

Pain is not a characteristic feature. There may be no pain at all or only a mild discomfort. When present it may be felt diffusely in the epigastrium or left hypochondrium, or in the precordial region, or in the back. Only rarely is it severe at this early stage.

The pain tends to be most pronounced an hour or so after meals, but there are many exceptions to this. Unlike the pain of simple gastric ulcer, it is usually but little relieved by the taking of food. It may, however, be relieved by an alkaline powder, probably because the gas evolved leads to belching and thus brings about some reduction in tension within the stomach.

Apart from this usual clinical picture the disease sometimes shows itself by symptoms related to the precise site of the lesion. For example, a growth situated close to the cardia may give rise to symptoms like those of carcinoma of the oesophagus, with dysphagia and regurgitation immediately after swallowing. In the same way a growth situated close to the pylorus gives rise to symptoms through obstruction of that orifice, and the whole clinical picture will be that of progressive pyloric obstruction, with retention vomiting and rapid loss of weight through dehydration.

Bleeding from the growth is a common feature. There may be frank haematemesis, though this is infrequent. Much more often there is an insidious loss of blood, leading to secondary (hypochromic) anaemia, which may be severe. In such cases the main burden of the symptomatology is related directly to the anaemia, with weakness, dizziness, and shortness of breath as the chief features. In a middle-aged man gastric carcinoma is the commonest cause of an obscure anaemia.

Lastly, it is important to note that some disturbance of the action of the bowels may provide the first clinical evidence of the disease. Sometimes there is constipation, probably attributable to the reduced intake of food; or there may be post-prandial diarrhoea of gastrogenous type, resulting from the achlorhydria or perhaps from infection of the ulcerated growth. In a middle-aged person of regular bowel habit the development of constipation or diarrhoea should arouse suspicion.

Clinical Examination

It is unfortunate that in the past medical teaching has concentrated so much on the late manifestations of gastric carcinoma, such as the characters of the palpable tumour, the large liver studded with umbilicated nodules, the secondary ascites, and the presence of metastases at the umbilicus, on the pelvic floor, or at the root of the neck. Important as these features are in relation to prognosis and treatment, they do not, of course, have any bearing on the early diagnosis, while emphasis on these gross manifestations tends to obscure the fact that in the early operable stage of the disease none of them should be in evidence.

In most cases the carcinoma originates on, or close to, the lesser curvature. When a person of average build is examined in the recumbent position a growth in this situation lies deep under cover of the liver and costal margin and can be reached only with difficulty, if at all, by the palpating finger. Abdominal examination may therefore be entirely negative.

Differential Diagnosis

Peptic Ulcer

The problem of distinguishing carcinoma of the stomach from simple peptic ulcer arises in two quite distinct types of case. First, there is the patient who gives a long history of indigestion and who is suspected, or perhaps known, to have had a simple ulcer of the stomach for a considerable time. In such a patient the possibility of malignant change in the ulcer may arise—the so-called ulcer-cancer. This problem is discussed below.

Secondly, there is the patient with no previous history of indigestion, in whom the question arises: Are the recent symptoms indicative of carcinoma or are they due to the recent development of a simple peptic ulcer (gastric or duodenal)? In this latter type of case the age incidence of the two diseases is important. Simple peptic ulcer arises most often before the age of 40, and it is comparatively infrequent for simple ulcer to occur for the first time in a patient over 50; whereas in carcinoma, as is well known, the incidence increases with advancing years. Consequently, in a middle-aged person whose symptoms are of recent origin the weight of probability is on the side of malignant disease.

In simple ulcer the pain tends to be more severe than in carcinoma, its relation to the taking of food is more precise, and as a rule the relief obtained by food or alkalis is more pronounced. There are cases, however, in which the symptoms of ulcer closely mimic those of carcinoma and in which such features as loss of appetite and loss of weight add to the confusion.

In such cases much information is obtained by a thorough investigation. A duodenal ulcer can generally be excluded without much difficulty by x-ray examination and by means of gastric analysis, for a high acid curve is almost invariable in duodenal ulcer and quite exceptional in carcinoma. A gastric ulcer is more difficult, for the x-ray examina-

tion may be equivocal, while hypochlorhydria is the rule. Gastrosopy is helpful; indeed, it is in just such cases that this method of examination provides its greatest field of usefulness.

If all these methods of investigation still leave the final diagnosis in some doubt, the therapeutic test may sometimes be employed. For this purpose the patient is placed on an intensive course of medical treatment, preferably by the continuous intragastric milk drip, for three weeks, after which he is submitted to further investigation by x-ray examination and gastrosopy. If at the end of this time there is clear evidence of healing, the medical regimen may be continued, otherwise operation should be advised without further delay. It should be emphasized that in such doubtful cases the therapeutic test must be interpreted in the light of strictly controlled x-ray and gastrosopic investigations; it must not be judged on the basis of clinical features, for the symptoms of carcinoma no less than of simple ulcer may improve greatly as a result of rest in bed with adequate medication.

Ulcer-Cancer

The development of malignant change in a benign chronic ulcer of the stomach is now regarded as infrequent, but this small group of cases is of great importance from the point of view of early diagnosis. Here we have a lesion of some chronicity—often it has been present for years—with a well-established liability to malignancy. It should offer a highly favourable field for the early treatment or even the prevention of the malignant complication. Yet, lamentable as it may seem, until recent years the prognosis of ulcer-cancer has been even graver than that of a newly developed cancer of the stomach.

The reasons are not far to seek. The patient with a chronic gastric ulcer has become inured by long experience to the discomforts of indigestion and does not notice the gradual worsening of his condition. The medical attendant, accustomed to the doleful tale and sorrowful aspect, may fail to perceive the early signs of malignant change. These early signs, moreover, are by no means characteristic. The malignant transformation is heralded by no new symptoms but simply an aggravation of the existing picture. Indeed, sometimes the pain, which has been the main feature of the ulcer during its period of benignity, now actually diminishes, though more often it continues much as before, but with the notable feature that it tends to become more continuous and less constantly relieved by food or alkalis, while such features as anorexia and loss of weight gradually make themselves apparent.

Attempts have been made in the past to establish definite criteria by which the development of malignancy in a simple gastric ulcer may be recognized at an early stage, but so far no precise test has been evolved. The size of the ulcer is no longer regarded as dependable evidence, for it is recognized that many small ulcers become malignant, while some large ones, even an inch or more in diameter, remain benign. It is true that x-ray examination and gastrosopy will generally furnish evidence when the malignant change is well established, but only too often by this time the growth has already spread widely in the stomach wall and in the lymphatic channels.

The conclusion to which one is led by these observations is inescapable. The proper treatment of ulcer-cancer is by prophylactic surgery. In other words, it is necessary to regard all cases of chronic gastric ulcer with a critical eye, and to recommend partial gastrectomy in those in which

malignant change may be regarded as a possible development. This is particularly so if the ulcer is situated within 2 in. (5 cm.) of the pylorus or if it does not heal readily and completely (as judged by x-ray examination and gastroscopy) after an adequate course of medical treatment.

Achlorhydric Dyspepsia ("Chronic Gastritis")

Apart from its possible significance as a factor predisposing to gastric carcinoma, achlorhydric gastritis is of some importance in the differential diagnosis. Generally achlorhydria is without clinical effects, but when symptoms do occur they usually consist of irregular dyspepsia with anorexia, loss of energy, and perhaps some loss of weight. In most cases their long duration is enough to rule out carcinoma as the original cause of the symptoms, but it is by no means easy on clinical grounds to exclude the possibility of malignant change of recent origin. X-ray examination, however, will be negative, while the true diagnosis can readily be made on gastroscopy.

Other Digestive Disturbances

Symptoms suggesting gastric carcinoma arise occasionally from many other causes, such as alcoholic gastritis, chronic gall-bladder disease, or even simple constipation. They may also be mimicked by flatulent distension complicating heart failure or in uraemia or other chronic toxic processes. In all such cases x-ray investigation may be required to exclude carcinoma.

Anaemia

It has already been shown that the whole clinical picture of gastric carcinoma may hinge upon anaemia due to insidious loss of blood. In such cases it is fatally easy to attribute the symptoms to the anaemia, and to be content with prescribing iron or even liver extract. Examination of the stools for occult blood will, of course, point immediately to the source of the blood loss and indicate the lines of investigation to be followed, but hospital experience shows that this simple test is sometimes omitted.

The converse error of mistaking a primary anaemia for carcinoma of the stomach is equally possible but much less serious, for in such a case a full investigation will be called for, in the course of which the proper diagnosis will be reached.

Methods of Investigation

In coming to an exact diagnosis of gastric carcinoma we rely more and more on laboratory methods of investigation. To interpret them properly the practitioner should appreciate not only the scope of such investigations but also—and even more important—their limitations.

The test for *faecal occult blood* has already been mentioned as being of special value in directing attention to the source of blood loss in a patient whose presenting feature is anaemia. Its usefulness is not limited to such cases, however, and it should be performed as a routine whenever carcinoma of the stomach is suspected. A positive result indicates the presence of an ulcerative lesion in the alimentary tract, and to this extent supports the diagnosis. But it must be emphasized that a negative result carries no significance whatever, even when repeated on several occasions, for quite often a carcinoma, especially of the scirrhus variety, bleeds not at all or only at long intervals.

Gastric analysis after a test meal is established as a valuable method of investigation in obscure dyspepsias, but in the diagnosis of carcinoma it has but a limited usefulness. It is true that in most cases of carcinoma there is complete

or almost complete absence of free hydrochloric acid. But complete achlorhydria may occur in health, or in association with other diseases, notably pernicious anaemia, while, conversely, a normal or even a high acid curve does not exclude carcinoma. Nor are such other features as a high content of combined acid, the presence of sarcinae, Boas-Oppler bacilli, and other organisms of positive value in the diagnosis. It is true that prolonged microscopical examination of the sediment may reveal the presence of malignant cells, and this finding admits of little doubt; but again a negative result has no significance. Gastric analysis is better omitted altogether, because of the time and trouble involved, the limited information gained, and, most important, the risk of interpreting falsely a negative result.

X-ray examination after a barium meal is, of course, the standard method of investigation, and in the hands of a competent radiologist a high degree of accuracy can be expected. Indeed, with a growth at the usual site in the lesser curvature, few methods of examination are more reliable. Difficulty arises occasionally, however, with a growth at an unusual site. Thus, a small lesion at the fundus or on the greater curve may be missed unless particular care is taken in the technical performance of the examination, while a small growth at the pylorus, giving rise to pyloric obstruction, is notoriously difficult to distinguish from a simple ulcer.

Gastroscopy has been introduced in recent years, and now holds a definite place in diagnosis, but the examination should be carried out by a specialist in this particular technique. Gastroscopy cannot be performed in all cases. It is obviously contraindicated in the presence of aortic aneurysm or cardiac enlargement, or when the thoracic spine is deformed or rigid through disease; sometimes the instrument will not readily enter the stomach, and in such circumstances to force it is to court disaster; if the growth is invading the lower end of the oesophagus the examination entails a certain risk; and finally in some cases, though the stomach is entered satisfactorily, a clear view of the pylorus cannot be achieved. Subject to these limitations, however, a great deal of information can be obtained from this method of investigation. Thus there is usually but little difficulty in distinguishing between carcinoma and a benign ulcer of the lesser curvature, while achlorhydric gastritis can be recognized at once by the thin, pale, atrophic character of the mucous membrane. Lastly, gastroscopy is sometimes the only method of examination by which, given a good view of the whole stomach, the absence of carcinoma can confidently be asserted.

Treatment

Apart from such obvious features as hepatomegaly, ascites, enlarged cervical glands, or pelvic metastases, there is no precise clinical evidence by which to determine if the tumour is capable of being removed. Formerly the site of the growth was a determining factor, and only those limited to the pyloric end of the stomach were deemed suitable for operation; but this no longer obtains, for by means of the thoracic or a combination of the abdominal and thoracic approaches the fundus and cardia are equally accessible. The size of the tumour is not a reliable criterion, for the main factor determining operability is the extent of lymphatic spread, and in this respect the large fungating tumour is sometimes less aggressive than the small scirrhus growth.

X-ray examination may provide valuable evidence. In some cases the fixity of the growth, as judged on screen

examination, may preclude operation; in others the lesser curvature may be visibly distorted by enlarged glands in the lesser omentum.

Such precise evidence is not often available, however, and in most cases the decision to operate is based more on hope than on expectation that resection will be possible.

Palliative Treatment.—As a rule the most distressing features in inoperable cases are retching, nausea, and vomiting, due in part to obstruction of the gastric pathway and in part to infection of the fungating growth. Much relief may be obtained in such cases by gastric lavage. For this purpose a stomach tube should be chosen of diameter midway between a Ryle's tube and one of full size. A No. 16 size is most satisfactory, for it can be introduced with little discomfort and yet is large enough to permit evacuation of blood clot and debris. A dilute solution of sodium bicarbonate may be used for the lavage, which may be performed daily, or even twice daily, with benefit.

PROGRESS IN CANCER RESEARCH ANNUAL REPORT OF B.E.C.C.

The 27th annual report of the British Empire Cancer Campaign covering the year 1949 was presented on July 12 at the annual meeting of the Campaign, over which the Duke of Gloucester presided. The editor of the report, Sir Heneage Ogilvie, summarizes the year's work as "steady progress," research having been conducted at an increasing pace.

Evidence has been accumulated at the Chester Beatty Research Institute to substantiate the carcinogenic action of the aromatic nitrogen mustards. Work at the Animal Supply and Research Units of the B.E.C.C. has shown that mammary-tumour tissue of the mouse has a higher content of the enzyme β -glucuronidase than healthy breast tissue.

It has been confirmed at Leeds University Department of Experimental Pathology and Cancer Research that tissue from mouse sarcomata can be frozen and dried and yet retain the capacity to reproduce the same type of tumour in susceptible animals. But it was considered that this may not constitute evidence of the existence of a virus as the infective agent, for sarcoma cells may survive freezing and desiccation in the presence of 5.3% glucose. If this is the case, the resulting tumours may be directly derived from the surviving cells. Work also carried out at Leeds does not suggest that mothers of breast-cancer patients are more liable to cancer in general, or that they are more liable to cancer of the breast than the average woman in England and Wales.

Chemotherapy

From the Westminster Hospital considerable improvement, in some cases lasting two years, is reported in certain cases of carcinoma of the breast with widespread visceral and skeletal metastases treated with large doses of testosterone propionate. The effects of hormones in cases of advanced cancer of the breast are being investigated at Guy's Hospital. As a general rule, young patients with skeletal deposits are given androgens, and old patients with soft tissue deposits are given oestrogens.

At the Christie Hospital and Holt Radium Institute nitrogen mustard continues to be used in the treatment of Hodgkin's disease, particularly when the disease is so widespread that x-ray therapy would probably fail to palliate the symptoms much. In apparently very early cases when the disease is localized to one region nitrogen mustard is followed up by x-ray therapy. The Courtauld Institute of Biochemistry continues to investigate the effects of terofterin on tumour growth. Some degree of inhibition has been obtained in animal experiments with α -methopterin, a folic acid antagonist.

Cancer of the Lung

Analysis of samples of air for benzpyrene is being carried out at St. Bartholomew's Hospital, and it is hoped to improve the technique of sampling. No evidence has yet been found of the formation of benzpyrene by the combustion of tobacco. Experiments are also being carried out to determine the amount of arsenic in tobacco, and how much of it is inhaled. A wide variation in different brands of tobacco has been found. The most likely source of the arsenic seems to be the insecticides with which the plants are sprayed.

THE POPULAR LECTURE EPIDEMICS, OLD AND NEW

The final event of the Annual Meeting at Liverpool was the Popular Lecture, which was delivered in the evening of July 21, in the Picton Hall, to an audience of about 250. The President, Sir Henry Cohen, was in the chair. The lecturer was Dr. C. O. Stallybrass, and his subject was "Epidemics, Old and New." The lecture was illustrated by lantern-slides.

Dr. Stallybrass began by reminding his audience of some of the great pestilences of the past. Pestilence had often followed in the wake of military operations and had thwarted such operations. Again, the first attempt to pierce the Isthmus of Panama was defeated, not by engineering difficulties, but by malaria and yellow fever, and not until the studies of Laveran and of Manson and Ross, who laid bare the part played by mosquitoes in the spread of malaria, and of Reed, Carroll, and Lazear (the last of whom died in these researches), who showed that yellow fever also was conveyed only by certain mosquitoes, did the piercing of the isthmus become possible and the Tropics become habitable by white peoples. Men of all nations had combined in the unending warfare against disease, and thanks to them the average span of life had been lengthened by tens of years.

In the combat against a particular epidemic disease account must be taken of its individual characteristics. Without such knowledge campaign after campaign had failed. The lecturer took as an example typhus fever, the scourge of every sustained military campaign for 500 years. In 1528 the French army besieging Naples was turned back by typhus, 30,000 soldiers being struck down. Spanish soldiers conveyed it to Mexico, with devastating effects. During the Thirty Years' War (1618-1648) typhus had its widest diffusion, Germany suffering the most. Russia and Poland for long harboured typhus. Between 1816 and 1819 some 700,000 cases of typhus were said to have occurred in Ireland. In Liverpool, when Duncan, the first medical officer of health, was appointed 103 years ago, typhus was rampant. Duncan and his successors waged a somewhat unsuccessful war against typhus for 70 years, for they were fighting in the dark. The last indigenous cases of typhus in Liverpool were in 1918, but the lecturer saw a further small group of cases in Birkenhead in 1921, the infection having been derived from Glasgow. But by this time the method by which typhus was conveyed by lice was known, and the germ, the *Rickettsia prowazekii*, named after its two independent discoverers, Ricketts, an American, and von Prowazek, an Austrian, had been found. Both these men died of typhus in the course of their studies.

Serbia, after having turned back the Austrian army in 1915, was overcome by typhus—a more powerful foe—and 150,000 fell ill in six months. General Hunter, of the British Army, was seconded to Serbia, and in a few months' scientific campaign he had eradicated the disease, so much so that when in 1916 the lecturer was in charge of the beds for infectious disease in the Serbian Army he never saw a case. Even greater success attended the efforts of the United Nations in the second world war, when typhus broke out in Naples but was rapidly controlled. There was a threatening situation at the end of the war when the inmates of concentration and prisoner-of-war camps were released, but by that time D.D.T. and other chemicals to destroy