washings. Henle (quoted by Finn *et al.*, 1949), in complement-fixation tests for the viruses of influenza and choriomeningitis, had negative results.

The absence of any other infective agent and the nature of the outbreaks suggest a virus as the probable cause. In the present series complement-fixation tests for the Coxsackie 2 strain of virus were kindly performed by Dr. G. M. Findlay in March, 1950 (five months after the last recorded case). Sera from five cases were tested, and the results are listed in Table IV. Three cases were strongly positive and two faintly positive; two normal students taken as controls were negative.

TABLE IV.—Result of Complement-fixation Tests

Serum	Antigen		
	Coxsackie 1	Coxsackie 2	Normal Mouse
Case 4	—	+ 1 in 5	—
Case 7	—	+ 1 in 20	—
Case 9	—	+ 1 in 20	—
Case 10	—	+ 1 in 20	—
Case 11	—	+ 1 in 5	—
Control I	—	—	—
Control II	—	—	—

This virus, which is transmissible only through newborn mice, was originally isolated by Dalldorf and Sickles (1948) and later by Melnick *et al.* (1949) from the faeces of cases of "non-paralytic poliomyelitis" in the United States. In mice the infection involves skeletal muscles, producing severe myositis and paralysis, while the central nervous system is unaffected. Death commonly occurs within twenty-four hours of the onset of paralysis.

Transmission

The exact mode of transmission is unknown. Melnick *et al.* (1949) isolated the Coxsackie strains from sewage and from flies in areas where epidemics of "non-paralytic poliomyelitis" occurred.

It is interesting that the first case in the present series was the cook for the whole of House A, in which seven further cases occurred, and that Case 2, the first in House B, assisted her parents in preparing food for this household. While this strongly suggests an alimentary spread at the onset of the epidemic, later cases may have arisen from droplet infection owing to the close proximity of their living quarters. On inquiry, however, no cases could be traced outside the two households among their day-to-day contacts at work.

Coincident with the epidemic, one sporadic case was also seen during the illness. A transitory bilateral pleural rub was present and the complement-fixation reaction to the Coxsackie 2 strain was strongly positive (1 in 20). This case also had a minimal unilateral pleuritic reaction on x-ray examination which did not persist.

Summary

An explosive outbreak of benign dry pleurisy is reported in two near-by households under common ownership. Two cases are described in detail.

The epidemiological aspects of the outbreak are discussed.

Five months after the illness the sera from five cases were shown by Dr. G. M. Findlay to give positive complement-fixation reactions to the Coxsackie 2 strain of virus as compared with negative results in two normal controls.

It is suggested that, in view of the seasonal incidence and the possible relationship to "non-paralytic poliomyelitis," the mode of spread may be similar to that of poliomyelitis itself and is possibly alimentary.

I wish to thank Dr. G. M. Findlay for his advice and for performing the serological tests, and Professor R. Cruickshank and Dr. T. A.

Kemp, physician in charge of the Student Health Service, St. Mary's Hospital, for their guidance and helpful criticism.

REFERENCES

- Attlee, W., Amsler, A. M., and Beaumont, D. C. (1924). *Lancet*, 2, 492.
 Carter, A. H. (1933). *British Medical Journal*, 2, 1186.
 Dalldorf, G., and Sickles, G. M. (1948). *Science*, 108, 61.
 Finn, J. J. (1947). *New Engl. J. Med.*, 237, 621.
 —, Weller, T. H., and Morgan, H. R. (1949). *Arch. intern. Med.*, 83, 305.
 Howard, C. R. G. (1938). *British Medical Journal*, 2, 1203.
 Jamieson, W. M., and Prinsley, D. M. (1947). *Ibid.*, 2, 47.
 Lloyd, E. I. (1924). *Lancet*, 2, 272.
 McConnell, J. (1945). *Amer. J. med. Sci.*, 209, 41.
 Melnick, J. L., Shaw, E. W., and Curnen, E. C. (1949). *Proc. Soc. exp. Biol., N.Y.*, 71, 344.
 Pickles, W. N. (1933). *British Medical Journal*, 2, 817.
 — (1939). *Epidemiology in Country Practice*. Wright, Bristol.
 — (1948). *Lancet*, 1, 201.
 Scadding, J. G. (1946). *Ibid.*, 1, 763.
 Small, J. C. (1924). *Amer. J. med. Sci.*, 168, 570.
 Sylvest, E. (1933). *Den Bornholmske Syge: Myalgia Epidemica*. Levin and Munksgaard, Copenhagen. English translation, *Epidemic Myalgia*, 1934. London.
 Williamson, B. (1924). *Lancet*, 2, 64.

ADRENAL INSUFFICIENCY AFTER SURAMIN TREATMENT OF PEMPHIGUS

BY

L. J. MAHONEY, B.A., M.D.

AND

H. J. BARRIE, B.M., B.Ch.

(From the Department of Pathology, University of Toronto, Toronto, Canada)

[WITH SPECIAL PLATE]

Suramin ("germanin") has been used for the treatment of pemphigus since 1931. It is a carbamide of a complex salt of trisulphonic acid, is also known as "naphuride" and "Bayer 205," and was originally introduced for the treatment of trypanosomiasis. It is known to have toxic effects on the renal tubules and haemopoietic system. Wells *et al.* (1937) have shown that when given to guinea-pigs in amounts comparable with human doses it causes damage to the renal tubules and adrenal cortex, but the full picture of so-called adrenal atrophy was not achieved in their experiments. Two cases of acute adrenal insufficiency following suramin treatment have been reported (Wells *et al.*, 1937; Tomlinson and Cameron, 1938), but any conclusion that these adrenal changes were solely due to the drug is confused by the fact that many of the features of pemphigus, especially in acute cases, strongly resemble those of adrenal insufficiency and are sometimes improved by the administration of cortical extracts.

Furthermore, Goldzieher (1945) has shown that signs of severe organic damage may be found in the adrenal cortex in deaths from pemphigus. None of the six cases he studied had received suramin,* and he regarded their lesions as terminal secondary processes such as occur in other severe debilitating diseases.

The paradox is a familiar pharmaceutical one—that a drug which mimics the effects of a disease should be beneficial for it—and perhaps the problem is one of correct dosage for the individual.

Wells *et al.* state that none of the persons who have been followed up for some time after suramin therapy for trypanosomiasis have shown any delayed symptoms attributable to adrenal insufficiency. We are therefore in somewhat

*Personal communication.

the same position as we once were with agranulocytosis and sulphonamide therapy. From consideration of only a few cases it is hard to be certain whether the disease or the drug plays the dominant part in producing the lesion. It seems to us likely that in the case of suramin it is a summation effect. We think that this further experience with suramin should therefore be reported. No further comment is necessary.

Case Report

A 37-year-old woman of Scottish-Canadian ancestry was perfectly well until seven months before death, when, in the second month of pregnancy, she developed typical lesions of pemphigus. During the next four months two courses of suramin were started but were stopped after one injection (0.25 g.) because of the development of fever. Three months before death she was delivered of a normal seven-months male infant by elective caesarean section and two weeks later developed rapid mood swings, hallucinations, and delusions, and was admitted to hospital.

Examination revealed a dehydrated, confused, belligerent, agitated, and hallucinated woman whose skin was covered with lesions in all stages of healing, varying from patches of erythema to bullae measuring 2 in. (5 cm.) in diameter. Her temperature varied from 99 to 100° F. (37.2 to 37.8° C.). Her blood pressure was 108/60. The haemoglobin was 70% and the leucocytes numbered 16,000 per c.mm. The urine contained many white cells but no albumin, and cultured *Bact. coli*. Her N.P.N. was 27 mg. per 100 ml.

She was treated with electric shock, and a third course of suramin was completed five weeks before death (1 g. every two days for ten doses), with definite improvement in her mental condition and her skin lesions. No albumin appeared in the urine immediately after the course of suramin. Because of a falling haemoglobin 500 ml. of whole blood was given. At this time she developed an upper respiratory infection which persisted. Two weeks before death abnormal pigmentation of her skin and buccal mucous membrane was noticed, and over a period of a few days she developed anorexia, nausea, and vomiting, and her temperature rose suddenly to 104° F. (40° C.), her pulse to 130, her respirations to 30, and her blood pressure fell to 60/46. Her urine now contained 1+ albumin and her blood chlorides were 489 mg. per 100 ml.

Because of the signs of adrenal insufficiency 10 ml. of "cortin" was given every four hours, at first intravenously and then intramuscularly, together with a continuous intravenous infusion of 5% glucose in normal saline and 100,000 units of penicillin intramuscularly twice a day. The nausea and vomiting decreased, her blood pressure remained around 90/50, and her temperature varied from 99 to 100° F. with occasional elevations up to 102° F. (38.9° C.). Her serum chloride level remained around 540 mg. per 100 ml.

Five days before death her temperature rose to 105° F. (40.6° C.) and then fell to normal, and, though her blood pressure was 100/50 and she appeared to be getting no worse, she died suddenly in the middle of the night.

Necropsy Findings

At necropsy typical Addisonian pigmentation of the buccal mucous membrane and skin was found, and large branny brown macules were scattered over the trunk and extremities.

The heart was soft and flabby; the spleen (470 g.) and liver (2,200 g.) were enlarged and both were congested. Several of the abdominal lymph nodes were enlarged and firm. The adrenals were thin, weighed 10 g. combined, and on cut section all laminae were narrow. The lipid zone was even and contained lipid which varied slightly in brightness. It was separated from the grey medulla by a vivid, sharply demarcated reddish black stripe which was much darker than the usual pigment zone.

There was moderate oedema of the brain (1,420 g.), with some herniation of the hippocampal unci and cerebellar tonsils, and moderate compression of the ventricles. All other organs appeared normal.

Microscopical Examination

Fragmentation of the myocardium, congestion of the liver and spleen, and oedema of the abdominal lymph nodes were found, but nothing else of importance except in the adrenals and thyroid.

Adrenals.—Sections of each adrenal were alike. The capsule and its blood vessels appeared normal. The most obvious change was a band of extreme congestion and haemorrhage occupying the inner half of the cortex (Plate, Fig. 1). In it could be seen isolated ghosts of pigmented cells. Peripherally there were radial bars of dilated sinusoids traversing the rest of the cortex, but the capillaries of the medulla were only slightly dilated and then only at the line of junction with the cortex. Because of the variable arrangement of the laminae in the centre of the adrenal it was difficult to be certain that none of the medullary cells were necrotic; most of them showed no evidence of it but were rather widely separated by oedema. In the outer half of the cortex the sinusoids could be recognized, and the spaces between were occupied by vacuoles separated by thin wisps of cytoplasm. Sometimes one could be certain that these vacuoles were in cells and sometimes it was difficult to be sure. At rare intervals there were cells without lipid occurring singly and in clumps, and many of these showed an acute autolytic form of necrosis and the adjoining sinusoids were enormously dilated (Fig. 2). In frozen sections the vacuoles were found to contain abundant lipid, most of which was doubly refractile. In summary, there was complete necrosis of the zona reticularis, and vacuolar degeneration and patchy necrosis of the remainder of the cortex.

Thyroid.—The thyroid acini were small, lined by flattened epithelium, and contained little but rather darkly staining colloid. There were many bands in which the acini had atrophied and had been replaced by lymphoid tissue.

Summary

A case is reported of a woman of 37 who developed pemphigus in pregnancy and died of acute adrenal necrosis five weeks after finishing a course of suramin therapy.

We would like to thank Dr. W. Garbe for permission to publish this case.

REFERENCES

- Goldzieher, J. W. (1945). *Arch. Derm. Syph., Chicago*, **52**, 369.
Tomlinson, C. C., and Cameron, O. J. (1938). *Ibid.*, **38**, 555.
Wells, H. G., et al. (1937). *J. Amer. med. Ass.*, **109**, 490.

A professional conference organized by the Royal College of Nursing and attended by some hundreds of nurses was held in London on June 28 to discuss the educational scope and potentialities of the Nurses Act, 1949. Miss Ethel Strudwick presided, and the principal address was given by Miss Enid Russell Smith, Under Secretary, Ministry of Health. Miss Russell Smith said that the Act provided for the first time for the financing of nurse training separately from the finances of the hospital. It ensured that the money for such training would not be diverted to any other object. The General Nursing Council had been reconstituted and strengthened, educationally and financially, to enable it to carry out its new function. There had previously been a two-tier system, consisting of the General Nursing Council and some 600 training schools: this was bound to be inflexible. Under the Act there would be an intermediate tier of 14 area management committees, each with responsibility for about 40 training schools. Miss Joan B. Price, sister-tutor at the Sheffield Hospitals School of Nursing, said that this school was an educational establishment for four hospitals with 1,200 beds, meeting the educational needs of 500 to 600 nurses. It was financially and administratively independent of the group of hospitals with which it was associated. The teaching staff were appointed by the school itself. There was a most valuable connexion with the medical faculty of the university. Student nurses attended the university and used the lecture theatre, and university lecturers visited the school.

GEORGE LUMB: DIFFUSE EOSINOPHILIC ARTERITIS

FIG. 1.—Radiograph of chest taken on June 20, 1949.

FIG. 2.—Lung. Fibrosis, giant-cell formation, and eosinophilic infiltration.

FIG. 3.—Lung. Necrosis and eosinophilic infiltration of the wall of a vessel. ($\times 150$.)

FIG. 5.—Kidney. Necrosis of convoluted tubules, capillary thrombosis in a glomerulus, and eosinophilic infiltration of the interstitial tissue. ($\times 150$.)

FIG. 4.—Heart. High-power view of eosinophils. ($\times 240$.)

L. J. MAHONEY AND H. J. BARRIE: ADRENAL INSUFFICIENCY AFTER SURAMIN

FIG. 1.—Adrenal. Vacuolation of zona fasciculata and haemorrhagic necrosis of zona reticularis. H. and E. ($\times 45$.)

FIG. 2.—Adrenal. Vacuolar degeneration of zona fasciculata with isolated islands of autolysing cells. H. and E. ($\times 115$.)